

# Absence of absolutely continuous spectrum of Floquet operators

Alain Joye  
Centre de Physique Théorique  
C.N.R.S. Marseille  
Luminy Case 907  
13288 Marseille Cedex 9  
and  
PHYMAT  
Université de Toulon et du Var  
B.P. 132  
83957 La Garde Cedex

## Abstract

The spectrum of the Floquet operator associated with time periodic perturbations of discrete hamiltonians is considered. If the gap between successive eigenvalues  $\lambda_j$  of the unperturbed hamiltonian grows as  $\lambda_j - \lambda_{j-1} \simeq j^\alpha$  and the multiplicity of  $\lambda_j$  grows as  $j^\beta$  with  $\alpha > \beta \geq 0$  as  $j$  tends to infinity, then the corresponding Floquet operator possesses no absolutely continuous spectrum provided the perturbation is smooth enough.

CPT-93/P.2957  
October 1993  
K.W.

# 1 Introduction

Consider a positive self-adjoint hamiltonian  $H_0$  on a separable Hilbert space  $\mathcal{H}$  with discrete spectrum  $\{\lambda_j\}_{j=1,\dots,\infty}$  and  $W(t)$  a symmetric time dependent periodic perturbation

$$W(t + 2\pi) = W(t), \quad \forall t \in \mathbf{R}. \quad (1.1)$$

The associated Floquet operator, defined by

$$F = -i \frac{\partial}{\partial t} + H_0 + W(t) \quad (1.2)$$

on  $L^2[0, 2\pi] \otimes \mathcal{H}$  with periodic boundary conditions in  $t$ , has been the object of considerable interest recently. More precisely, the nature of the spectrum of  $F$ ,  $\sigma(F)$ , has been investigated thoroughly for specific models and for more general situations as well, as reviewed by Bellissard [B] and more recently by Jauslin [J]. The interest of such detailed studies for physics lies in the fact that the long time behaviour of the solutions  $\psi(t)$  of the time dependent Schrödinger equation

$$i \frac{\partial}{\partial t} \psi(t) = (H_0 + W(t))\psi(t), \quad \psi(0) = \varphi \quad (1.3)$$

is closely related to the spectral properties of the Floquet operator. This asymptotic behaviour of the solutions of (1.3) is quite relevant for the study of quantum stability or quantum chaos [B], [J]. Let  $U(t)$  be the unitary evolution associated with (1.3) such that

$$\psi(t) = U(t)\varphi. \quad (1.4)$$

As a consequence of the periodicity of the hamiltonian  $H_0 + W(t)$ , the solution of (1.3) satisfies

$$\psi(n2\pi) = U(2\pi)^n \varphi, \quad \forall n \in \mathbf{N} \quad (1.5)$$

where  $U(2\pi)$  is the monodromy operator. The large  $n$  behaviour of  $\psi(n2\pi)$  is thus clearly dependent on the spectral subspace (pure point, absolutely or singular continuous) of  $U(2\pi)$  in which the initial condition  $\varphi$  is chosen, as discussed in [B] and [J]. On the other hand, the spectral properties of the Floquet operator  $F$  and of the monodromy operator  $U(2\pi)$  are equivalent, as established in [Y], [H1]. Note that this type of approach can be generalized in order to study the stability of the quantum dynamics when the perturbation  $W(t)$  is quasiperiodic in time, as demonstrated recently in [BJL].

From the mathematical point of view, these considerations pertain to the perturbation theory of operators with dense pure point spectrum. Indeed, if  $W(t) \equiv 0$ , the spectrum of the Floquet operator is given by  $\{n + \lambda_j\}_{j=1,\dots,\infty}^{n \in \mathbf{N}}$  which is generically dense on the real line. Correspondingly, the spectrum of the associated monodromy operator  $U_0(2\pi)$  consists in the set  $\{e^{-2\pi i \lambda_j}\}_{j=1,\dots,\infty}$  which generically fills the unit circle densely. The question is the following: does this pure point spectrum remain stable after perturbation by the time periodic operator  $W(t)$ ? Although this problem is in general rather delicate [H2], a rigorous complete positive answer can be given for certain specific models, in some range of parameters. For example, the pulsed rotor considered by Bellissard in [B], some time dependent quadratic hamiltonians studied by Hagedorn, Loss and Slawny [HLS], a class of time dependent perturbations of the harmonic oscillator and discrete hamiltonians kicked periodically by some rank one perturbations as shown by Combescure in [C1] and [C2]. On the other hand, results valid for general systems are scarce and they provide a partial answer only to the above question. These results are based on the search of general criteria allowing to exclude the presence of

absolutely continuous spectrum in the Floquet operators. This approach was initiated and developed by Howland in the series of papers [H3], [H4], [H5]. Such results yield a partial answer to our question in the sense that they say nothing about singular continuous spectrum. However, the absence of absolutely continuous component  $\sigma_{ac}(F)$  in the spectrum of  $F$  is sometimes already considered as stability result. The criteria obtained are of the following form. Assume the hamiltonian  $H_0$  has simple eigenvalues  $\lambda_1 < \lambda_2 < \dots < \lambda_j < \dots$  satisfying the growth condition

$$\lambda_j - \lambda_{j-1} \simeq j^\alpha \tag{1.6}$$

for some  $\alpha > 0$  and suppose that  $W(t)$  is uniformly bounded. Then it is proven in [H4] that  $\sigma_{ac}(F) = \emptyset$  for any  $\alpha > 0$  provided  $W(t)$  is smooth enough. Actually, this result is also true for some class of unbounded perturbations  $W(t)$ , as discussed in [H5], and it is even shown in [H3] that  $\sigma(F)$  is generically dense pure point, in an appropriate probabilistic sense, provided  $\alpha > 2$ . However, the restriction imposed on the eigenvalues of  $H_0$  to be non degenerate was conjectured in [H4] to be technical only. Indeed, the result is expected to hold for the pulsed rotor of Bellissard, a case where the corresponding eigenvalues satisfy (1.6) but are doubly degenerate. Accordingly, it was proven recently by Nenciu [N1] that  $\sigma_{ac}(F) = \emptyset$  if the growth condition (1.6) is satisfied, and if the multiplicity  $m_j$  of the eigenvalues  $\lambda_j$  is uniformly bounded in  $j$ , provided  $W(t)$  is smooth enough. However, to prove this result, it was necessary to impose another technical condition, namely  $\alpha > 1/2$ .

In this paper we reconsider the absence of absolutely continuous spectrum of Floquet operators in the generalization due to Nenciu [N1] of the framework designed by Howland [H3], [H4], [H5] (see below). We improve the preceding results in two ways. First, we remove the technical condition  $\alpha > 1/2$  to give a complete proof of the conjecture of Howland [H4] in the degenerate case for any positive  $\alpha$ . Second, and more important, we can allow the degeneracy  $m_j$  of the eigenvalues  $\lambda_j$  to increase with  $j$  as  $m_j \simeq j^\beta$ , with  $\beta < \alpha$ . This generalization may be of interest for the study of systems of more than one degree of freedom since in such cases the degeneracy is likely to increase with the principal quantum number. See however the remarks following the main theorem 2.1.

## 2 Result and strategy

Let  $H_0$  be an operator satisfying the following spectral hypothesis **S**:

- $H_0$  is densely defined on  $D \subset \mathcal{H}$ , self-adjoint and positive
- $\sigma(H_0) = \bigcup_{j=1}^{\infty} \sigma_j$
- $\sigma_j$  consists in a finite number of finitely degenerate eigenvalues such that
- $\max_{\lambda, \mu \in \sigma_j} |\lambda - \mu| \leq c_0$
- $\text{dist}(\sigma_j, \sigma_{j-1}) \geq c_1 j^\alpha$ ,  $\alpha > 0$
- $\text{mult}(\sigma_j) \leq c_2 j^\beta$ ,  $\beta \geq 0$

where  $c_0, c_1, c_2, \alpha, \beta$  are independent of  $j$ .

Let  $W(t)$  be an operator satisfying the regularity condition **R<sub>k</sub>**:

- $W(t)$  is bounded and symmetric  $\forall t \in \mathbf{R}$
- $W(t)$  is strongly  $C^k$   $\forall t \in \mathbf{R}$ .

The operator  $H_0 + W(t)$  is thus self-adjoint, densely defined on  $D$  (thm 4.3, p.287 [Ka]) and strongly  $C^k$ , with bounded derivatives. If  $k \geq 1$ , there exists a unitary evolution operator  $U(t)$ , strongly  $C^1$  on  $D$ , which maps  $D$  into  $D$  and satisfies for any  $\varphi$  in  $D$  and  $t \in \mathbf{R}$

$$i \frac{\partial}{\partial t} U(t) \varphi = (H_0 + W(t)) U(t) \varphi, \quad U(0) = \mathbf{I}, \quad (2.1)$$

as can be deduced from theorem X70 [RS].

**Theorem 2.1** *Let  $H_0$  satisfy  $S$  and  $W(t)$  be  $2\pi$ -periodic in  $t$  and satisfy  $R_k$ , with  $k \geq 1$ .*

*If  $\alpha > \beta \geq 0$  and  $k \geq \left\lceil \frac{1+\beta}{\alpha} \right\rceil + 1$ , then  $\sigma_{ac}(U(2\pi)) = \emptyset$ .*

**Remarks:**

Setting  $\beta = 0$ , we obtain the conjecture of Howland [H4] for any  $\alpha > 0$ .

It is possible to weaken the spectral hypothesis  $S$  somehow since the result also holds if the size of the spectral sets  $\sigma_j$  grows as

$$\max_{\lambda, \mu \in \sigma_j} |\lambda - \mu| \leq c_0 j^\alpha. \quad (2.2)$$

As already noticed, the above theorem describes  $\sigma(U(2\pi))$  only partially. However, the result is not of perturbative nature, in the sense that the norm of the operator  $W(t)$  can be arbitrarily large. By contrast, the complete characterization of  $\sigma(F)$  performed by Bellissard [B] and Combescure [C1] on their models can be achieved in regimes where  $\|W(t)\|$  is small enough.

This result can also be useful for some cases where  $W(t)$  is unbounded. Indeed, it is shown in [H5] how to reduce the study of the Floquet operator  $-i \frac{\partial}{\partial t} + H_0 + W_u(t)$ , where  $W_u(t)$  belongs to a certain class of unbounded operators, to the study of  $-i \frac{\partial}{\partial t} + H_0 + W_b(t)$ , where  $W_b(t)$  is bounded. And, according to the final remark of §2 in [H5], a similar procedure can be applied when  $H_0$  has degenerate eigenvalues.

As noted in [J], this type of result is likely to apply to one degree of freedom systems essentially, because of the growth condition on the gaps between successive eigenvalues in hypothesis  $S$ . This impression is strengthened by the supplementary condition imposed on the growth of the multiplicity of eigenvalues. Indeed, consider the simple two degrees of freedom system given by a free rotator in  $\mathbf{R}^3$ . The hamiltonian of the system is  $H_0 = \mathbf{J}^2$ , where  $\mathbf{J}$  denotes the angular momentum operator. The gaps between the eigenvalues  $\lambda_j = j(j+1)$  of  $H_0$  do satisfy the growth condition with exponent  $\alpha = 1$ ,

$$j(j+1) - (j-1)j = 2j, \quad (2.3)$$

however the multiplicity of these eigenvalues grows as

$$\text{mult}(\lambda_j) = 2j + 1 \quad (2.4)$$

so that the exponent  $\beta = 1$  and the condition  $\alpha > \beta$  is not satisfied. This is nevertheless in agreement with the fact that multidimensional systems seem to be more inclined to instabilities [J]. Note that if we consider formally  $H_0 = \mathbf{J}^4$ , the corresponding eigenvalues  $\lambda_j = j^2(j+1)^2$  and their multiplicities behave in the proper way with exponents  $\alpha = 3$  and  $\beta = 1$ .

The strategy followed to prove results of this type, which is common to [H4], [N1] and the present work, is based on the few general theorems of stability of absolutely continuous spectrum. In [H4] where the Floquet operator  $F$  is considered, Howland shows by means of

a KAM inspired procedure that  $F$  is unitarily equivalent to an operator  $F_0 + R$ , where  $F_0$  is self-adjoint and has a pure point spectrum and  $R$  is trace class. Then it remains to invoke the Kato-Rosenblum theorem (thm. 4.4 p.540 [Ka]) stating that the absolutely continuous subspaces of self-adjoint operators are unitarily equivalent if they differ by a trace class operator. In [N1], Nenciu deals with the monodromy operator  $U(2\pi)$  instead of  $F$ , and uses recently developed tools in the adiabatic theory [N2] to approximate  $U(2\pi)$  by  $V + R$  where  $V$  is a unitary operator having pure point spectrum and  $R$  is trace class. The result is thus achieved by virtue of the Birman-Krein theorem [BK], an equivalent of the Rosenblum-Kato theorem for unitary operators, which assesses that the absolutely continuous spectral subspaces of unitary operators are unitarily equivalent if they differ by a trace class operator. Our proof of theorem 2.1 above follows the latter method proposed by Nenciu, however we make use of another adiabatic approximation technique developed in [JP1],[JP2]. In the present context, where no small parameter appears, this method proves to be very efficient as well. Indeed, it's relative simplicity makes it possible for us to obtain the accurate estimates which are needed to bound operators in the trace norm and to extend the previous results as described in theorem 2.1.

**Acknowledgments:** It is a pleasure to thank P.Briet, J.-M.Combes, P.Duclos, and J.Howland for many informative discussions at various stages of this work.

## 3 Proof

We present in this section the proof of theorem 2.1 based on the adiabatic techniques developed in [JP2]. In doing so, we make use of some intermediate results which will be proven in the next technical section.

### 3.1 Preliminaries

Let us first consider the stability of the spectral hypotheses  $S$ .

**Lemma 3.1** *Let  $H_0$  satisfy  $S$  and let  $B(t)$  satisfy  $R_0$ . Then  $H_0 + B(t)$  satisfies  $S$  uniformly in  $t \in [0, 2\pi]$  with the same exponents  $\alpha$  and  $\beta$ .*

**Proof:** The uniform boundedness of  $B(t)$  and the growth condition on the gaps ensure that  $\sigma(H_0 + B(t))$  consists in the disjoint union of new sets  $\bigcup_{k=1}^{\infty} \sigma'_k(t)$ . Hence, taking into account a possible relabelling, the gaps between these sets behave as

$$\text{dist}(\sigma'_k(t), \sigma'_{k-1}(t)) \geq c_1(k+r)^\alpha \geq c'_1 k^\alpha \quad (3.1)$$

for large  $k$ 's. Moreover, it is readily seen by considering the interpolating operator  $H_0 + xB(t)$ , where  $x$  ranges in  $[0, 1]$ , that

$$\text{mult}(\sigma'_k(t)) = \text{mult}(\sigma_{k+r}) \leq c'_2 k^\beta, \quad (3.2)$$

for  $k$  large enough. □

**Notation:** There will appear several constants in the sequel which we shall denote generically by the same symbol  $c$ . From now on, the time derivative will be denoted by a prime.

Consider the operator  $H_0 + B(t)$  where  $H_0$  satisfies  $S$  and  $B(t)$  satisfies  $R_0$ . Let  $R(t, \lambda) = (H_0 + B(t) - \lambda)^{-1}$ . By the above lemma, for  $t \in [0, 2\pi]$ , the spectrum  $\sigma(t)$  of  $H_0 + B(t)$  consists in spectral sets  $\sigma_j(t)$  and the associated spectral projectors  $P_j(t)$  can be written as

$$P_j(t) = -\frac{1}{2\pi i} \oint_{\Gamma_j} R(t, \lambda) d\lambda \quad (3.3)$$

where the non-intersecting paths  $\Gamma_j$  surrounding  $\sigma_j(t)$  are chosen in such a way that

$$\text{long}(\Gamma_j) \equiv |\Gamma_j| \leq c j^\alpha \quad (3.4)$$

and

$$\text{dist}(\Gamma_j, \sigma(t)) \geq c j^\alpha. \quad (3.5)$$

**Remark:** Both estimates (3.4) and (3.5) are true for appropriate paths  $\Gamma_j$  if the length of the sets  $\sigma_j(t)$  grows as  $j^\alpha$ , in the spirit of the first remark below theorem 2.1.

**Proposition 3.1** *Let  $H_0$  satisfy  $S$ ,  $B(t)$  satisfy  $R_n$ ,  $n \geq 1$ , and  $P_j(t)$  be defined by (3.3). Then, if  $\alpha > \beta \geq 0$  the operator*

$$K(t) = \sum_{j=1}^{\infty} P_j(t) P_j'(t)$$

*is bounded, strongly  $C^{n-1}$ ,  $n \geq 1$ , on  $[0, 2\pi]$  and such that  $K(t)^* = -K(t)$ .*

**Remark:** The content of this proposition is nontrivial for  $\alpha \leq 1/2$ , see [N1].

Actually, the proposition is a consequence of the following technical

**Lemma 3.2** *Assume  $H_0$  satisfies  $S$ ,  $B(t)$  satisfies  $R_0$  and let*

$$G(t) = \sum_{j=1}^{\infty} P_j(t) \frac{1}{2\pi i} \oint_{\Gamma_j} A_j(t, \lambda) R(t, \lambda) d\lambda$$

*where*

$$\sup_{t \in [0, 2\pi]} \sup_{\lambda \in \Gamma_j} \|A_j(t, \lambda)\| \leq \frac{c}{j^\alpha}.$$

*Then, if  $\alpha > \beta \geq 0$ ,  $G(t)$  is bounded and strongly continuous  $\forall t \in [0, 2\pi]$ . If, furthermore,  $B(t)$  satisfies  $R_1$  and  $A_j(t, \lambda)$  is strongly  $C^1$  with*

$$\sup_{t \in [0, 2\pi]} \sup_{\lambda \in \Gamma_j} \|A_j'(t, \lambda)\| \leq \frac{c}{j^\alpha},$$

*then,  $G(t)$  is strongly  $C^1 \forall t \in [0, 2\pi]$  and*

$$G'(t) = G_1(t) - G_0(t)G(t)$$

*where  $G_0(t)$  and  $G_1(t)$  have the same form as  $G(t)$  with*

$$A_j^0(t, \lambda) = R(t, \lambda) B'(t)$$

*and*

$$A_j^1(t, \lambda) = P_j'(t) A_j(t, \lambda) + A_j'(t, \lambda) - A_j(t, \lambda) R(t, \lambda) B'(t)$$

*in place of  $A_j(t, \lambda)$ .*

**Remark:** The operator  $G_0(t)$  coincides with the operator  $K(t)$  of the proposition since

$$R'(t, \lambda) = -R(t, \lambda)B'(t)R(t, \lambda). \quad (3.6)$$

**Proof of proposition 3.1:** Since  $H_0 + B(t)$  satisfies  $S$  uniformly in  $t \in [0, 2\pi]$ , we have by (3.5)

$$\|R(t, \lambda)|_{\lambda \in \Gamma_j}\| \leq \frac{c}{j^\alpha} \quad (3.7)$$

which yields the required bound on  $A_j^0(t, \lambda)$ . Hence  $K(t)$  is bounded provided  $\alpha > \beta \geq 0$ . Now, if  $B(t)$  is strongly  $C^n$ , the same is true for  $R(t, \lambda)$ , see (3.6), and using the Leibnitz formula or eq.(2.36) in [JP2] we get

$$\left\| \left( \frac{\partial}{\partial t} \right)^m R(t, \lambda)|_{\lambda \in \Gamma_j} \right\| \leq \frac{c}{j^\alpha}, \quad \forall m \leq n. \quad (3.8)$$

Consequently,  $A_j^0(t, \lambda)$  is strongly  $C^{n-1}$  and

$$\left\| \left( \frac{\partial}{\partial t} \right)^m A_j^0(t, \lambda) \right\| \leq \frac{c}{j^\alpha} \quad (3.9)$$

uniformly in  $\lambda \in \Gamma_j$  and  $t \in [0, 2\pi]$ , for all  $m \leq n - 1$ . Thus, as easily checked, the formula of lemma 3.2 can be iterated since, using (3.4), (3.6) and (3.8),

$$\left\| \left( \frac{\partial}{\partial t} \right)^m P_j(t) \right\| \leq \frac{c}{j^\alpha}, \quad \forall 1 \leq m \leq n. \quad (3.10)$$

The identity  $K(t)^* = -K(t)$  results from the self adjointness of the projectors  $P_j(t)$  and the identity

$$0 = \sum_{j=0}^{\infty} P_j'(t) = \sum_{j=0}^{\infty} (P_j'(t)P_j(t) + P_j(t)P_j'(t)). \quad (3.11)$$

□

## 3.2 Adiabatic formalism

We introduce in this section an iteration scheme which, in the adiabatic context, is also known as superadiabatic renormalization [JP1], [JP2]. For  $t \in [0, 2\pi]$ , we start with

$$H_0(t) \equiv H_0 + W(t) \quad (3.12)$$

where  $H_0$  satisfies  $S$  and  $W(t)$  satisfies  $R_k$ ,  $k \geq 1$ , so that lemma 3.1 applies. The spectral projectors of  $H_0(t)$  are denoted by

$$P_j^0(t) = -\frac{1}{2\pi i} \oint_{\Gamma_j^0} R_0(t, \lambda) d\lambda \quad (3.13)$$

where  $R_0(t, \lambda) = (H_0(t) - \lambda)^{-1}$  and  $\Gamma_j^0$  encircles the spectral set  $\sigma_j^0(t)$  in such a way that (3.4) and (3.5) hold (with the obvious change of notation). We define the operator

$$K_0(t) = \sum_{j=1}^{\infty} P_j^0(t)P_j^{0'}(t) \quad (3.14)$$

which is bounded and strongly  $C^{k-1}$  by proposition 3.1. At the  $q^{\text{th}}$  step,  $k-1 \geq q \geq 1$ , we set

$$H_q(t) \equiv H_0(t) + iK_{q-1}(t) \quad (3.15)$$

which satisfies  $S$  as well. Thus we can define its spectral projectors by

$$P_j^q(t) = -\frac{1}{2\pi i} \oint_{\Gamma_j^q} R_q(t, \lambda) d\lambda \quad (3.16)$$

where  $R_q(t, \lambda) = (H_q(t) - \lambda)^{-1}$  and  $\Gamma_j^q$  encircles the spectral set  $\sigma_j^q(t)$  in such a way that (3.4) and (3.5) hold. Similarly, we define

$$K_q(t) = \sum_{j=1}^{\infty} P_j^q(t) P_j^{q'}(t). \quad (3.17)$$

Using proposition 3.1 iteratively, we find that  $H_q(t)$  is strongly  $C^{k-q}$  whereas  $K_q(t)$  is strongly  $C^{k-q-1}$ , so that this scheme is well defined provided  $q \leq k-1$ .

Let  $V_q(t)$  be the solution of the Schrödinger-like equation for  $t \in [0, 2\pi]$

$$iV_q'(t) = (H_q(t) - iK_q(t))V_q(t), \quad V_q(0) = \mathbf{I}. \quad (3.18)$$

**Lemma 3.3** *For any  $q \leq k-1$ , the operator  $V_q(t)$  is unitary, maps  $D$  into  $D$  and satisfies (3.18) strongly on  $D$ . Moreover,*

$$V_q(t) P_j^q(0) = P_j^q(t) V_q(t)$$

for any  $t \in [0, 2\pi]$  and any  $j = 1, \dots, \infty$ .

**Remark:** The first part of the lemma is nontrivial for  $q = k-1$  since  $K_{k-1}(t)$  is not differentiable. The second part generalizes standard results [Ka], [Kr] which hold for a finite number of projectors.

**Corollary 3.1** *If  $W(t)$  is  $2\pi$ -periodic and  $q \leq k-1$ ,*

$$\sigma_{ac}(V_q(2\pi)) = \emptyset.$$

**Proof:** The operators  $H_q(t)$  are  $2\pi$ -periodic since their construction is local. Hence we have

$$P_j^q(2\pi) = P_j^q(0) \quad (3.19)$$

so that

$$[V_q(2\pi), P_j^q(0)] = 0, \quad \forall j = 1, \dots, \infty. \quad (3.20)$$

Since the orthogonal subspaces  $P_j^q(0)\mathcal{H}$  are finite dimensional,  $V_q(2\pi)$  has pure point spectrum.  $\square$

Let us now evaluate the difference between  $U(2\pi)$  and  $V_q(2\pi)$ . For  $\varphi \in D$  we compute, see (3.12),

$$\begin{aligned} i \left( V_q^{-1}(t) U(t) \varphi \right)' &= V_q^{-1}(t) (-H_q(t) + iK_q(t) + H_0(t)) U(t) \varphi \\ &= iV_q^{-1}(t) (K_q(t) - K_{q-1}(t)) U(t) \varphi \end{aligned} \quad (3.21)$$



hence

$$U(t) - V_q(t) = V_q(t) \int_0^t ds V_q^{-1}(s) (K_q(s) - K_{q-1}(s)) U(s). \quad (3.22)$$

In order to apply the Birman-Krein theorem [BK], it remains to show that the trace norm of  $U(2\pi) - V_q(2\pi)$  is finite. This will be true if we show that

$$\sup_{t \in [0, 2\pi]} \|K_q(t) - K_{q-1}(t)\|_1 \leq c, \quad (3.23)$$

where  $\|\cdot\|_1$  stands for the trace norm.

### 3.3 Estimations in the trace norm

We first note that

$$K_q(t) - K_{q-1}(t) = \sum_{j=1}^{\infty} \left( P_j^q(t) P_j^{q'}(t) - P_j^{q-1}(t) P_j^{q-1'}(t) \right) \quad (3.24)$$

where the operators  $P_j^q(t) P_j^{q'}(t) - P_j^{q-1}(t) P_j^{q-1'}(t)$  are degenerate. And since  $H_q(t)$  and  $H_{q-1}(t)$  satisfy the spectral hypothesis  $S$ , we have

$$\dim \text{Ran}(P_j^q(t) P_j^{q'}(t) - P_j^{q-1}(t) P_j^{q-1'}(t)) \leq c j^\beta. \quad (3.25)$$

Our main tools to perform estimations in the trace norm are the following lemmas to be found in [Ka], pp. 521.

#### Lemmas

- i) If  $T$  is degenerate,  $\|T\|_1 \leq \dim \text{Ran}(T) \|T\|$ .
- ii) If  $\|T_n - T\| \rightarrow 0$  as  $n \rightarrow \infty$  and  $\|T_n\|_1 \leq M$  uniformly in  $n$ , then  $\|T\|_1 \leq M$ .

We now state the main proposition of this section.

**Proposition 3.2** *The projectors  $P_j^q(t)$  defined by (3.16) satisfy*

$$\left\| \left( \frac{\partial}{\partial t} \right)^n P_j^{q-1}(t) \left( P_j^q(t) - P_j^{q-1}(t) \right) \right\| \leq \frac{c}{j^{(q+1)\alpha}}$$

for any  $n$  and  $q$  such that  $n + q \leq k$ , and

$$\left\| \left( \frac{\partial}{\partial t} \right)^n P_j^q(t) \left( P_j^q(t) P_j^{q'}(t) - P_j^{q-1}(t) P_j^{q-1'}(t) \right) \right\| \leq \frac{c}{j^{(q+1)\alpha}}$$

for any  $n$  and  $q$  such that  $n + q + 1 \leq k$ .

**Corollary 3.2** *For  $q \leq k - 1$*

$$\left\| P_j^q(t) P_j^{q'}(t) - P_j^{q-1}(t) P_j^{q-1'}(t) \right\| \leq \frac{c}{j^{(q+1)\alpha}}.$$

**Proof:**

$$\begin{aligned}
& P_j^q(t)P_j^{q'}(t) - P_j^{q-1}(t)P_j^{q-1'}(t) = \\
& P_j^q(t) \left( P_j^q(t)P_j^{q'}(t) - P_j^{q-1}(t)P_j^{q-1'}(t) \right) + \\
& P_j^q(t)P_j^{q-1}(t)P_j^{q-1'}(t) - P_j^{q-1}(t)P_j^{q-1}(t)P_j^{q-1'}(t) = \\
& P_j^q(t) \left( P_j^q(t)P_j^{q'}(t) - P_j^{q-1}(t)P_j^{q-1'}(t) \right) + \left[ P_j^{q-1}(t) \left( P_j^q(t) - P_j^{q-1}(t) \right) \right]^* P_j^{q-1'}(t)
\end{aligned} \tag{3.26}$$

where  $\|P_j^{q-1'}(t)\| \leq c$  (see (3.10)).  $\square$

We can now end the proof of theorem 2.1. On the one hand, using lemma i) and the above corollary

$$\left\| \sum_{j=1}^n \left( P_j^q(t)P_j^{q'}(t) - P_j^{q-1}(t)P_j^{q-1'}(t) \right) \right\|_1 \leq c \sum_{j=1}^n \frac{j^\beta}{j^{(q+1)\alpha}} \leq M < \infty \tag{3.27}$$

where  $M$  is independent of  $n$ , provided  $(q+1)\alpha - \beta > 1$ . On the other hand

$$\begin{aligned}
& \left\| \sum_{j=1}^n \left( P_j^q(t)P_j^{q'}(t) - P_j^{q-1}(t)P_j^{q-1'}(t) \right) - \sum_{j=1}^{\infty} \left( P_j^q(t)P_j^{q'}(t) - P_j^{q-1}(t)P_j^{q-1'}(t) \right) \right\| \\
& \leq c \sum_{j=n+1}^{\infty} \frac{1}{j^{(q+1)\alpha}} \rightarrow 0 \text{ as } n \rightarrow \infty
\end{aligned} \tag{3.28}$$

provided  $(q+1)\alpha > 1$ . In view of lemma ii) and corollary 3.1, we can conclude by the Birman-Krein theorem [BK] that  $\sigma_{ac}(U(2\pi)) = \emptyset$  if  $k\alpha - \beta > 1$ , since  $q \leq k-1$ .  $\square$

## 4 Technicalities

We present in this final section the proofs of the results stated in the previous section.

**Proof of lemma 3.2:** Let us show that  $G(t)$  is bounded for any  $\alpha > \beta \geq 0$ . We consider  $G(t)$  as an infinite matrix in the orthonormal basis

$$\left\{ \psi_j^{n_j}(t) \right\}_{j=1, \dots, \infty}^{n_j=1, \dots, \text{mult}(\sigma_j)}, \quad \text{s.t. } P_j(t)\psi_j^{n_j}(t) = \psi_j^{n_j}(t). \tag{4.1}$$

Its matrix elements are given by

$$\begin{aligned}
g_{j,k}^{n_j, n_k}(t) & \equiv \langle \psi_j^{n_j}(t) | G(t) \psi_k^{n_k}(t) \rangle \\
& = \langle \psi_j^{n_j}(t) | \frac{1}{2\pi i} \oint_{\Gamma_j} A_j(t, \lambda) R(t, \lambda) d\lambda P_k(t) \psi_k^{n_k}(t) \rangle.
\end{aligned} \tag{4.2}$$

If  $j \neq k$ , we have, using the first resolvent identity and the Cauchy formula,

$$\begin{aligned}
& \oint_{\Gamma_j} A_j(t, \lambda) R(t, \lambda) d\lambda P_k(t) = \\
& -\frac{1}{2\pi i} \oint_{\Gamma_j} \oint_{\Gamma_k} A_j(t, \lambda) R(t, \lambda) R(t, \mu) d\lambda d\mu = \\
& -\frac{1}{2\pi i} \oint_{\Gamma_j} \oint_{\Gamma_k} \left( \frac{A_j(t, \lambda) R(t, \lambda)}{\lambda - \mu} - \frac{A_j(t, \lambda) R(t, \mu)}{\lambda - \mu} \right) d\lambda d\mu = \\
& \frac{1}{2\pi i} \oint_{\Gamma_j} \oint_{\Gamma_k} \frac{A_j(t, \lambda) R(t, \mu)}{\lambda - \mu} d\lambda d\mu.
\end{aligned} \tag{4.3}$$

As a consequence of the spectral hypothesis  $S$  (see [H4]),

$$|\lambda - \mu| \geq c((j+1)^\alpha + (j+2)^\alpha + \dots + k^\alpha) \geq c|j^{\alpha+1} - k^{\alpha+1}|, \quad (4.4)$$

so that by virtue of the hypothesis and our choice of paths (3.4), (3.5) we get the estimate

$$g_{j,k}^{n_j, n_k}(t) \leq \frac{c}{|j^{\alpha+1} - k^{\alpha+1}|}. \quad (4.5)$$

If  $j = k$ , we have

$$g_{j,j}^{n_j, n_j}(t) \leq \frac{c}{j^\alpha}. \quad (4.6)$$

According to the Schur condition, [Ka] example 2.3 p.143,

$$\|G(t)\| \leq \max \left( \sup_{\substack{j, n_j \\ n_k}} \sum_{k=1}^{\infty} |g_{j,k}^{n_j, n_k}(t)|, \sup_{\substack{k, n_k \\ n_j}} \sum_{j=1}^{\infty} |g_{j,k}^{n_j, n_k}(t)| \right). \quad (4.7)$$

Thus, since  $n_k \leq ck^\beta$ , we have to show that

$$\sup_j \sum_{\substack{k=1 \\ k \neq j}}^{\infty} \frac{k^\beta}{|j^{\alpha+1} - k^{\alpha+1}|} < \infty \quad \text{and} \quad \sup_j \frac{j^\beta}{j^\alpha} < \infty. \quad (4.8)$$

But it is proven in the appendix of [H4] that for  $\alpha > \beta$ ,

$$\sum_{\substack{k=1 \\ k \neq j}}^{\infty} \frac{k^\beta}{|j^{\alpha+1} - k^{\alpha+1}|} = \mathcal{O}(j^{\beta-\alpha} \ln j) \quad (4.9)$$

so that (4.8) is indeed true provided  $\alpha > \beta \geq 0$ . To prove the strong continuity of  $G(t)$ , we introduce the strongly continuous approximations

$$G_N(t) = \sum_{j=1}^N P_j(t) \frac{1}{2\pi i} \oint_{\Gamma_j} A_j(t, \lambda) R(t, \lambda) d\lambda. \quad (4.10)$$

Applying the Schur condition again, and (4.9), we obtain

$$\lim_{N \rightarrow \infty} \sup_{t \in [0, 2\pi]} \|G_N(t) - G(t)\| = 0 \quad (4.11)$$

if  $\alpha > \beta \geq 0$ , which shows that  $G(t)$  is strongly continuous. Indeed, the conditions to verify are

$$\sup_{j > N} \left( \sum_{\substack{k=1 \\ k \neq j}}^{\infty} \frac{k^\beta}{|j^{\alpha+1} - k^{\alpha+1}|} + \frac{j^\beta}{j^\alpha} \right) \xrightarrow{N \rightarrow \infty} 0 \quad (4.12)$$

$$\sup_k \sum_{\substack{j=N+1 \\ j \neq k}}^{\infty} \frac{j^\beta}{|j^{\alpha+1} - k^{\alpha+1}|} + \sup_{j > N} \frac{j^\beta}{j^\alpha} \xrightarrow{N \rightarrow \infty} 0. \quad (4.13)$$

According to (4.9), (4.12) is true if  $\alpha > \beta \geq 0$  and the sum in (4.13) can be approximated by

$$\begin{aligned} & \sup_{k \leq N} \sum_{\substack{j=N+1 \\ j \neq k}}^{\infty} \frac{j^\beta}{|j^{\alpha+1} - k^{\alpha+1}|} + \sup_{k > N} \sum_{\substack{j=N+1 \\ j \neq k}}^{\infty} \frac{j^\beta}{|j^{\alpha+1} - k^{\alpha+1}|} \leq \\ & \sum_{j=N+1}^{\infty} \frac{j^\beta}{|j^{\alpha+1} - N^{\alpha+1}|} + \sup_{k > N} \sum_{\substack{j=N+1 \\ j \neq k}}^{\infty} \frac{j^\beta}{|j^{\alpha+1} - k^{\alpha+1}|} \end{aligned} \quad (4.14)$$

where both terms are  $\mathcal{O}(N^{\beta-\alpha} \ln N)$  again.

To consider the differentiability of  $G(t)$  when  $B(t)$  and  $A_j(t, \lambda)$  are strongly  $C^1$ , we also introduce the projector

$$\Pi_N(t) = \mathbf{I} - \sum_{j=1}^N P_j(t). \quad (4.15)$$

By (3.6) and a standard application of the Cauchy formula we can write

$$\Pi'_N(t) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} i d\eta R(t, \xi_N + i\eta) B'(t) R(t, \xi_N + i\eta) \quad (4.16)$$

where  $\xi_N$  lies on the real axis between  $\sigma_N$  and  $\sigma_{N+1}$ . Hence

$$\|\Pi'_N(t)\| \leq c \|B'(t)\| \int_{-\infty}^{\infty} d\eta \frac{1}{cN^{2\alpha} + \eta^2} \leq \frac{c}{N^\alpha} \int_{-\infty}^{\infty} dy \frac{1}{1+y^2} \leq \frac{c}{N^\alpha}. \quad (4.17)$$

Let us compute

$$\begin{aligned} & \frac{\partial}{\partial t} \left\{ \sum_{j=1}^N P_j(t) \frac{1}{2\pi i} \oint_{\Gamma_j} A_j(t, \lambda) R(t, \lambda) d\lambda \right\} = \\ & \sum_{j=1}^N \left( P'_j(t) P_j(t) + P_j(t) P'_j(t) \right) \frac{1}{2\pi i} \oint_{\Gamma_j} A_j(t, \lambda) R(t, \lambda) d\lambda + \\ & \sum_{j=1}^N P_j(t) \frac{1}{2\pi i} \oint_{\Gamma_j} \left( A'_j(t, \lambda) - A_j(t, \lambda) R(t, \lambda) B'(t) \right) R(t, \lambda) d\lambda = \\ & \sum_{j=1}^N P_j(t) \frac{1}{2\pi i} \oint_{\Gamma_j} \left( P'_j(t) A_j(t, \lambda) + A'_j(t, \lambda) - A_j(t, \lambda) R(t, \lambda) B'(t) \right) R(t, \lambda) d\lambda + \\ & \sum_{k=1}^N P'_k(t) P_k(t) \sum_{j=1}^N P_j(t) \frac{1}{2\pi i} \oint_{\Gamma_j} A_j(t, \lambda) R(t, \lambda) d\lambda. \end{aligned} \quad (4.18)$$

Now, using the definition (4.15),

$$\sum_{j=1}^N P'_j(t) + \Pi'_N(t) = \sum_{j=1}^N P'_j(t) P_j(t) + P_j(t) P'_j(t) + \Pi'_N(t) \equiv 0 \quad (4.19)$$

so that

$$\sum_{j=1}^N P'_j(t) P_j(t) = - \sum_{j=1}^N P_j(t) P'_j(t) - \Pi'_N(t). \quad (4.20)$$

Consequently, the last term of (4.18) can be written as

$$-\sum_{k=1}^N P_k(t)P'_k(t)G_N(t) - \Pi'_N(t)G_N(t) \equiv -G_{0,N}(t)G_N(t) - \Pi'_N(t)G_N(t), \quad (4.21)$$

where  $G_{0,N}$  approximates  $G_0(t)$  as in (4.10) (see the remark below lemma 3.2). We have thus proved the formula, with a similar notation,

$$G'_N(t) = G_{1,N}(t) - G_{0,N}(t)G_N(t) - \Pi'_N(t)G_N(t) \quad (4.22)$$

where  $G_{1,N}(t)$  tends to  $G_1(t)$  in norm and uniformly in  $t \in [0, 2\pi]$  by our hypothesis on  $\|A'_j(t, \lambda)\|$ . On the other hand,

$$\begin{aligned} G(s)|_{t_0}^t &= G_N(s)|_{t_0}^t + (G(s) - G_N(s))|_{t_0}^t = \\ &= \int_{t_0}^t G'_N(s)ds + (G(s) - G_N(s))|_{t_0}^t = \\ &= \int_{t_0}^t (G_{1,N}(s) - G_{0,N}(s)G_N(s) - \Pi'_N(s)G_N(s)) ds + (G(s) - G_N(s))|_{t_0}^t \end{aligned} \quad (4.23)$$

where the integrand tends to  $G_1(s) - G_0(s)G(s)$  in norm and uniformly in  $s \in [0, 2\pi]$ , by (4.11) and (4.17). Thus we can take the limit  $N \rightarrow \infty$  inside the integral in the above equation which eventually yields

$$G(t) - G(t_0) = \int_{t_0}^t G_1(s) - G_0(s)G(s)ds. \quad (4.24)$$

□

**Proof of lemma 3.3:** In order to simplify the notation, we drop the indices  $q$  so that we are lead to consider

$$H(t) = H_0 + B(t) \quad (4.25)$$

where  $H_0$  satisfies  $S$  with  $\alpha > \beta \geq 0$ ,  $B(t)$  satisfies  $R_1$  and the operator

$$K(t) = \sum_{j=1}^{\infty} P_j(t)P'_j(t) \quad (4.26)$$

where  $P_j(t)$  is defined by (3.3). We first have to show that there exists a unitary operator  $V(t)$  strongly  $C^1$  on  $D$ , mapping  $D$  into  $D$  such that for any  $\varphi \in D$  and  $t \in \mathbf{R}$

$$iV'(t)\varphi = (H(t) - iK(t))V(t)\varphi, \quad V(0) = \mathbf{I}. \quad (4.27)$$

Note that as  $K(t)$  is strongly continuous only, we cannot invoke theorem X70 of [RS] directly. We shall instead make use of a theorem of perturbation of evolution operators, theorem 3.4 p.198 in [Kr]. Since  $H'(t) = B'(t)$  is bounded, there exists a unitary  $U(t)$  with the above properties such that

$$iU'(t)\varphi = H(t)U(t)\varphi, \quad U(0) = \mathbf{I}, \quad (4.28)$$

if  $\varphi \in D$ . Then, according to the above mentionned theorem, there exists a unitary  $V(t)$  associated with (4.27) possessing the required properties, provided the perturbation  $K(t)$  and the operator  $H(t)K(t)R(t, 0)$  are bounded and strongly continuous. Let  $\varphi \in D$ . We have for any  $j \geq 1$

$$H(t)P_j(t)\varphi = P_j(t)H(t)\varphi \quad (4.29)$$

so that (applying lemma 1.3 p.178 [Kr] for example)

$$(H'(t)P_j(t) + H(t)P_j'(t))\varphi = (P_j'(t)H(t) + P_j(t)H'(t))\varphi. \quad (4.30)$$

Hence we compute, using (4.25) and the completeness of the projectors  $P_j(t)$ ,

$$\begin{aligned} H(t)K(t)R(t,0) &= \sum_{j=1}^{\infty} P_j(t)H(t)P_j'(t)R(t,0) \\ &= \sum_{j=1}^{\infty} P_j(t) \left( P_j'(t)H(t) + P_j(t)B'(t) - B'(t)P_j(t) \right) R(t,0) \\ &= \sum_{j=1}^{\infty} \left( P_j(t)P_j'(t) + P_j(t)B'(t)R(t,0) - P_j(t)B'(t)R(t,0)P_j(t) \right) \\ &= K(t) + B'(t)R(t,0) - \sum_{j=1}^{\infty} P_j(t)B'(t)R(t,0)P_j(t). \end{aligned} \quad (4.31)$$

The last operator is bounded,

$$\begin{aligned} &\left\| \sum_{j=1}^{\infty} P_j(t)B'(t)R(t,0)P_j(t)\varphi \right\|^2 = \sum_{j=1}^{\infty} \|P_j(t)B'(t)R(t,0)P_j(t)\varphi\|^2 \\ &\leq \sum_{j=1}^{\infty} \|P_j(t)B'(t)R(t,0)\|^2 \|P_j(t)\varphi\|^2 \leq \|B'(t)R(t,0)\|^2 \sum_{j=1}^{\infty} \|P_j(t)\varphi\|^2 \\ &= \|B'(t)R(t,0)\|^2 \|\varphi\|^2, \end{aligned} \quad (4.32)$$

and strongly continuous since it is a sum of strongly continuous operators converging uniformly in  $t \in [0, 2\pi]$ . Indeed, it is sufficient to note that

$$\begin{aligned} &\sum_{j=N+1}^{\infty} \|P_j(t)\varphi\|^2 = \|(\mathbf{I} - \Pi_N(t))\varphi\|^2 = \\ &\left\| \left( \mathbf{I} - \Pi_N(0) - \int_0^t ds \Pi_N'(s) \right) \varphi \right\|^2 \leq \|(\mathbf{I} - \Pi_N(0))\varphi\|^2 + \frac{c}{N^\alpha} \|\varphi\|^2 \xrightarrow{N \rightarrow \infty} 0 \end{aligned} \quad (4.33)$$

uniformly in  $t \in [0, 2\pi]$ , according to (4.17).

To prove the intertwining property

$$V(t)P_j(0) = P_j(t)V(t), \quad \forall t \in [0, 2\pi] \text{ and } \forall j \geq 1, \quad (4.34)$$

we approximate  $V(t)$  by evolution operators  $V_N(t)$ ,  $N = 1, \dots, \infty$ , associated with the equations

$$iV_N'(t)\varphi = (H(t) - iK_N(t) - i\Pi_N(t)\Pi_N'(t))V_N(t)\varphi, \quad V_N(0) = \mathbf{I} \quad (4.35)$$

where  $\varphi \in D$  and  $t \in \mathbf{R}$  and

$$K_N(t) = \sum_{j=1}^N P_j(t)P_j'(t) \equiv G_{0,N}(t). \quad (4.36)$$

As above, we deduce that  $V_N(t)$  are unitary, strongly  $C^1$  on  $D$  and map  $D$  into  $D$  since  $\Pi_N(t)$  commutes with  $H(t)$  and

$$\sum_{j=1}^N P_j(t) + \Pi_N(t) = \mathbf{I}. \quad (4.37)$$

Moreover, since we deal with a finite number of projectors, we have the standard property, see e.g.[Kr] §3, Chap. IV,

$$\begin{aligned} V_N(t)P_j(0) &= P_j(t)V_N(t) \quad \forall j = 1, \dots, N \\ V_N(t)\Pi_N(0) &= \Pi_N(t)V_N(t) \quad \forall t \in [0, 2\pi]. \end{aligned} \quad (4.38)$$

We compute

$$\begin{aligned} V(t)P_j(0) - P_j(t)V(t) &= \\ (V(t) - V_N(t))P_j(0) - P_j(t)(V(t) - V_N(t)) &+ (V_N(t)P_j(0) - P_j(t)V_N(t)) \end{aligned} \quad (4.39)$$

where the last bracket vanishes if  $j \leq N$ . Then, for any  $\varphi \in D$ ,

$$i \frac{\partial}{\partial t} \left( V_N^{-1}(t)V(t)\varphi \right) = iV_N^{-1}(t) (K_N(t) - K(t) + \Pi_N(t)\Pi'_N(t)) V(t)\varphi \quad (4.40)$$

so that

$$\|V(t) - V_N(t)\| \leq \int_0^t ds \|K_N(s) - K(s) + \Pi_N(s)\Pi'_N(s)\|. \quad (4.41)$$

But the integrand tends to zero uniformly in  $s$  as  $N$  tends to infinity (see (4.11) and (4.17)) so that

$$\lim_{N \rightarrow \infty} \|V(t) - V_N(t)\| = 0 \quad \forall t \in [0, 2\pi]. \quad (4.42)$$

Consequently, we deduce from (4.39) that  $V(t)P_j(0) - P_j(t)V(t) \equiv 0$  for any  $j \geq 1$  and any  $t \in [0, 2\pi]$ .  $\square$

**Proof of proposition 3.2:** The proof is by induction. Let us recall that  $W(t)$  is strongly  $C^k$  and  $q \leq k - 1$ . We first note that, at the cost of a possible relabelling, we can assume that for  $j$  large enough the spectral sets  $\sigma_j^s(t)$ ,  $s = 0, 1, \dots, q + 1$ , can be surrounded by the same path  $\Gamma_j$  such that

$$\begin{aligned} |\Gamma_j| &\leq cj^\alpha \\ \text{dist}(\Gamma_j, \cup_{j=1}^\infty \sigma_j^s(t)) &\geq cj^\alpha, \quad s = 0, 1, \dots, q + 1. \end{aligned} \quad (4.43)$$

Indeed,  $H_s(t) = H_0(t) + iK_{s-1}(t)$  where

$$\max_{s=1, \dots, q+1} \sup_{t \in [0, 2\pi]} \|K_{s-1}(t)\| \leq c. \quad (4.44)$$

We set, using the second resolvent identity,

$$\begin{aligned} T_j^q(t) &\equiv P_j^{q+1}(t) - P_j^q(t) \\ &= -\frac{1}{2\pi i} \oint_{\Gamma_j} (R_{q+1}(t, \lambda) - R_q(t, \lambda)) d\lambda \\ &= \frac{1}{2\pi} \oint_{\Gamma_j} R_q(t, \lambda) (K_q(t) - K_{q-1}(t)) R_{q+1}(t, \lambda) d\lambda. \end{aligned} \quad (4.45)$$

**Lemma 4.1**

- a)  $T_j^q(t) = P_j^{q+1}(t)T_j^q(t) + T_j^q(t)P_j^q(t)$
- b)  $P_j^{q+1}(t)T_j^q(t) = P_j^{q+1}(t)P_j^q(t)T_j^q(t)(\mathbf{I} - T_j^q(t))^{-1}$ .

**Proof:**

a) Since the operators  $P_j^s(t)$  are projectors we can write

$$P_j^{q+1}(t) = \left( P_j^q(t) + T_j^q(t) \right)^2 = P_j^q(t) + P_j^q(t)T_j^q(t) + T_j^q(t)P_j^q(t) + T_j^q(t)T_j^q(t) \quad (4.46)$$

hence

$$T_j^q(t) = \left( P_j^q(t) + T_j^q(t) \right) T_j^q(t) + T_j^q(t)P_j^q(t). \quad (4.47)$$

b) For the same reason we have

$$\begin{aligned} P_j^{q+1}(t)T_j^q(t) &= P_j^{q+1}(t) \left( P_j^q(t) + T_j^q(t) \right) T_j^q(t) \\ &= P_j^{q+1}(t)P_j^q(t)T_j^q(t) + P_j^{q+1}(t)T_j^q(t)T_j^q(t) \end{aligned} \quad (4.48)$$

so that

$$P_j^{q+1}(t)T_j^q(t)(\mathbf{I} - T_j^q(t)) = P_j^{q+1}(t)P_j^q(t)T_j^q(t). \quad (4.49)$$

From (4.45) and (4.43) we obtain the estimate

$$\begin{aligned} \|T_j^q(t)\| &\leq c \left\| \oint_{\Gamma_j} R_q(t, \lambda)(K_q(t) - K_{q-1}(t))R_{q+1}(t, \lambda)d\lambda \right\| \\ &\leq c|\Gamma_j| \frac{1}{j^{2\alpha}} \|K_q(t) - K_{q-1}(t)\| \leq \frac{c}{j^\alpha} \end{aligned} \quad (4.50)$$

which shows that for  $j$  large enough the operator  $(\mathbf{I} - T_j^q(t))$  is invertible.  $\square$

**Remark:** We can deduce easily from this lemma and the self adjointness of  $P_j^s(t)$  the following estimate for  $j$  large enough

$$\|T_j^q(t)\| \leq c \|P_j^q(t)T_j^q(t)\| \quad (4.51)$$

which yields the sharper estimate on  $\|T_j^q(t)\|$

$$\|T_j^q(t)\| \leq \frac{c}{j^{(q+2)\alpha}}, \quad (4.52)$$

provided proposition 3.2 is true.

Let us assume the proposition holds for the index  $q$  and let us check its validity for  $q + 1$ . Thus we have to estimate for any  $n$  such that  $n + q + 1 \leq k$ , see (4.45),

$$\left( \frac{\partial}{\partial t} \right)^n P_j^q(t)T_j^q(t) = \frac{1}{2\pi} \oint_{\Gamma_j} \left( \frac{\partial}{\partial t} \right)^n R_q(t, \lambda)P_j^q(t)(K_q(t) - K_{q-1}(t))R_{q+1}(t, \lambda)d\lambda. \quad (4.53)$$

We compute

$$\begin{aligned} &P_j^q(t)(K_q(t) - K_{q-1}(t)) = \\ &P_j^q(t) \left( P_j^q(t)K_q(t) - P_j^{q-1}(t)K_{q-1}(t) \right) + P_j^q(t)P_j^{q-1}(t)K_{q-1}(t) - P_j^q(t)K_{q-1}(t) = \\ &P_j^q(t) \left( P_j^q(t)P_j^{q'}(t) - P_j^{q-1}(t)P_j^{q-1'}(t) \right) - P_j^q(t)T_j^{q-1}(t)K_{q-1}(t). \end{aligned} \quad (4.54)$$



Using property b), we get

$$\begin{aligned}
& \left( \frac{\partial}{\partial t} \right)^n P_j^q(t) T_j^q(t) = \\
& \frac{1}{2\pi} \oint_{\Gamma_j} \left( \frac{\partial}{\partial t} \right)^n R_q(t, \lambda) P_j^q(t) \left( P_j^q(t) P_j^{q'}(t) - P_j^{q-1}(t) P_j^{q-1'}(t) \right) R_{q+1}(t, \lambda) d\lambda \\
& - \frac{1}{2\pi} \oint_{\Gamma_j} \left( \frac{\partial}{\partial t} \right)^n R_q(t, \lambda) P_j^q(t) P_j^{q-1}(t) T_j^{q-1}(t) \left( \mathbf{I} - T_j^{q-1}(t) \right)^{-1} K_{q-1}(t) R_{q+1}(t, \lambda) d\lambda.
\end{aligned} \tag{4.55}$$

By the Leibnitz formula or eq.(2.36) in [JP2], the  $n^{\text{th}}$  time derivative of  $R_s(t, \lambda)$  for  $s = 0, \dots, q+1$  is given by a sum of products of operators  $R_s(t, \lambda)$  and

$$\left( \frac{\partial}{\partial t} \right)^l H_s(t) = \left( \frac{\partial}{\partial t} \right)^l W(t) + i \left( \frac{\partial}{\partial t} \right)^l K_{s-1}(t), \quad l = 1, \dots, n. \tag{4.56}$$

Hence, provided  $n + q + 1 \leq k$  (see section 3.2), and making use of (4.43) we get

$$\left\| \left( \frac{\partial}{\partial t} \right)^n R_s(t, \lambda) |_{\lambda \in \Gamma_j} \right\| \leq \frac{c}{j^\alpha} \quad \text{and} \quad \left\| \left( \frac{\partial}{\partial t} \right)^n P_j^s(t) \right\| \leq \frac{c}{j^\alpha}, \quad n \geq 1 \tag{4.57}$$

for  $s = 0, \dots, q+1$ . The same argument can be applied to estimate the  $n^{\text{th}}$  time derivative of  $\left( \mathbf{I} - T_j^{q-1}(t) \right)^{-1}$ . Indeed, for  $j$  large enough,

$$\left( \mathbf{I} - T_j^{q-1}(t) \right)^{-1'} = \left( \mathbf{I} - T_j^{q-1}(t) \right)^{-1} T_j^{q-1'}(t) \left( \mathbf{I} - T_j^{q-1}(t) \right)^{-1} \tag{4.58}$$

and

$$\left( \frac{\partial}{\partial t} \right)^l T_j^{q-1}(t) = \frac{1}{2\pi} \oint_{\Gamma_j} \left( \frac{\partial}{\partial t} \right)^l R_{q-1}(t, \lambda) (K_{q-1}(t) - K_{q-2}(t)) R_q(t, \lambda) d\lambda \tag{4.59}$$

so that

$$\left\| \left( \frac{\partial}{\partial t} \right)^n \left( \mathbf{I} - T_j^{q-1}(t) \right)^{-1} \right\| \leq c, \quad \forall n + q + 1 \leq k. \tag{4.60}$$

Thus, invoking the induction hypothesis and (4.57), we get

$$\left\| \left( \frac{\partial}{\partial t} \right)^n P_j^q(t) T_j^q(t) \right\| \leq \frac{c}{j^{(q+2)\alpha}}, \quad \forall n + q + 1 \leq k. \tag{4.61}$$

Now, if  $q + 2 \leq k$  we compute

$$\begin{aligned}
& P_j^{q+1}(t) \left( P_j^{q+1}(t) P_j^{q+1'}(t) - P_j^q(t) P_j^{q'}(t) \right) = \\
& P_j^{q+1}(t) \left( T_j^q(t) P_j^{q'}(t) + P_j^{q+1}(t) T_j^{q'}(t) \right) = \\
& \left( P_j^{q+1}(t) T_j^q(t) \right) P_j^{q'}(t) + P_j^{q+1}(t) \left( P_j^{q+1}(t) T_j^q(t) \right)' - P_j^{q+1}(t) P_j^{q+1'}(t) T_j^q(t)
\end{aligned} \tag{4.62}$$

where the last term can be written as

$$P_j^{q+1}(t) P_j^{q+1'}(t) T_j^q(t) = P_j^{q+1'}(t) (T_j^q(t) P_j^q(t)) \tag{4.63}$$

using the identity  $Q(t)Q'(t)Q(t) \equiv 0$  for any projector  $Q(t)$  and lemma 4.1 a). Then, by lemma 4.1 b),

$$\begin{aligned}
& P_j^{q+1}(t) \left( P_j^{q+1}(t) P_j^{q+1'}(t) - P_j^q(t) P_j^{q'}(t) \right) = \\
& P_j^{q+1}(t) (P_j^q(t) T_j^q(t)) \left( \mathbf{I} - T_j^q(t) \right)^{-1} P_j^{q'}(t) + \\
& P_j^{q+1}(t) \left\{ P_j^{q+1}(t) (P_j^q(t) T_j^q(t)) \left( \mathbf{I} - T_j^q(t) \right)^{-1} \right\}' - P_j^{q+1'}(t) (P_j^q(t) T_j^q(t))^*.
\end{aligned} \tag{4.64}$$

Finally, considering the Leibnitz formula again, (4.61), (4.60), (4.57) and

$$\left(\frac{\partial}{\partial t}\right)^l (P_j^q(t)T_j^q(t))^* = \left\{ \left(\frac{\partial}{\partial t}\right)^l (P_j^q(t)T_j^q(t)) \right\}^* \quad (4.65)$$

we obtain similarly

$$\left\| \left(\frac{\partial}{\partial t}\right)^n P_j^{q+1}(t) \left( P_j^{q+1}(t)P_j^{q+1'}(t) - P_j^q(t)P_j^{q'}(t) \right) \right\| \leq \frac{c}{j^{(q+2)\alpha}} \quad (4.66)$$

for any  $n$  such that  $n + q + 2 \leq k$ . At last, the induction hypothesis is readily verified for  $q = 1$  with  $n + 1 \leq k$  on

$$\begin{aligned} P_j^0(t)T_j^0(t) &= \frac{1}{2\pi} \oint_{\Gamma_j} R_0(t, \lambda) P_j^0(t) K_0(t) R_1(t, \lambda) d\lambda \\ &= \frac{1}{2\pi} \oint_{\Gamma_j} R_0(t, \lambda) P_j^0(t) P_j^{0'}(t) R_1(t, \lambda) d\lambda, \end{aligned} \quad (4.67)$$

see (4.57) and (4.43), which implies, as above, that it is satisfied by

$$P_j^1(t) \left( P_j^1(t)P_j^{1'}(t) - P_j^0(t)P_j^{0'}(t) \right) \quad (4.68)$$

with  $n + 2 \leq k$ . □

## References

- [B] J.Bellissard: Stability and Instability in Quantum Mechanics in "Trends and developments in the eighties", S.Albeverio, Ph.Blanchard Edts, World Scientific, Singapore 1985.
- [J] H.R.Jauslin: Stability and Chaos in Classical and Quantum Hamiltonian Systems, II Granada Seminar on Computational Physics, P.Garrido, J.Marro Edts, World Scientific, Singapore 1983.
- [Y] K.Yajima: Scattering Theory for Schrödinger Equations with Potential Periodic in Time, *J.Math.Soc.Japan***29** (1977) 729-743.
- [H1] J.Howland: Scattering Theory for Hamiltonians Periodic in Time, *Indiana Univ.Math.J.***28** (1979) 471-494.
- [BJL] P.M.Blekher, H.R.Jauslin, J.L.Lebowitz: Floquet Spectrum for Two-Level Systems in Quasiperiodic Time-Dependent Fields, *J.Stat.Phys.***68** (1992) 271-310.
- [H2] J.Howland: Perturbation Theory of Dense Point Spectra, *J.Funct.Anal.***74** (1987) 52-80.
- [HLS] G.Hagedorn, M.Loss, J.Slawny: Non Stochasticity of Time-dependent Quadratic Hamiltonians and Spectra of Cononical Transformations, *J.Phys.A.***19** (1986) 521-531.
- [C1] M.Combescure: Quantum Stability Problem for Time Periodic Perturbations of the Harmonic Oscillator, *Ann.Inst.H.Poincaré***47** (1987) 63-83.
- [C2] M.Combescure: Spectral Properties of a Periodically Kicked Quantum Hamiltonian, *J.Stat.Phys.***59** (1990) 679-690.
- [H3] J.Howland: Floquet Operator with Singular Spectrum I, *Ann.Inst.H.Poincaré***49** (1989) 309-323.

- [H4] J.Howland: Floquet Operator with Singular Spectrum II, *Ann.Inst.H.Poincaré***49** (1989) 325-334.
- [H5] J.Howland: Stability of Quantum Oscillators, *J.Phys.A.***25** (1992) 5177-5181.
- [N1] G.Nenciu: Floquet Operators without Absolutely Continuous Spectrum, *Preprint Zürich ETH-TH/92-5* (1992).
- [RS] M.Reed, B.Simon: "Methods of Modern Mathematical Physics, II Fourier Analysis, Self-Adjointness", Academic Press, New York, San Francisco, London 1975.
- [Ka] T.Kato: "Perturbation Theory for Linear Operators", Springer-Verlag, Berlin, Heidelberg, New York 1966.
- [N2] G.Nenciu: Asymptotic Invariant Subspaces, Adiabatic Theorems and Block Diagonalisation in "Recent Developments in Quantum Mechanics", A.Boutet de Monvel et.al. Edts, Kluwer Academic Publishers, Dordrecht 1991.
- [BK] M.S.Birman, M.G.Krein: On the Theory of Wave Operators and Scattering Theory, *Soviet.Math.Dokl.***3** (1962) 740-744.
- [JP1] A.Joye, C.Pfister: Full Asymptotic Expansion of Transition Probabilities in the Adiabatic Limit, *J.Phys.A.***24** (1991) 753-766.
- [JP2] A.Joye, C.Pfister: Superadiabatic Evolution and Adiabatic Transition Probability between Two Non-degenerate Levels Isolated in the Spectrum, *J.Math.Phys.***34** (1993) 454-479.
- [Kr] S.G.Krein: "Linear Differential Equations in Banach Spaces", American Mathematical Society Vol.29, Providence 1971.