
TAMAGAWA NUMBERS OF DIAGONAL CUBIC SURFACES, NUMERICAL EVIDENCE*

by

Emmanuel Peyre & Yuri Tschinkel

Abstract. — A refined version of Manin’s conjecture about the asymptotics of points of bounded height on Fano varieties has been developed by Batyrev and the authors. We test numerically this refined conjecture for some diagonal cubic surfaces.

Résumé. — Une version raffinée d’une conjecture de Manin sur le comportement asymptotique des points de hauteur bornée sur les variétés de Fano a été proposée par Batyrev et les auteurs. Nous testons numériquement cette conjecture pour diverses surfaces cubiques.

1. Introduction

The aim of this paper is to test numerically a refined version of a conjecture of Manin concerning the asymptotic for the number of rational points of bounded height on Fano varieties (see [BM] or [FMT] for Manin’s conjecture and [Pe1] or [BT3] for its refined versions).

Let V be a smooth Fano variety over a number field F and ω_V^{-1} its anticanonical line bundle. Let $\text{Pic}(V)$ be the Picard group and $\text{NS}(V)$ the Néron-Severi group of V . We denote by $\text{Val}(F)$ the set of all places of F and by F_v the v -adic completion of F . Let $(\|\cdot\|_v)_{v \in \text{Val}(F)}$ be an adelic metric on ω_V^{-1} . By definition, this is a family of v -adically continuous metrics on $\omega_V^{-1} \otimes F_v$ which for almost all valuations v are given by a smooth model of V (see [Pe2]). These data define a

2000 Mathematics Subject Classification. — primary 11D25; secondary 14G05, 14J25.

*Math. Comp. **70** (2000), n° 233, 367–387

height \mathbf{H} on the set of rational points $V(F)$ given by

$$\forall x \in V(F), \quad \forall y \in \omega_V^{-1}(x), \quad \mathbf{H}(x) = \prod_{v \in \text{Val}(F)} \|y\|_v^{-1}.$$

For every open subset $U \subset V$ and every real number H we have

$$n_{U, \mathbf{H}}(H) = \#\{x \in U(F) \mid \mathbf{H}(x) \leq H\} < \infty.$$

The problem is to understand the asymptotic behavior of $n_{U, \mathbf{H}}(H)$ as H goes to infinity. It is expected that at least for Del Pezzo surfaces the following asymptotic formula holds:

$$n_{U, \mathbf{H}}(H) = \theta_{\mathbf{H}}(V) H (\log H)^{t-1} (1 + o(1))$$

as $H \rightarrow \infty$, over appropriate finite extensions E/F of the groundfield. Here the open set U is the complement to exceptional curves, $\theta_{\mathbf{H}}(V) > 0$ and t is the rank of the Picard group of V over E . We have counter-examples to this conjecture in every dimension ≥ 3 [BT2] (see [BT3] for a discussion of higher dimensional varieties).

In this paper we focus on the constant $\theta_{\mathbf{H}}(V)$. On the one hand, there is a theoretical description

$$(1) \quad \theta_{\mathbf{H}}(V) = \alpha(V) \beta(V) \tau_{\mathbf{H}}(V)$$

where $\tau_{\mathbf{H}}(V)$ is a Tamagawa number associated to the metrized anticanonical line bundle [Pe1], $\alpha(V)$ is a rational number defined in terms of the cone of effective divisors [Pe1] and the integer $\beta(V)$ is a cohomological invariant, which first appeared in asymptotic formulas in [BT1].

On the other hand, let us consider a diagonal cubic surface $V \subset \mathbf{P}_{\mathbf{Q}}^3$ given by

$$ax^3 + by^3 + cz^3 + dt^3 = 0,$$

with $a, b, c, d \in \mathbf{Z}$ and $abcd \neq 0$. Our counting problem can be formulated as follows: find all quadruples of integers (x, y, z, t) with

$$\text{g.c.d.}(x, y, z, t) = 1 \quad \text{and} \quad \max\{|x|, |y|, |z|, |t|\} \leq H$$

which satisfy the equation above. Quadruples differing by a sign are counted once. A *proof* of an asymptotic of the type (1) for smooth cubic surfaces seems to be out of reach of available methods, but one can numerically search for solutions of bounded height. The cubics with coefficients $(1, 1, 1, 2)$ and $(1, 1, 1, 3)$ and height $H \leq 2000$ were treated by Heath-Brown in [HB]. In both cases weak approximation fails. Swinnerton-Dyer made substantial progress towards an interpretation of the constant $\tau_{\mathbf{H}}(V)$ [SD]. In particular, he suggested that

the adelic integral defining $\tau_{\mathbf{H}}(V)$ should be taken over the closure of rational points $\overline{V(F)} \subset V(\mathbf{A}_F)$, rather than the whole adelic space.

Our goal is to compute the theoretical constant $\theta_{\mathbf{H}}(V)$ explicitly for certain diagonal cubic surfaces with and without obstruction to weak approximation and to compare the result with numerical data (with height $H \leq 100000$). We observe a very good accordance.

In section 2 we define the Tamagawa number. This definition is slightly different from the one in [Pe1], but the numbers coincide conjecturally. In sections 3, 4 and 5 we explain how to compute it. There is a subtlety at the places of bad reduction, notable at 3, overlooked previously. In section 6 we compute the Brauer-Manin obstruction to weak approximation. And in section 7 we present the numerical results. These computations were made using a program of Bernstein which is described in [Be].

2. Conjectural constant

Notations 2.1. — If \mathcal{V} is a scheme over a ring A and B an A -algebra, we denote by \mathcal{V}_B the product $\mathcal{V} \times_{\text{Spec } A} \text{Spec } B$ and by $\mathcal{V}(B)$ the set of B -points, that is $\text{Hom}_{\text{Spec } A}(\text{Spec } B, \mathcal{V})$. For any field E , we denote by \overline{E} a fixed algebraic closure and by \overline{V} the variety $V_{\overline{E}}$.

If F is a number field, we identify the set of finite places with the set of prime ideals in \mathcal{O}_F . We denote by d_F the absolute value of its discriminant. If \mathfrak{p} is a finite place of F , then $\mathcal{O}_{\mathfrak{p}}$ is the ring of integers in $F_{\mathfrak{p}}$ and $\mathbf{F}_{\mathfrak{p}}$ its residue field.

In the sequel we will always assume that V is a smooth projective geometrically integral variety over a number field F satisfying the following conditions:

- (i) The group $H^i(V, \mathcal{O}_V)$ is trivial for $i = 1$ or 2 ,
- (ii) $\text{Pic}(\overline{V})$ has no torsion,
- (iii) ω_V^{-1} belongs to the interior of the cone of classes of effective divisors $\Lambda_{\text{eff}}(V)$.

Since V is projective, the adelic space $V(\mathbf{A}_F)$ of V coincides with the product

$$\prod_{v \in \text{Val}(F)} V(F_v).$$

One says that *weak approximation* holds for V if the diagonal map from $V(F)$ to $V(\mathbf{A}_F)$ has a dense image. Our definition of the conjectural asymptotic constant $\theta_{\mathbf{H}}(V)$ uses the notion of the Brauer-Manin obstruction to weak approximation, which we now recall.

Notations 2.2. — Let $\mathrm{Br}(V)$ be the étale cohomology group $H_{\text{ét}}^2(V, \mathbf{G}_m)$. If A belongs to $\mathrm{Br}(V)$ and E is a field over F then, for any P in $V(E)$, we denote by $A(P)$ the evaluation of A at P . For any class A , there exists a finite set of places S of F such that

$$\forall v \notin S, \quad \forall P_v \in V(F_v), \quad A(P_v) = 0,$$

(see, for example, [CT2, lemma 1]). For any v in $\mathrm{Val}(F)$, let $\mathrm{inv}_v : \mathrm{Br}(F_v) \rightarrow \mathbf{Q}/\mathbf{Z}$ be the invariant given by local class field theory normalized so that the sequence

$$0 \rightarrow \mathrm{Br}(F) \rightarrow \bigoplus_{v \in \mathrm{Val}(F)} \mathrm{Br}(F_v) \xrightarrow{\sum \mathrm{inv}_v} \mathbf{Q}/\mathbf{Z} \rightarrow 0$$

is exact. Let ρ_A be the composite map

$$V(\mathbf{A}_F) \rightarrow \bigoplus_{v \in \mathrm{Val}(F)} \mathrm{Br}(F_v) \xrightarrow{\sum \mathrm{inv}_v} \mathbf{Q}/\mathbf{Z}.$$

Then one defines

$$V(\mathbf{A}_F)^{\mathrm{Br}} = \bigcap_{A \in \mathrm{Br} V} \ker(\rho_A) \subset V(\mathbf{A}_F).$$

The above exact sequence gives an inclusion $V(F) \subset V(\mathbf{A}_F)^{\mathrm{Br}}$. The *Brauer-Manin obstruction to weak approximation*, introduced by Manin in [Ma1] and by Colliot-Thélène and Sansuc in [CTS] is defined as the condition

$$V(\mathbf{A}_F)^{\mathrm{Br}} \subsetneq V(\mathbf{A}_F).$$

Remark 2.1. — It is conjectured that the closure of the set of rational points $\overline{V(F)} \subset V(\mathbf{A}_F)$ in fact coincides with $V(\mathbf{A}_F)^{\mathrm{Br}}$, at least for Del Pezzo surfaces. This has been proved, for example, by Salberger and Skorobogatov for a smooth complete intersection of two quadrics in \mathbf{P}^4 if $V(F)$ is not empty (see [SaSk]). It would be very interesting to see an example of a cubic surface V with $t = \mathrm{rk} \mathrm{Pic}(V) = 1$ where weak approximation holds, or where one could actually prove that $\overline{V(F)} = V(\mathbf{A}_F)^{\mathrm{Br}}$, assuming that $V(F)$ is Zariski dense, which by a result of B. Segre (see [Ma2, §29, §30]) is equivalent to $V(F) \neq \emptyset$.

Notations 2.3. — Let $(\|\cdot\|_v)_{v \in \mathrm{Val}(F)}$ be an adelic metric on ω_V^{-1} and \mathbf{H} the associated height function on $V(F)$. The adelic metrization of the anticanonical line bundle yields for any place v of F a measure $\omega_{\mathbf{H},v}$ on the locally compact

space $V(F_v)$, given by the local formula

$$\omega_{\mathbf{H},v} = \left\| \frac{\partial}{\partial x_{1,v}} \wedge \cdots \wedge \frac{\partial}{\partial x_{n,v}} \right\|_v dx_{1,v} \cdots dx_{n,v}.$$

where $x_{1,v}, \dots, x_{n,v}$ are local v -adic analytic coordinates, $\frac{\partial}{\partial x_{1,v}} \wedge \cdots \wedge \frac{\partial}{\partial x_{n,v}}$ is seen as a section of $\omega_{\mathcal{V}}^{-1}$ and the Haar measures $dx_{j,v}$ (for $j = 1, \dots, n$) are normalized by

- if v is a finite place then $\int_{\mathcal{O}_v} dx_{j,v} = 1$,
- if v is real then $dx_{j,v}$ is the standard Lebesgue measure,
- if v is complex then $dx_{j,v} = dz d\bar{z}$.

We choose a finite set S of bad places containing the archimedean ones and a smooth projective model \mathcal{V} of V over the ring of S -integers \mathcal{O}_S . For any \mathfrak{p} in $\text{Val}(F) - S$, the local term of the L -function corresponding to the Picard group is defined by

$$L_{\mathfrak{p}}(s, \text{Pic}(\overline{\mathcal{V}})) = \frac{1}{\text{Det}(1 - (\#\mathbf{F}_{\mathfrak{p}})^{-s} \text{Fr} \mid \text{Pic}(\mathcal{V}_{\overline{\mathbf{F}}_{\mathfrak{p}}}) \otimes \mathbf{Q})},$$

where Fr is the Frobenius map. The corresponding global L -function is given by

$$L_S(s, \text{Pic}(\overline{\mathcal{V}})) = \prod_{\mathfrak{p} \in \text{Val}(F) - S} L_{\mathfrak{p}}(s, \text{Pic}(\overline{\mathcal{V}})),$$

it converges for $\text{Re}(s) > 1$ and has a meromorphic continuation to \mathbf{C} with a pole of order $t = \text{rk Pic}(V)$ at 1. The local convergence factors are defined by

$$\lambda_v = \begin{cases} L_v(1, \text{Pic}(\overline{\mathcal{V}})) & \text{if } v \in \text{Val}(F) - S, \\ 1 & \text{otherwise.} \end{cases}$$

The Weil conjectures, proved by Deligne, imply that the adelic measure

$$\prod_{v \in \text{Val}(F)} \lambda_v^{-1} \omega_{\mathbf{H},v}$$

converges on $V(\mathbf{A}_F)$ (see [Pe1, proposition 2.2.2]).

Definition 2.4. — The *Tamagawa measure* corresponding to \mathbf{H} is defined by

$$\omega_{\mathbf{H}} = \frac{1}{\sqrt{d_F}^{\dim V}} \lim_{s \rightarrow 1} (s-1)^t L_S(s, \text{Pic}(\overline{\mathcal{V}})) \prod_{v \in \text{Val}(F)} \lambda_v^{-1} \omega_{\mathbf{H},v}.$$

The *Tamagawa number* is defined by

$$\tau_{\mathbf{H}}(V) = \omega_{\mathbf{H}}(V(\mathbf{A}_F)^{\text{Br}}).$$

The cohomological constant is given by

$$\beta(V) = \#H^1(F, \text{Pic}(\overline{V})).$$

Let $\text{NS}(V)^{\vee}$ be the lattice dual to $\text{NS}(V)$. It defines a natural Lebesgue measure $\mathbf{d}\mathbf{y}$ on $\text{NS}(V)^{\vee} \otimes \mathbf{R}$. Denote by $\Lambda_{\text{eff}}(V) \subset \text{NS}(V) \otimes \mathbf{R}$ the cone of effective divisors and by $\Lambda_{\text{eff}}(V)^{\vee} \subset \text{NS}(V)^{\vee} \otimes \mathbf{R}$ the dual cone.

Definition 2.5. — We define

$$\alpha(V) = \frac{1}{(t-1)!} \int_{\Lambda_{\text{eff}}(V)^{\vee}} e^{-\langle \omega_V^{-1}, \mathbf{y} \rangle} \mathbf{d}\mathbf{y}.$$

Remarks 2.2. — (i) Of course, for nonsplit cubic surfaces with $\text{rk Pic}(V) = 1$ the constant $\alpha(V) = 1$. However, it is a challenge to compute this constant for a split cubic surface with $\text{rk Pic}(V) = 7$.

(ii) As Salberger, we use $\omega_{\mathbf{H}}(V(\mathbf{A}_F)^{\text{Br}})$ instead of $\omega_{\mathbf{H}}(\overline{V(k)})$ in the definition of $\tau_{\mathbf{H}}(V)$. By remark 2.1 these numbers are conjecturally the same, but only the first one is computable for a general cubic. Also we use the convention of [Pe1, §2.2.5] for the definition of $\alpha(V)$.

Definition 2.6. — We define the constant corresponding to V and \mathbf{H} as

$$\theta_{\mathbf{H}}(V) = \alpha(V)\beta(V)\tau_{\mathbf{H}}(V).$$

3. Measures and density

In this section we relate the local volumes of the variety with the density of solutions modulo \mathfrak{p}^n . Lemma 5.4.6 in [Pe1] relates the local volume for $\omega_{\mathbf{H},\mathfrak{p}}$ to the volume for Leray's measure. We now compare the latter to the density modulo \mathfrak{p}^n .

Notations 3.1. — Let F be a number field and V a smooth complete intersection in \mathbf{P}_F^N defined by m homogeneous polynomials f_i in the algebra $\mathcal{O}_F[X_0, \dots, X_N]$. Let $\delta = N + 1 - \sum_{i=1}^m \deg f_i$. We assume that $\delta \geq 1$. We denote

by $W \subset \mathbf{A}_F^{N+1} - \{0\}$ the cone above V and by $\mathbf{f} : \mathbf{A}_{\mathcal{O}_F}^{N+1} \rightarrow \mathbf{A}_{\mathcal{O}_F}^m$ the map induced by the f_i . Then the Leray form on W is defined locally by

$$(2) \quad \omega_L = (-1)^{Nm - \sum_{j=1}^m k_j} \left(\text{Det} \left(\frac{\partial f_i}{\partial X_{k_j}} \right)_{1 \leq i, j \leq m} \right)^{-1} \\ \times dX_0 \wedge \cdots \wedge \widehat{dX_{k_1}} \wedge \cdots \wedge \widehat{dX_{k_m}} \wedge \cdots \wedge dX_N$$

where $0 \leq k_1 < \cdots < k_m \leq N$. For any v in $\text{Val}(F)$, this form yields a measure $\omega_{L,v}$ on $W(F_v)$.

The following result is well known in the setting of the circle method (see for example [Lac, proposition 1.14]) where it is generally proved using a Fourier inversion formula. It may also be deduced from a more general result of Salberger [Sal, theorem 2.13]. We prove it here in a direct and elementary way.

Proposition 3.1. — *We fix a finite place $v = v_{\mathfrak{p}}$ of F . If all f_i have the same degree, then*

$$\int_{\{x \in \mathcal{O}_{\mathfrak{p}}^{N+1} \mid f(x) \not\equiv 0\}} \omega_{L,v} = \lim_{r \rightarrow +\infty} \frac{\#\{x \in (\mathcal{O}_{\mathfrak{p}/\mathfrak{p}^r})^{N+1} \mid \mathbf{f}(x) = 0 \text{ in } (\mathcal{O}_{\mathfrak{p}/\mathfrak{p}^r})^m\}}{(\#\mathbf{F}_{\mathfrak{p}})^{r \dim W}}.$$

This proposition follows from the next two lemmata. In fact, in the explicit computations, we shall only use lemma 3.2 and the first assertion of lemma 3.4.

Lemma 3.2. — *For any $r > 0$ we consider the set*

$$W^*(\mathcal{O}_{\mathfrak{p}/\mathfrak{p}^r}) = \{x \in (\mathcal{O}_{\mathfrak{p}/\mathfrak{p}^r})^{N+1} - (\mathfrak{p}/\mathfrak{p}^r)^{N+1} \mid \mathbf{f}(x) = 0 \text{ in } (\mathcal{O}_{\mathfrak{p}/\mathfrak{p}^r})^m\}$$

and put $N^*(\mathfrak{p}^r) = \#W^*(\mathcal{O}/\mathfrak{p}^r)$. Then there is an integer $r_0 > 0$ such that

$$\int_{\{x \in \mathcal{O}_{\mathfrak{p}}^{N+1} - \mathfrak{p}^{N+1} \mid f(x) \not\equiv 0\}} \omega_{L,v} = \frac{N^*(\mathfrak{p}^r)}{(\#\mathbf{F}_{\mathfrak{p}})^{r \dim W}}$$

if $r \geq r_0$.

Remark 3.3. — It will follow from the proof that it is in fact sufficient to take r_0 to be

$$2 \inf \{r \in \mathbf{Z}_{>0} \mid \forall x \in \mathcal{O}_{\mathfrak{p}}^{N+1} - \mathfrak{p}^{N+1}, f(x) \equiv 0 \pmod{\mathfrak{p}^r} \Rightarrow (\mathfrak{p}^r)^m \subset \text{Im}(d\mathbf{f}_x)\} + 1.$$

Proof. — For any $r > 0$,

$$\begin{aligned} \int_{\{x \in \mathcal{O}_{\mathfrak{p}}^{N+1} - \mathfrak{p}^{N+1} \mid \mathbf{f}(x) \in \mathfrak{p}\}} \omega_{L, \mathfrak{p}}(x) &= \sum_{x \in (\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^r)^{N+1} - (\mathfrak{p}/\mathfrak{p}^r)^{N+1}} \int_{\{y \in \mathcal{O}_{\mathfrak{p}}^{N+1} \mid \mathbf{f}(y) \in \mathfrak{p} \text{ and } [y]_r = x\}} \omega_{L, \mathfrak{p}}(y) \\ &= \sum_{x \in W^*(\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^r)} \int_{\{y \in \mathcal{O}_{\mathfrak{p}}^{N+1} \mid \mathbf{f}(y) \in \mathfrak{p} \text{ and } [y]_r = x\}} \omega_{L, \mathfrak{p}}(y) \end{aligned}$$

where for any y in $\mathcal{O}_{\mathfrak{p}}^{N+1}$ we denote by $[y]_r$ its class modulo \mathfrak{p}^r . Since V is smooth, the cone W does not intersect the cone defined by the equations

$$\det \left(\frac{\partial f_i}{\partial X_{k_j}} \right)_{1 \leq i, j \leq m} = 0 \quad \text{for } 0 \leq k_1 < \dots < k_m \leq N.$$

Therefore, for r big enough and for any x in $(\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^r)^{N+1} - (\mathfrak{p}/\mathfrak{p}^r)^{N+1}$ such that $\mathbf{f}(x) = 0$ in $(\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^r)^m$ one has that

$$\inf_{(k_j)_j} v_{\mathfrak{p}} \left(\det \left(\frac{\partial f_i}{\partial X_{k_j}} \right)_{1 \leq i, j \leq m} \right)$$

is finite and constant on the class defined by x . Let c be its value. We may assume that $r > c$ and choose a family $0 \leq k_1 < \dots < k_m \leq N$ which realizes this minimum. We may assume that $k_j = N - m + j$. Then if $y \in \mathcal{O}_{\mathfrak{p}}^{N+1}$ represents x and if z belongs to $\mathcal{O}_{\mathfrak{p}}^{N+1}$, one has

$$(3) \quad f_i(y+z) = f_i(y) + \sum_{j=0}^N \frac{\partial f_i}{\partial X_j}(y) z_j + \sum_{j, k=0}^N P_{i, j, k}(y, z) z_j z_k.$$

where the $P_{i, j, k}$ are polynomials in $2N + 2$ variables with coefficients in $\mathcal{O}_{\mathfrak{p}}$. Let L_y be the image of the linear map defined by $\left(\frac{\partial f_i}{\partial X_j}(y) \right)$ on $(\mathcal{O}_{\mathfrak{p}})^{N+1}$, then one has the inclusions

$$(\mathfrak{p}^r)^m \subset (\mathfrak{p}^c)^m \subset L_y \subset (\mathcal{O}_{\mathfrak{p}})^m$$

and $\#((\mathcal{O}_{\mathfrak{p}})^m/L_y) = (\#\mathbf{F}_{\mathfrak{p}})^c$. In particular, for any z in $(\mathfrak{p}^r)^{N+1}$ one has $L_{y+z} = L_y$. We put $L = L_y$. By (3) we have that for any z in $(\mathfrak{p}^r)^{N+1}$,

$$\mathbf{f}(y+z) - \mathbf{f}(y) \in \mathfrak{p}^r L.$$

Therefore, the image of $\mathbf{f}(y)$ in $\mathcal{O}_{\mathfrak{p}}^m/\mathfrak{p}^r L$ depends only on x and we denote it by $\mathbf{f}^*(x)$. If $\mathbf{f}^*(x) \neq 0$ then the set

$$\{u \in \mathcal{O}_{\mathfrak{p}}^{N+1} \mid \mathbf{f}(u) = 0 \text{ and } [u]_r = x\}$$

is empty and the integral is trivial. On the other hand, the set

$$\{u \in (\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^{r+c})^{N+1} \mid \mathbf{f}(u) = 0 \text{ in } (\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^{r+c})^m \text{ and } [u]_r = x\}$$

is also empty. If $\mathbf{f}^*(x) = 0$ then it follows from Hensel's lemma that the coordinates X_0, \dots, X_{N-m} define an isomorphism from

$$\{u \in \mathcal{O}_{\mathfrak{p}}^{N+1} \mid \mathbf{f}(u) = 0 \text{ and } [u]_r = x\}$$

to $(y_0, \dots, y_{N-m}) + (\mathfrak{p}^r)^{N-m+1}$. Therefore, using (2) and the definition of c , we get that

$$\begin{aligned} \int_{\{y \in \mathcal{O}_{\mathfrak{p}}^{N+1} \mid \mathbf{f}(y) = 0 \text{ and } [y]_r = x\}} \omega_{L/\mathfrak{p}}(y) &= \int_{(y_0, \dots, y_{N-m}) + (\mathfrak{p}^r)^{N-m+1}} (\#\mathbf{F}_{\mathfrak{p}})^c du_{0,\mathfrak{p}} \dots du_{N-m,\mathfrak{p}} \\ &= \#\mathbf{F}_{\mathfrak{p}}^{c-r \dim W}. \end{aligned}$$

Let x/\mathfrak{p}^{r+c} be the set

$$\{u \in (\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^{r+c})^{N+1} \mid [u]_r = x\}.$$

Then \mathbf{f} induces a map from x/\mathfrak{p}^{r+c} to $(\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^{r+c})^m$ given by

$$\mathbf{f}([y+z]_{r+c}) = [\mathbf{f}(y)]_{r+c} + \sum_{j=0}^N \frac{\partial \mathbf{f}}{\partial X_j}(y) z_j$$

the image of which is $\mathfrak{p}^r L/(\mathfrak{p}^{r+c})^m$. Therefore, we obtain

$$\begin{aligned} \#\{u \in x/\mathfrak{p}^{r+c} \mid \mathbf{f}(u) = 0 \text{ in } (\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^{r+c})^m\} \\ = (\mathfrak{p}^r L/(\mathfrak{p}^{r+c})^m)^{-1} \times \#(\mathfrak{p}^r/\mathfrak{p}^{r+c})^{N+1} \\ = \#\mathbf{F}_{\mathfrak{p}}^{c+c \dim W} \end{aligned}$$

and

$$\frac{\#\{u \in x/\mathfrak{p}^{r+c} \mid \mathbf{f}(u) = 0 \text{ in } (\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^{r+c})^m\}}{\#\mathbf{F}_{\mathfrak{p}}^{(r+c) \dim W}} = \#\mathbf{F}_{\mathfrak{p}}^{c-r \dim W}.$$

Finally, we get the result. \square

Lemma 3.4. — *With notation as in proposition 3.1, one has*

$$\int_{\{x \in \mathcal{O}_{\mathfrak{p}}^{N+1} - \mathfrak{p}^{N+1} \mid f(x) \not\equiv 0\}} \omega_{L, \mathfrak{p}} = \left(1 - \frac{1}{\#\mathbf{F}_{\mathfrak{p}}^{\delta}}\right) \int_{\{x \in \mathcal{O}_{\mathfrak{p}}^{N+1} \mid f(x)=0\}} \omega_{L, \mathfrak{p}}$$

and

$$\begin{aligned} & \lim_{r \rightarrow +\infty} \frac{N^*(\mathfrak{p}^r)}{(\#\mathbf{F}_{\mathfrak{p}})^{r \dim W}} \\ &= \left(1 - \frac{1}{\#\mathbf{F}_{\mathfrak{p}}^{\delta}}\right) \lim_{r \rightarrow +\infty} \frac{\#\{x \in (\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^r)^{N+1} \mid f(x) = 0 \text{ in } (\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^r)^m\}}{(\#\mathbf{F}_{\mathfrak{p}})^{r \dim W}}. \end{aligned}$$

Proof. — By definition, one has for any λ in $F_{\mathfrak{p}}^*$ the relation

$$\omega_{L, \mathfrak{p}}(\lambda U) = |\lambda|_{\mathfrak{p}}^{\delta} \omega_{L, \mathfrak{p}}(U)$$

which implies the first assertion.

For the second one, let d be the common degree of the f_i . If $r \geq id + 1$, one has the relations

$$\begin{aligned} & \#\{x \in (\mathfrak{p}^i/\mathfrak{p}^r)^{N+1} - (\mathfrak{p}^{i+1}/\mathfrak{p}^r)^{N+1} \mid f(x) \equiv 0 \pmod{\mathfrak{p}^r}\} \\ &= \#\{x \in (\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^{r-i})^{N+1} - (\mathfrak{p}/\mathfrak{p}^{r-i})^{N+1} \mid f(x) \equiv 0 \pmod{\mathfrak{p}^{r-id}}\} \\ &= \#\mathbf{F}_{\mathfrak{p}}^{(N+1)(d-1)i} \#\{x \in (\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^{r-id})^{N+1} - (\mathfrak{p}/\mathfrak{p}^{r-id})^{N+1} \mid f(x) \equiv 0 \pmod{\mathfrak{p}^{r-id}}\}. \end{aligned}$$

Thus we get

$$\begin{aligned} & \#\{x \in (\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^r)^{N+1} \mid f(x) = 0 \text{ in } (\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^r)^m\} \\ &= \sum_{0 \leq i \leq a} \#\mathbf{F}_{\mathfrak{p}}^{(N+1)(d-1)i} N^*(\mathfrak{p}^{r-id}) \\ & \quad + \#\{x \in (\mathfrak{p}^{a+1}/\mathfrak{p}^r)^{N+1} \mid f(x) \equiv 0 \pmod{\mathfrak{p}^r}\} \end{aligned}$$

where $r - r_0 = ad + b$ with $b < d$. We have $a > (r - r_0 - d)/d$ and $N + 1 - md \geq 1$. Thus

$$\#(\mathfrak{p}^{a+1}/\mathfrak{p}^r)^{N+1} \leq \#\mathbf{F}_{\mathfrak{p}}^{(N+1)(r-(r-r_0)/d)} \leq \#\mathbf{F}_{\mathfrak{p}}^{r(N+1-m-1/d)+(N+1)r_0/d}$$

Dividing by $\#\mathbf{F}_p^{r \dim W}$ and using the previous lemma, we get that

$$\frac{\#\{x \in (\mathcal{O}_p/\mathfrak{p}^r)^{N+1} \mid \mathbf{f}(x) = 0 \text{ in } (\mathcal{O}_p/\mathfrak{p}^r)^m\}}{(\#\mathbf{F}_p)^{r \dim W}} = \left(1 - \frac{1}{\#\mathbf{F}_p^\delta}\right)^{-1} \frac{N^*(\mathfrak{p}^r)}{\#\mathbf{F}_p^{r \dim W}} + O(\#\mathbf{F}_p^{-r/d}). \quad \square$$

The equations $(f_i)_{1 \leq i \leq m}$ define an isomorphism

$$\omega_V^{-1} \xrightarrow{\sim} \mathcal{O}_V(\delta).$$

Therefore, for any place v of F the metric on $\mathcal{O}_V(\delta)$ induced by the monomials of degree δ defines a metric $\|\cdot\|_v$ on ω_V^{-1} . The height \mathbf{H} defined by the corresponding metrized line bundle $(\omega_V^{-1}, (\|\cdot\|_v)_{v \in \text{Val}(F)})$ verifies

$$\forall x \in V(F), \quad \mathbf{H}(x) = \prod_{v \in \text{Val}(F)} \sup_{0 \leq i \leq N} (|x_i|_v)^\delta.$$

Corollary 3.5. — *With notations as in proposition 3.1 one has for any finite place \mathfrak{p} of F*

$$\omega_{\mathbf{H}, \mathfrak{p}}(V(F_{\mathfrak{p}})) = \frac{1 - \#\mathbf{F}_p^{-\delta}}{1 - \#\mathbf{F}_p^{-1}} \lim_{r \rightarrow +\infty} \frac{\#\{x \in (\mathcal{O}_p/\mathfrak{p}^r)^{N+1} \mid \mathbf{f}(x) = 0 \text{ in } (\mathcal{O}_p/\mathfrak{p}^r)^m\}}{\#\mathbf{F}_p^{r \dim W}}.$$

Proof. — This follows from proposition 3.1 and [Pe1, lemme 5.4.6]. \square

Remark 3.6. — In particular, a factor $1/3$ is erroneously introduced in the first formula giving the constant $C_{\text{HB}}(V)$ on page 148 in [Pe1] (see also [SD, p. 374]) and therefore a factor 3 is missing in proposition 5.6.1 of [Pe1]. In fact, if V is the cubic surface defined by the equation

$$X_0^3 + X_1^3 + X_2^3 = kX_3^3$$

with $k = 2$ or 3 , one gets the equality

$$\mathfrak{S}_k = \alpha(V)\beta(V)\tau_{\mathbf{H}}(V),$$

where \mathfrak{S}_k is the constant defined by Heath-Brown in [HB]. Therefore, the numerical experiments made by Heath-Brown are compatible with the constant $\theta_{\mathbf{H}}(V)$ as in definition 2.6 and the remark 2.3.2 in [Pe1] has to be corrected accordingly.

4. Points on cubics over \mathbf{F}_p

We now describe explicitly the cardinal of $V(\mathbf{F}_p)$ when V is the diagonal cubic surface given by the equation

$$(4) \quad X_0^3 + q^2 X_1^3 + qr X_2^3 + r^2 X_3^3 = 0$$

where $q, r \in \mathbf{Z}_{>1}$ are squarefree and coprime. We put $K_1 = \mathbf{Q}(q^{1/3})$, $K_2 = \mathbf{Q}(r^{1/3})$ and $K_3 = \mathbf{Q}((qr)^{1/3})$ and consider

$$v_{q,r}(p) = \#\{i \mid p \text{ is totally split in } K_i\}.$$

Proposition 4.1. — *if $p \nmid 3qr$, then*

$$\frac{\#V(\mathbf{F}_p)}{p^2} = \begin{cases} 1 + \frac{1}{p} + \frac{1}{p^2} & \text{if } p \equiv 2 \pmod{3}, \\ 1 + \frac{3v_{q,r}(p)-2}{p} + \frac{1}{p^2} & \text{otherwise.} \end{cases}$$

Remark 4.2. — By a result of Weil (see [Ma2, theorem 23.1]),

$$\frac{\#V(\mathbf{F}_p)}{p^2} = 1 + \text{Tr}(\text{Fr}_p \mid \text{Pic } \overline{V})_p + p^2$$

and the only difficulty is to determine $\text{Tr}(\text{Fr}_p \mid \text{Pic } \overline{V})$. We have chosen to avoid this computation by using a general formula valid for diagonal hypersurfaces.

Remark 4.3. — If $p \equiv 1 \pmod{3}$ then \mathbf{F}_p contains the cubic roots of 1. Therefore $v_{q,r}(p)$ is either 3, 1 or 0. In other words, the possible values in this case are

$$1 + \frac{7}{p} + \frac{1}{p^2}, \quad 1 + \frac{1}{p} + \frac{1}{p^2}, \quad 1 - \frac{2}{p} + \frac{1}{p^2}.$$

Proof. — Let $N(p)$ be the number of solutions of (4) in $(\mathbf{F}_p)^4$. By [IR, §8.7, theorem 5], we have the formula

$$N(p) = p^3 + \sum \overline{\chi}_1(1) \overline{\chi}_2(q^2) \overline{\chi}_3(r^2) \overline{\chi}_4(qr) J_0(\chi_1, \dots, \chi_4)$$

where the sum is taken over the quadruples of nontrivial cubic characters $\chi_1, \chi_2, \chi_3, \chi_4$ from \mathbf{F}_p^* to \mathbf{C}^* such that $\chi_1 \chi_2 \chi_3 \chi_4$ is the trivial character and where

$$J_0(\chi_1, \dots, \chi_4) = \sum_{t_1 + \dots + t_4 = 0} \prod_{i=1}^4 \chi_i(t_i),$$

with the convention $\chi_i(0) = 0$. If $p \equiv 2 \pmod{3}$, then there are no nontrivial characters and we get that

$$\#V(\mathbf{F}_p) = \frac{N(p) - 1}{p - 1} = 1 + p + p^2.$$

Otherwise, there are exactly two nontrivial characters which are conjugate and will be denoted by χ and $\bar{\chi}$. By [IR, §8.5, theorem 4], we have

$$|J_0(\chi \chi \bar{\chi} \bar{\chi})| = p(p - 1).$$

But, by definition, this complex number may be written as

$$\begin{aligned} J_0(\chi \chi \bar{\chi} \bar{\chi}) &= \sum_{t_1 + \dots + t_4 = 0} \chi(t_1 t_2) \bar{\chi}(t_3 t_4) \\ &= \sum_{a \in \mathbf{F}_p} \left| \sum_{t_1 + t_2 = a} \chi(t_1 t_2) \right|^2 \end{aligned}$$

and is a positive real number. Finally we get

$$N(p) = p^3 + p(p - 1) \sum \chi_1(q^2) \chi_2(r^2) \chi_3(qr),$$

where the sum is taken over all nontrivial cubic characters such that $\chi_1 \chi_2 \chi_3$ is nontrivial. This sum may be written as

$$\begin{aligned} \sum \chi_1(q^2) \chi_2(r^2) \chi_3(qr) &= \chi(q^2) \chi(r^2) \bar{\chi}(qr) + \chi(q^2) \bar{\chi}(r^2) \chi(qr) \\ &\quad + \bar{\chi}(q^2) \chi(r^2) \chi(qr) + \chi(q^2) \bar{\chi}(r^2) \bar{\chi}(qr) \\ &\quad + \bar{\chi}(q^2) \chi(r^2) \bar{\chi}(qr) + \bar{\chi}(q^2) \bar{\chi}(r^2) \chi(qr) \\ &= \chi(q) + \bar{\chi}(q) + \chi(r) + \bar{\chi}(r) + \chi(qr) + \bar{\chi}(qr). \end{aligned}$$

Observe that for any integer n prime to p , one has

$$\chi(n) + \bar{\chi}(n) = \begin{cases} -1 & \text{if } p \text{ is not split in } \mathbf{Q}(n^{1/3}), \\ 2 & \text{otherwise. } \square \end{cases}$$

Lemma 4.4. — *With notation as above, if $p \equiv 2 \pmod{3}$, $p \neq 2$, and $p|qr$ then*

$$\frac{N^*(p^t)}{p^{3t}} = 1 - \frac{1}{p} \quad \text{if } t > 0.$$

Proof. — We may assume that $p|r$. Let $\mathbf{x} = (x_0, x_1, x_2, x_3)$ be a solution of the equation (4) in the set $(\mathbf{Z}/p^t \mathbf{Z})^4 - (p\mathbf{Z}/p^t \mathbf{Z})^4$. If $p|x_1$ then by the equation $p|x_0$ and then $\mathbf{x} \in (p)^4$ which gives a contradiction. Since the group of invertible

elements in $\mathbf{Z}/p^t\mathbf{Z}$ is isomorphic to $\mathbf{Z}/p^{t-1}(p-1)\mathbf{Z}$, any element in this group has a unique cubic root. Therefore, the set of solutions is parametrized by the $(x_1, x_2, x_3) \in \mathbf{Z}/p^t\mathbf{Z}$ such that $p \nmid x_1$. \square

Lemma 4.5. — *With notations as above, if $q \equiv \pm r \pmod{9}$ and $3 \nmid qr$, then the possible values for $N^*(3^2)/3^6$ are given by the following table:*

$q, r \pmod{9}$	± 1	± 2	± 4
$N^*(3^2)/3^6$	2	$2^2/3$	$2/3$

Proof. — Up to multiplication by units, the equation in this case may be written over \mathbf{Q}_3 as

$$X^3 + q^2 Y^3 + q^2 Z^3 + q^2 T^3 = 0$$

which is equivalent to

$$X^3 + Y^3 + Z^3 + qT^3 = 0$$

and the result follows from [HB] or a direct computation. \square

5. Convergence factors and residues

As in Heath-Brown [HB], for the explicit computation of the constant we need a family of convergence factors related to zeta functions of cubic extensions of \mathbf{Q} . If V is defined by (4), it follows from [CTKS, p. 12] that $t = \text{rk Pic } V = 1$.

Proposition 5.1. — *If V is the diagonal cubic given by the equation (4) and K_i are the fields defined in the previous paragraph, then the measure $\omega_{\mathbf{H}}$ coincides with the measure*

$$\lim_{s \rightarrow 1} (s-1) \frac{\prod_{i=1}^3 \zeta_{K_i}(s)}{\zeta_{\mathbf{Q}}(s)^2} \prod_{v \in \text{Val}(\mathbf{Q})} \lambda'_v \omega_{\mathbf{H},v}.$$

where

$$\lambda'_p = \frac{\prod_{i=1}^3 \prod_{\{\mathfrak{P} \in \text{Val}(K_i) \mid \mathfrak{P} \mid p\}} (1 - \#\mathbf{F}_{\mathfrak{P}}^{-1})}{(1-p^{-1})^2} \quad \text{if } p \text{ is a prime number and } \lambda'_{\mathbf{R}} = 1.$$

Remark 5.2. — If p does not divide $3qr$, we can use the term $\lambda'_p \omega_{\mathbf{H},p}(V(\mathbf{Q}_p))$ which by lemma 3.2, lemma 3.4, and lemma 5.4.6 in [Pe1] coincides with

$\lambda'_p \#V(\mathbf{F}_p)/p^2$ (see also [Pe1, lemme 2.2.1]) and by proposition 4.1 is equal to

$$\begin{aligned} & \left(1 - \frac{1}{p}\right)^7 \left(1 + \frac{7}{p} + \frac{1}{p^2}\right) && \text{if } p \equiv 1 \pmod{3} \text{ and } v_{q,r}(p) = 3 \\ & \left(1 - \frac{1}{p}\right) \left(1 - \frac{1}{p^3}\right)^2 \left(1 + \frac{1}{p} + \frac{1}{p^2}\right) && \text{if } p \equiv 1 \pmod{3} \text{ and } v_{q,r}(p) = 1 \\ & \left(1 - \frac{1}{p}\right) \left(1 + \frac{1}{p} + \frac{1}{p^2}\right)^3 \left(1 - \frac{2}{p} + \frac{1}{p^2}\right) && \text{if } p \equiv 1 \pmod{3} \text{ and } v_{q,r}(p) = 0 \\ & \left(1 - \frac{1}{p}\right) \left(1 - \frac{1}{p^2}\right)^3 \left(1 + \frac{1}{p} + \frac{1}{p^2}\right) && \text{if } p \equiv 2 \pmod{3} \end{aligned}$$

and the good places yield a product $C_1 C_2 C_3$ where

$$\begin{aligned} C_1 &= \prod_{\substack{p \nmid 3qr \\ p \equiv 1 \pmod{3} \\ v_{q,r}(p) = 3}} \left(1 - \frac{1}{p}\right)^7 \left(1 + \frac{7}{p} + \frac{1}{p^2}\right) \\ C_2 &= \prod_{\substack{p \nmid 3qr \\ p \equiv 1 \pmod{3} \\ v_{q,r}(p) \neq 3}} \left(1 - \frac{1}{p^3}\right)^3 \\ C_3 &= \prod_{\substack{p \nmid 3qr \\ p \equiv 2 \pmod{3}}} \left(1 - \frac{1}{p^3}\right) \left(1 - \frac{1}{p^2}\right)^3 \end{aligned}$$

Proof. — It follows from the proof of proposition 4.1 that, if $p \nmid 3qr$,

$$(5) \quad \frac{\#V(\mathbf{F}_p)}{p^2} = 1 + \frac{a(p)}{p} + \frac{1}{p^2}$$

where

$$(6) \quad a(p) = \begin{cases} 1 & \text{if } p \equiv 2 \pmod{3}, \\ 1 + \chi(q) + \bar{\chi}(q) + \chi(r) + \bar{\chi}(r) + \chi(qr) + \bar{\chi}(qr) & \text{otherwise,} \end{cases}$$

where χ is a nontrivial cubic character of \mathbf{F}_p , if $p \equiv 1 \pmod{3}$. It follows from a theorem of Weil (see [Ma2, theorem 23.1]) that, outside a finite set of places,

$$\#V(\mathbf{F}_p) = 1 + \text{Tr}(\text{Fr}_p | \text{Pic } \bar{V})_p + p^2.$$

Since the action of the Galois group on the Picard group splits over a finite extension of k , the eigenvalues of the Frobenius map for this action are roots of unity. Therefore we get that

$$(7) \quad L_p(s, \text{Pic}(\overline{V}))^{-1} = 1 - \frac{a(p)}{p^s} + R_p \left(\frac{1}{p^s} \right)$$

where the R_p are polynomials of order at least 2 with uniformly bounded coefficients with respect to p .

But for any cubefree integer n , the zeta function of $\mathbf{Q}(n^{1/3})$ may be written as an Euler product with local terms $\zeta_{\mathbf{Q}(n^{1/3}),p}(s)$ which for $p \nmid 3n$ are given by

$$\zeta_{\mathbf{Q}(n^{1/3}),p}(s)^{-1} = \begin{cases} \left(1 - \frac{1}{p^s}\right) \left(1 - \frac{1}{p^{2s}}\right) & \text{if } p \equiv 2 \pmod{3} \\ \left(1 - \frac{1}{p^s}\right) \left(1 - \frac{\chi(n)}{p^s}\right) \left(1 - \frac{\bar{\chi}(n)}{p^s}\right) & \text{if } p \equiv 1 \pmod{3} \end{cases}$$

Thus, by (6), for almost all places, the local terms of the zeta functions verify

$$\left(\zeta_{\mathbf{Q},p}(s) \prod_{i=1}^3 \frac{\zeta_{K_i,p}(s)}{\zeta_{\mathbf{Q},p}(s)} \right)^{-1} = 1 - \frac{a(p)}{p^s} + Q_p \left(\frac{1}{p^s} \right)$$

where the Q_p are polynomials of order at least 2 with bounded coefficients. Using (5) we get that the product of measures given in the proposition defines a Borel measure on the adelic points of V and by (7) that the Euler product defining the quotient

$$L_S(s, \text{Pic}(\overline{V})) \Big/ \frac{\prod_{i=1}^3 \zeta_{K_i}(s)}{\zeta_{\mathbf{Q}}(s)^2}$$

converges absolutely in $s = 1$. Since $d_{\mathbf{Q}} = 1$, we get that

$$\begin{aligned}
 \omega_{\mathbf{H}} &= \lim_{s \rightarrow 1} \left((s-1)L_S(s, \text{Pic } \bar{V}) \right) \prod_{v \in \text{Val}(\mathbf{Q})} \lambda_v^{-1} \omega_{\mathbf{H},v} \\
 &= \lim_{s \rightarrow 1} \left((s-1)L_S(s, \text{Pic } \bar{V}) \right) \prod_{v \in \text{Val}(\mathbf{Q})} \frac{\lambda_v^{-1}}{\lambda'_v} \prod_{v \in \text{Val}(\mathbf{Q})} \lambda'_v \omega_{\mathbf{H},v} \\
 &= \lim_{s \rightarrow 1} \left((s-1)L_S(s, \text{Pic } \bar{V}) \right) \lim_{s \rightarrow 1} \frac{\left(\prod_{i=1}^3 \zeta_{K_i}(s) \right) / \zeta_{\mathbf{Q}}(s)^2}{L_S(s, \text{Pic } \bar{V})} \prod_{v \in \text{Val}(\mathbf{Q})} \lambda'_v \omega_{\mathbf{H},v} \\
 &= \lim_{s \rightarrow 1} (s-1) \frac{\prod_{i=1}^3 \zeta_{K_i}(s)}{\zeta_{\mathbf{Q}}(s)^2} \prod_{v \in \text{Val}(\mathbf{Q})} \lambda'_v \omega_{\mathbf{H},v}. \quad \square
 \end{aligned}$$

6. Brauer-Manin obstruction to weak approximation

In this paragraph, using the work of Colliot-Thélène, Kanevsky and Sansuc [CTKS], we shall compute the quotient

$$(8) \quad \omega_{\mathbf{H}}(V(\mathbf{A}_{\mathbf{Q}})^{\text{Br}}) / \omega_{\mathbf{H}}(V(\mathbf{A}_{\mathbf{Q}}))$$

when V is the diagonal cubic defined by the equation (4) and $V(\mathbf{A}_{\mathbf{Q}}) \neq \emptyset$.

In the following, we assume that q and r are distinct prime numbers such that $3 \nmid qr$. It follows from [CTKS, p. 28] that $V(\mathbf{A}_{\mathbf{Q}}) \neq \emptyset$ if and only if the following condition is satisfied

$$(9) \quad (q \equiv 2 \pmod{3} \text{ or } r \in \mathbf{F}_q^{*3}) \text{ and } (r \equiv 2 \pmod{3} \text{ or } q \in \mathbf{F}_r^{*3})$$

Proposition 6.1. — *Under these assumptions, the value for the quotient (8) depends only on the classes of p and q modulo 9. These values are given in the following table:*

$r \backslash q$	1	2	4	5	7	8
1	1	1	1	1	1	1
2	1	$1/3$	0	0	$1/3$	1
4	1	0	$1/3$	$1/3$	0	1
5	1	0	$1/3$	$1/3$	0	1
7	1	$1/3$	0	0	$1/3$	1
8	1	1	1	1	1	1

Proof. — Let j be a primitive third root of unity, $k = \mathbf{Q}(j)$ and $K = k(q^{1/3}, r^{1/3})$. We have the following diagram of fields

$$\begin{array}{c}
 K \\
 \swarrow \quad | \quad \searrow \\
 k(q^{1/3}) \quad k((qr)^{1/3}) \quad k(r^{1/3}) \\
 \swarrow \quad | \quad \searrow \\
 k \\
 | \\
 \mathbf{Z}/2\mathbf{Z} \\
 | \\
 \mathbf{Q}
 \end{array}$$

and the group $G = \text{Gal}(K/\mathbf{Q})$ may be described as the semi-direct product $(\mathbf{Z}/3\mathbf{Z})^2 \rtimes \mathbf{Z}/2\mathbf{Z}$ where $\mathbf{Z}/2\mathbf{Z}$ acts by $-\text{Id}$ on $(\mathbf{Z}/3\mathbf{Z})^2$. By [CTKS, proposition 1, p. 7], we have that

$$\text{Br}(V \times_{\mathbf{Q}} k) / \text{Br}(k) \xrightarrow{\sim} H^1(k, \text{Pic}(\overline{V})) \xrightarrow{\sim} \mathbf{Z}/3\mathbf{Z}.$$

But the Hochschild-Serre spectral sequence gives an exact sequence

$$\begin{aligned}
 0 \rightarrow H^1(\mathbf{Z}/2\mathbf{Z}, (\text{Pic}(\overline{V}))^{(\mathbf{Z}/3\mathbf{Z})^2}) &\rightarrow H^1(\mathbf{Q}, \text{Pic}(\overline{V})) \\
 &\rightarrow (H^1(k, \text{Pic}(\overline{V})))^{\mathbf{Z}/2\mathbf{Z}} \rightarrow H^2(\mathbf{Z}/2\mathbf{Z}, (\text{Pic}(\overline{V}))^{(\mathbf{Z}/3\mathbf{Z})^2}).
 \end{aligned}$$

By [CTKS, p. 12], $\text{Pic}(\overline{V})^{\text{Gal}(\overline{k}/k)} = \mathbf{Z}$ and, since a map from a group killed by 3 to one killed by 2 is trivial, we obtain an isomorphism

$$H^1(\mathbf{Q}, \text{Pic}(\overline{V})) \xrightarrow{\sim} H^1(k, \text{Pic}(\overline{V}))^{\mathbf{Z}/2\mathbf{Z}}.$$

By the proof of [CTKS, lemme 3, p. 13], we have isomorphisms

$$H^1(k, \text{Pic}(\overline{V})) \xrightarrow{\sim} H^1(\mathbf{Z}/3\mathbf{Z}, \mathbf{Z} \oplus Q) \xrightarrow{\sim} H^1(\mathbf{Z}/3\mathbf{Z}, Q)$$

where Q is the $\mathbf{Z}/3\mathbf{Z}$ -module defined by the exact sequence

$$0 \rightarrow \mathbf{Z} \xrightarrow{N} \mathbf{Z}[\mathbf{Z}/3\mathbf{Z}] \rightarrow Q \rightarrow 0.$$

But for any cyclic group C and any C -module M there is canonical injection

$$H^1(C, M) \rightarrow \text{Hom}(C, M_C)$$

where M_C denotes the group of coinvariants and which sends the class of a cocycle γ onto the induced morphism $\bar{\gamma}$ from C to M_C , so that the diagram

$$\begin{array}{ccc} C & \xrightarrow{\gamma} & M \\ \parallel & & \downarrow \\ C & \xrightarrow{\bar{\gamma}} & M_C \end{array}$$

commutes. If C is a normal subgroup of a group H then the above morphism is compatible with the natural actions of H/C on $H^1(C, M)$ and $\text{Hom}(C, M_C)$. It follows from [CTKS, lemme 3] that $\mathbf{Z}/2\mathbf{Z}$ acts by $-\text{Id}$ on the torsion part of $Q_{\mathbf{Z}/3\mathbf{Z}}$ and by the above description of G as a semi-direct product, $\mathbf{Z}/2\mathbf{Z}$ acts by -1 on $\mathbf{Z}/3\mathbf{Z}$. Therefore the action of $\mathbf{Z}/2\mathbf{Z}$ on $H^1(\mathbf{Z}/3\mathbf{Z}, Q)$ is trivial and

$$H^1(\mathbf{Q}, \text{Pic}(\bar{V})) = \mathbf{Z}/3\mathbf{Z}.$$

For any prime p , the canonical pairing

$$\begin{aligned} \text{Br}(V) \times V(\mathbf{Q}_p) &\rightarrow \mathbf{Q}/\mathbf{Z} \\ ([A], P) &\mapsto \text{inv}_p(A(P)) \end{aligned}$$

defines an equivalence relation on $V(\mathbf{Q}_p)$ which is also denoted by Br . If V has good reduction in p , then by [CT1, theorem A (iii)] the rational equivalence on $V(\mathbf{Q}_p)$ is trivial and therefore the same is true of the Brauer equivalence. The condition (9) implies that if $p|qr$ then V is rational over \mathbf{Q}_p (see [CTKS, lemme 8, p. 41]) and $\#V(\mathbf{Q}_p)/\text{Br} = 1$. Using the same type of argument we get that $\#V(\mathbf{Q}_3)/\text{Br} = 1$ if q, r or qr is a cube modulo 9, that is one of them is 1 or -1 modulo 9. On the other hand, it follows from [CTKS, §5, p. 41] that this cardinal is 3 at 3 otherwise. This explains the dichotomy between integral and nonintegral values in the table.

We have to find an element $A \in \text{Br}(V)$ whose class spans $\text{Br}(V)/\text{Br}(\mathbf{Q})$. Then the Brauer equivalence and the Brauer obstruction may be computed in terms of the functions

$$\begin{aligned} i_p : V(\mathbf{Q}_p) &\rightarrow \mathbf{Z}/3\mathbf{Z} \\ P &\mapsto \text{inv}_p(A(P)). \end{aligned}$$

This is an intricate procedure. It is described in detail on pages 19-72 and summarized in an algorithm on pages 73-79 in [CTKS], which we are going to follow.

Let us first assume that q, r or qr is a cube modulo 9. Then all functions i_p are constant and A may be chosen so that the value i_p of the function i_p is trivial

except when p equals 3, q , or r . It remains to compute the constant values i_3 , i_q , and i_r for such an A . To this end, we use the additive norm rest symbols $[\cdot, \cdot]_p$ from k_v^* to $\mathbf{Z}/3\mathbf{Z}$, for v in $\text{Val}(k)$, dividing p (see [CTKS, p. 77]). They are biadditive, anticommutative and verify the relations

$$[j, p]_p = \frac{p-1}{3} \quad \text{if } p \equiv 1 \pmod{3}$$

$$[j, p]_p = -\frac{p^2-1}{3} \quad \text{if } p \equiv 2 \pmod{3}.$$

If r is a cube in \mathbf{Q}_3 , we get that A may be chosen so that

$$i_3 = 0, \quad i_q = 0, \quad i_r = [j, r]_r.$$

Since $r \equiv \pm 1 \pmod{9}$, we have that $i_r = 0$ and the Brauer-Manin obstruction is trivial. If q is a cube modulo 9 the result is similar and the quotient (8) equals 1.

If qr is a cube modulo 9 but q and r are not, then the situation is similar to the one considered in [CTKS, proposition 5, p. 67–69], and one writes the equation as

$$X^3 + qY^3 + q^2rZ^3 + q^4r^2T^3 = 0$$

and, by [CTKS, §8], A may be chosen so that

$$i_3 = 0, \quad i_q = 0 \quad \text{and} \quad i_r = [j, r]_r.$$

The values of i_r are given by the following table (see [CTKS, p. 78])

$r \pmod{9}$	2	4	5	7
$[j, r]_r$	2	1	1	2

and in this case there is a Brauer-Manin obstruction to the Hasse principle and the quotient (8) is zero.

If none of the q , r and qr is a cube in \mathbf{Q}_3 , then to complete the proof of proposition 6.1 we have only to prove that each class in $V(\mathbf{Q}_3)$ for the Brauer equivalence has the same volume for $\omega_{\mathbf{H},3}$.

Up to permutation and change of sign, we may assume that $q \equiv r \equiv 2 \pmod{9}$ or $q \equiv r \equiv 4 \pmod{9}$. Therefore the equation modulo 9 may be written as

$$X^3 + 4Y^3 + 4Z^3 + 4T^3 = 0$$

or

$$X^3 + 16Y^3 + 16Z^3 + 16T^3 = 0.$$

Therefore on \mathbf{Q}_3 the equation is equivalent to

$$X^3 + Y^3 + Z^3 + 2T^3 = 0$$

or

$$X^3 + Y^3 + Z^3 + 4T^3 = 0$$

A direct computation modulo 9 shows that exactly one of the three first coordinates has to be divisible by 3. But in the first case, it follows from [HB, proof of theorem 1] that the classes for the Brauer equivalence are determined by the coordinate which vanishes modulo 3. By a symmetry argument the volumes of the equivalence classes are the same.

By [CTKS, p. 49], there is a choice of \mathcal{A} so that the induced map i_3 over $V(\mathbf{Q}_3(j))$ is given as

$$(10) \quad i_3(x, y, z, t) = -[x + jy, v]_3,$$

where v is the coefficient of T^3 . Using the commutative diagram

$$\begin{array}{ccc} V(\mathbf{Q}_3) & \xrightarrow{i_3} & \mathbf{Z}/3\mathbf{Z} \\ \downarrow & & \downarrow \times 2 \\ V(\mathbf{Q}_3(j)) & \xrightarrow{i_3} & \mathbf{Z}/3\mathbf{Z} \end{array}$$

we get that the invariants for the second equation may be described, after reduction modulo 9, as the double of the ones for the first and we get a similar description of $V(\mathbf{Q}_3)/\text{Br}$. This implies the result in this case. \square

7. Numerical tests

The numerical tests for the number of points with bounded heights have been conducted using an efficient program of Dan Bernstein [Be]. We considered the following cubic surfaces


$$(S_1) \quad X_0^3 + 17^2 X_1^3 + 17 \times 53 X_2^3 + 53^2 X_3^3 = 0$$

$$(S_2) \quad X_0^3 + 71^2 X_1^3 + 71 \times 53 X_2^3 + 53^2 X_3^3 = 0$$

$$(S_3) \quad X_0^3 + 5^2 X_1^3 + 23 \times 5 X_2^3 + 23^2 X_3^3 = 0$$

$$(S_4) \quad X_0^3 + 11^2 X_1^3 + 29 \times 11 X_2^3 + 29^2 X_3^3 = 0$$

One can take for the open set U the whole surface V , as there are no rational points on the exceptional curves. The graphs of $n_{U, \mathbf{H}}$ are presented below.



Let us sum up the description of the theoretical constant. Let V be a diagonal cubic over \mathbf{Q} defined by the equation (4) with q and r distinct prime numbers such that $3 \nmid qr$ and such that the condition (9) is satisfied.

By proposition 5.1, the constant $\theta_{\mathbf{H}}(V)$ may be written as the product

$$\frac{\omega_{\mathbf{H}}(V(A_{\mathbf{Q}})^{\text{Br}})}{\omega_{\mathbf{H}}(V(A_{\mathbf{Q}}))} \#H^1(\mathbf{Q}, \text{Pic}(\overline{V})) \\ \times \prod_{i=1}^3 \zeta_{K_i}^*(1) \prod_{p|3qr} \lambda'_p \omega_{\mathbf{H},p}(V(\mathbf{Q}_p)) C_1 C_2 C_3 \omega_{\mathbf{H},\mathbf{R}}(V(\mathbf{R}))$$

where the first term which will be denoted by C_{Br} may be found in proposition 6.1, the cardinal of $H^1(\mathbf{Q}, \text{Pic}(\overline{V}))$ equals 3, the residues of the zeta functions $\zeta_{K_i}^*(1)$ have been computed using Dirichlet's class number formula and the system PARI (see also [Co, chapter 4]), λ'_p is defined in proposition 5.1, the volumes at the bad places are given in lemmata 4.4 and 4.5 and C_1, C_2, C_3 have been described as absolutely convergent Euler products (see remark 5.2). The volume at the real place may be computed directly using the definition of the Leray form.

The computations are summarized in the following table:

Surface	S_1	S_2	S_3	S_4
H	99999	99999	99999	99999
$n_{U, \mathbf{H}}(H)$	3773	1696	2353	2904
C_{Br}	1	1	1/3	1/3
$H^1(\mathbf{Q}, \text{Pic}(\overline{V}))$	3	3	3	3
$\zeta_{\mathbf{Q}(q^{1/3})}^*(1)$	1.4680	1.8172	1.1637	1.2284
$\zeta_{\mathbf{Q}(r^{1/3})}^*(1)$	1.8172	2.2035	1.1879	1.6792
$\zeta_{\mathbf{Q}((qr)^{1/3})}^*(1)$	1.9342	1.9925	1.0865	1.0543
$\lambda'_3 \omega_{\mathbf{H}}(V(\mathbf{Q}_3))$	0.5926	0.5926	0.6667	1.3333
$\lambda'_q \omega_{\mathbf{H}}(V(\mathbf{Q}_q))$	0.9379	0.9808	0.7680	0.9016
$\lambda'_r \omega_{\mathbf{H}}(V(\mathbf{Q}_r))$	0.9808	0.9857	0.9547	0.9644
C_1	0.9978	0.9989	0.9973	0.9812
C_2	0.9892	0.9892	0.9892	0.9893
C_3	0.3103	0.3072	0.3514	0.3158
$\omega_{\mathbf{H}}(V(\mathbf{R}))$	0.0148	0.0042	0.0917	0.0389
$\theta_{\mathbf{H}}(V)$	0.0382	0.0175	0.0234	0.0300
$n_{U, \mathbf{H}}(H)/\theta_{\mathbf{H}}(V)H$	0.9882	0.9669	1.0077	0.9665


The new program of Bernstein allows to increase the upper bound for the height of rational points on the cubic surfaces studied by Heath-Brown in [HB].

These cubics are defined by the equations

$$(S_5) \quad X_0^3 + X_1^3 + X_2^3 + 2X_3^3 = 0$$

$$(S_6) \quad X_0^3 + X_1^3 + X_2^3 + 3X_3^3 = 0.$$

The graphs of $n_{U, \mathbf{H}}$ are drawn below.



In this case, by remark 3.6 and [HB], the constant $\theta_{\mathbf{H}}(V)$ may be written as the product

$$\frac{\omega_{\mathbf{H}}(V(A_{\mathbf{Q}})^{\text{Br}})}{\omega_{\mathbf{H}}(V(A_{\mathbf{Q}}))} \#H^1(\mathbf{Q}, \text{Pic}(\overline{V})) \times \zeta_{\mathbf{Q}(q^{1/3})}^*(1)^3 \prod_{p|3q} \lambda'_p \omega_{\mathbf{H},p}(V(\mathbf{Q}_p)) C_1 C_2 C_3 \omega_{\mathbf{H},\mathbf{R}}(V(\mathbf{R}))$$

where the first factor is $1/3$ for these cubics, the second 3 , q is the last coefficient in the equation, the local factors at the bad places are given by

$$N^*(3^2) = \begin{cases} 2^2 3^5 & \text{if } q = 2 \\ 2^3 3^4 & \text{if } q = 3 \end{cases} \quad N^*(4) = 2^6 - 2^4 \quad \text{if } q = 2$$

and C_1 , C_2 and C_3 are defined exactly as in remark 5.2.

The results are given in the following table:

Surface	S_5	S_6
H	99999	99999
$n_{U, \mathbf{H}}(H)$	205431	115582
C_{Br}	1/3	1/3
$H^1(\mathbf{Q}, \text{Pic}(\bar{V}))$	3	3
$\zeta_{\mathbf{Q}(q^{1/3})}^*(1)$	0.814624	1.017615
$\lambda'_3 \omega_{\mathbf{H}}(V(\mathbf{Q}_3))$	1.333333	0.888889
$\lambda'_q \omega_{\mathbf{H}}(V(\mathbf{Q}_q))$	0.750000	1.000000
C_1	0.954038	0.976203
C_2	0.989387	0.989279
C_3	0.830682	0.306638
$\omega_{\mathbf{H}}(V(\mathbf{R}))$	4.921515	4.295619
$\theta_{\mathbf{H}}(V)$	2.086108	1.191539
$n_{U, \mathbf{H}}(H)/\theta_{\mathbf{H}}(V)H$	0.984767	0.970032

Therefore, the numerical tests for these cubic surfaces are compatible with an asymptotic behavior of the form

$$n_{U, \mathbf{H}}(H) \sim \theta_{\mathbf{H}}(V)H \quad \text{when } H \rightarrow +\infty.$$

Acknowledgements This work was done while both authors were enjoying the hospitality of the Newton Institute for Mathematical Sciences in Cambridge. We have benefited from conversations with Colliot-Thélène, Heath-Brown, Salberger and Swinnerton-Dyer. We thank the referee for his useful remarks and suggestions.

The second author was partially supported by NSA Grant MDA904-98-1-0023 and by EPSRC Grant K99015.

References

- [BM] V. V. Batyrev and Y. I. Manin, *Sur le nombre des points rationnels de hauteur bornée des variétés algébriques*, Math. Ann. **286** (1990), 27–43.
- [BT1] V. V. Batyrev and Y. Tschinkel, *Rational points of bounded height on compactifications of anisotropic tori*, Internat. Math. Res. Notices **12** (1995), 591–635.
- [BT2] ———, *Rational points on some Fano cubic bundles*, C. R. Acad. Sci. Paris Sér. I Math. **323** (1996), n° 1, 41–46.
- [BT3] ———, *Tamagawa numbers of polarized algebraic varieties*, Nombre et répartition de points de hauteur bornée, Astérisque, vol. 251, SMF, Paris, 1998, pp. 299–340.
- [Be] D. J. Bernstein, *Enumerating solutions to $p(a) + q(b) = r(c) + s(d)$* , to appear in Math. Comp. (1999).
- [Co] H. Cohen, *A course in computational algebraic number theory*, Graduate Texts in Math., vol. 138, Springer-Verlag, Berlin, Heidelberg and New York, 1993.
- [CT1] J.-L. Colliot-Thélène, *Hilbert’s theorem 90 for K_2 , with application to the Chow groups of rational surfaces*, Invent. Math. **71** (1983), 1–20.
- [CT2] ———, *The Hasse principle in a pencil of algebraic varieties*, Number theory (Tiruchirapalli, 1996), Contemp. Math., vol. 210, Amer. Math. Soc., Providence, 1998, pp. 19–39.
- [CTKS] J.-L. Colliot-Thélène, D. Kanevsky, et J.-J. Sansuc, *Arithmétique des surfaces cubiques diagonales*, Diophantine approximation and transcendence theory (Bonn, 1985), Lecture Notes in Math., vol. 1290, Springer-Verlag, Berlin, 1987, pp. 1–108.
- [CTS] J.-L. Colliot-Thélène et J.-J. Sansuc, *La descente sur une variété rationnelle définie sur un corps de nombres*, C. R. Acad. Sci. Paris Sér. A **284** (1977), 1215–1218.
- [FMT] J. Franke, Y. I. Manin, and Y. Tschinkel, *Rational points of bounded height on Fano varieties*, Invent. Math. **95** (1989), 421–435.
- [HB] D. R. Heath-Brown, *The density of zeros of forms for which weak approximation fails*, Math. Comp. **59** (1992), 613–623.
- [IR] K. Ireland and M. Rosen, *A classical introduction to modern number theory (second edition)*, Graduate texts in Math., vol. 84, Springer-Verlag, Berlin, Heidelberg and New York, 1990.
- [Lac] G. Lachaud, *Une présentation adélique de la série singulière et du problème de Waring*, Enseign. Math. (2) **28** (1982), 139–169.
- [Ma1] Y. I. Manin, *Le groupe de Brauer-Grothendieck en géométrie diophantienne*, Actes du congrès international des mathématiciens, Tome 1 (Nice, 1970), Gauthiers-Villars, Paris, 1971, pp. 401–411.

- [Ma2] ———, *Cubic forms (second edition)*, North-Holland Math. Library, vol. 4, North-Holland, Amsterdam, 1986.
- [Pe1] E. Peyre, *Hauteurs et mesures de Tamagawa sur les variétés de Fano*, Duke Math. J. **79** (1995), n° 1, 101–218.
- [Pe2] ———, *Terme principal de la fonction zêta des hauteurs et torseurs universels*, Nombre et répartition de points de hauteur bornée, Astérisque, vol. 251, SMF, Paris, 1998, pp. 259–298.
- [Sal] P. Salberger, *Tamagawa measures on universal torsors and points of bounded height on Fano varieties*, Nombre et répartition de points de hauteur bornée, Astérisque, vol. 251, SMF, Paris, 1998, pp. 91–258.
- [SaSk] P. Salberger and A. N. Skorobogatov, *Weak approximation for surfaces defined by two quadratic forms*, Duke Math. J. **63** (1991), n° 2, 517–536.
- [SD] H. P. F. Swinnerton-Dyer, *Counting rational points on cubic surfaces*, Classification of algebraic varieties (L'Aquila, 1992) (C. Ciliberto, E. L. Livorni, and A. J. Sommese, eds.), Contemp. Math., vol. 162, AMS, Providence, 1994, pp. 371–379.

October 5, 2022

EMMANUEL PEYRE, Institut de Recherche Mathématique Avancée, Université Louis Pasteur
et C.N.R.S., 7 rue René-Descartes, 67084 Strasbourg, France
E-mail: peyre@irma.u-strasbg.fr

YURI TSCHINKEL, Department of mathematics, University of Illinois in Chicago, 851 South
Morgan Street, Chicago IL 60607-7045, USA • *E-mail*: yuri@math.uic.edu