
COUNTING POINTS ON VARIETIES USING UNIVERSAL TORSORS

by

Emmanuel Peyre

Abstract. — Around 1989, Manin initiated a program to understand the asymptotic behaviour of rational points of bounded height on Fano varieties. This program led to the search of new methods to estimate the number of points of bounded height on various classes of varieties. Methods based on harmonic analysis were successful for compactifications of homogeneous spaces. However, they do not apply to other types of varieties. Universal torsors, introduced by Colliot-Thélène and Sansuc in connection with the Hasse principle and weak approximation, turned out to be a useful tool in the treatment of other varieties. The aim of this short survey is to describe the use of torsors in various representative examples.

Résumé. — Vers 1989, Manin initia un programme en vue de comprendre le comportement asymptotique des points rationnels de hauteur bornée sur les variétés de Fano. Ce programme amena à la recherche de nouvelles méthodes pour estimer le nombre de points de hauteur bornée sur différentes classes de variétés. Les méthodes basées sur l'analyse harmonique s'appliquent avec succès pour les compactifications d'espaces homogènes. Cependant, elles ne s'appliquent pas à d'autres types de variétés. Les toreseurs universels, introduits par Colliot-Thélène et Sansuc en relation avec le principe de Hasse et l'approximation faible se sont révélés utiles pour l'étude d'autres variétés. L'objet de cet article est de décrire l'utilisation de ces toreseurs pour divers exemples représentatifs.

1. Introduction

If the rational points of a variety V over a number field k are Zariski dense, it is natural to equip V with a height H and to study the asymptotic distribution of

Key words and phrases. — Rational points, heights, universal torsors.

the set of points of bounded height on V . In **[FMT]** and **[BM]**, Batyrev, Franke, Manin and Tschinkel gave strong evidence supporting conjectures relating the asymptotic behaviour of the number of points of bounded height on open subsets of V to geometrical invariants of V . This work motivated the development of several methods to estimate the number of points of bounded height for new classes of varieties. One of the most successful methods was the use of harmonic analysis on adelic groups. For example, it was used by Batyrev and Tschinkel in **[BT1]**, **[BT2]**, and **[BT4]** to handle the case of projective toric varieties, by Strauch and Tschinkel in **[ST1]** and **[ST2]** for toric bundles over flag varieties, and by Chambert-Loir and Tschinkel in **[CLT1]**, **[CLT2]**, and **[CLT3]** for equivariant compactifications of vector spaces. However, these kind of methods apply only to equivariant compactifications of homogeneous spaces. One may say that almost all other methods have one step in common, namely the lifting to universal torsors. Universal torsors have been introduced by Colliot-Thélène and Sansuc in **[CTS1]**, **[CTS2]**, **[CTS3]**, and **[CTS4]** to study the Hasse principle and the weak approximation. The interest of universal torsors is that, from an arithmetic point of view, these torsors should be much simpler than the variety itself. As an example, universal torsors over smooth projective toric varieties are open subsets of an affine space. When the Fano variety V is a smooth complete intersection of dimension bigger than three in the projective space, the universal torsor may be described as the cone over the variety. In that case, if the dimension of the variety is big enough, the conjectural formula of Manin may be deduced from the formula given by the classical circle method. This reduction, which is described in **[FMT]**, may be seen as a particular case of the lifting to the universal torsor. Salberger in **[Sal]** was the first to use explicitly universal torsors in relation with points of bounded height. In particular, he was able to give a new proof of the theorem of Batyrev and Tschinkel for smooth projective split toric varieties over \mathbf{Q} . This lifting to the universal torsor was then used by de la Bretèche in **[Bre1]** to give a better estimate for the number of points of bounded height on toric varieties. The lifting to universal torsors was later used by Salberger and de la Bretèche (see **[Bre2]**) to prove the asymptotic formula for the plane blown up in four points over \mathbf{Q} . In a more general setting, the author described in **[Pe2]** and **[Pe3]** how the conjectural asymptotic formula lifts naturally to universal torsors.

The aim of this short survey is to explain in a self-contained way the usefulness of universal torsors for counting points of bounded height. In Section 2, we describe the heights used throughout the paper, and in Section 3 we recall the empirical formula for the number of points on Fano varieties. In Section 4 we

give a short list of cases for which this formula holds, in Section 5 we describe the counter-example of Batyrev and Tschinkel. In Section 6 we describe briefly both methods: harmonic analysis and universal torsors. Section 7 is devoted to the case of a hypersurface in \mathbf{P}^n . The next section contains the definition of universal torsors in general. In Section 9 we describe Cox's construction of universal torsors for toric varieties and explain how Salberger used it and in Section 10 we turn to the case of the plane blown up in four points, in which case the universal torsor was described by Salberger and Skorobogatov. The last section contains a short description of the generalization of these lifting arguments to a larger class of varieties.

Acknowledgements: I am very grateful to John Voight who typed the notes of my talks at the American Institute of Mathematics on which this survey is based.

2. Heights on projective varieties

Definition 2.1. — The classical exponential height on the projective space over \mathbf{Q} is defined as follows:

$$H_N: \mathbf{P}^N(\mathbf{Q}) \rightarrow \mathbf{R}_{>0}$$

$$(x_0 : \dots : x_N) \mapsto \sup_{0 \leq i \leq N} |x_i|, \quad \text{if } \begin{cases} x_i \in \mathbf{Z}, \text{ and} \\ \gcd(x_i) = 1. \end{cases}$$

If K is a number field, one generalizes this construction in the following way:

$$H_N: \mathbf{P}^N(K) \rightarrow \mathbf{R}_{>0}$$

$$(x_0 : \dots : x_N) \mapsto \prod_{v \in \Omega_K} \sup_{0 \leq i \leq N} |x_i|_v$$

where Ω_K is the set of places of K and for any $x \in K_v$,

$$|x|_v = |N_{K_v/\mathbf{Q}_p}(x)|_p \quad \text{if } v | p.$$

Any morphism of varieties $\phi: V \rightarrow \mathbf{P}_K^N$ induces a height

$$H: V(K) \rightarrow \mathbf{R}_{>0}$$

$$x \mapsto H_N(\phi(x)).$$

If $U \subset V$ is an open subset, then we would like to describe and understand the asymptotic behavior of the counting function

$$N_{U,H}(B) = \#\{x \in U(K) \mid H(x) \leq B\}.$$


Let us first give a few examples:

Example 2.2. — If $V = \mathbf{P}^N(\mathbf{Q})$ and $\phi = \text{id}$, then an easy Möbius inversion formula gives that

$$N_{V,H}(B) \sim \frac{2^N}{\zeta_{\mathbf{Q}}(N+1)} B^{N+1}$$

as $B \rightarrow \infty$ (see Figure 1).

FIGURE 1. Projective space



This result was later generalized by Schanuel in [Sc] to the projective space over any number field (see Figure 2).


Example 2.3. — If $V = V_1 \times V_2$, and $H_i : V_i(K) \rightarrow \mathbf{R}_{>0}$ are heights as above, and $U_i \subset V_i$ is an open subset for each i , then the height $H : V \rightarrow \mathbf{R}_{>0}$ defined by $H(x_1, x_2) = H_1(x_1)H_2(x_2)$ corresponds to the Segre embedding of $V_1 \times V_2$. Assume that

$$N_{U_i, H_i}(B) = C_i B(\log B)^{t_i-1} + O(B(\log B)^{t_i-2}).$$

Then, by [FMT, Proposition 2]

$$N_{U_1 \times U_2, H}(B) \sim \frac{(t_1-1)!(t_2-1)!}{(t_1+t_2-1)!} C_1 C_2 B(\log B)^{t_1+t_2-1}$$

FIGURE 2. Projective line over $\mathbf{Q}(i)$




as $B \rightarrow \infty$. For $\mathbf{P}^1 \times \mathbf{P}^1$, we get

$$N_{\mathbf{P}^1 \times \mathbf{P}^1, H}(B) \sim CB^2 \log B$$

(see Figure 3).

FIGURE 3. Product of two projective lines



Example 2.4. — Let $V \rightarrow \mathbf{P}^2(\mathbf{Q})$ be the blow up of \mathbf{P}^2 at $P_1 = (1 : 0 : 0)$, $P_2 = (0 : 1 : 0)$, and $P_3 = (0 : 0 : 1)$. Then V may be seen as the hypersurface in $\mathbf{P}^1 \times \mathbf{P}^1 \times \mathbf{P}^1$ given by the equation $x_1x_2x_3 = y_1y_2y_3$. We put

$$H(P_1, P_2, P_3) = H_1(P_1)H_1(P_2)H_1(P_3)$$

which defines a height $H : V(\mathbf{Q}) \rightarrow \mathbf{R}_{>0}$.

On V , there are 6 exceptional lines E_{ij} : $x_i = 0, y_j = 0$ for $i \neq j$. Let $U = V - \cup_{i \neq j} E_{ij}$. We have


$$N_{E_{ij}, H}(B) \sim CB^2$$

and

$$N_{U, H}(B) \sim \frac{1}{6} \left(\prod_p \left(1 - \frac{1}{p} \right)^4 \left(1 + \frac{4}{p} + \frac{1}{p^2} \right) \right) B(\log B)^3$$

(see Figure 4).

FIGURE 4. The plane blown up



We see that $N_{U, H}(B) = o(N_{E_{ij}, H}(B))$. Thus, in this case, the dominant term of the asymptotic behaviour of $N_{V, H}(B)$ is given by the number of points on the six lines. Therefore it can not reflect the geometry of the whole of V . One of the basic ideas in the interpretation of the asymptotic behaviour of the number of points of bounded height is that one has to consider open subsets to be able to get a meaningful geometric interpretation.

In all examples known to the author for which it is possible to give a precise estimate of the number of points of bounded height, the asymptotic behaviour is of the form

$$N_{U, H}(B) \sim CB^a (\log B)^{b-1}$$

with $C \geq 0$, $a \geq 0$ and $b \in \frac{1}{2}\mathbf{Z}$, $b \geq 1$. Thus one wishes to give a geometric interpretation of a , b and C .

3. Manin's principle

We assume that V is a smooth, geometrically integral projective variety of dimension n over the number field K . We also assume that $\omega_V^{-1} = \Lambda^n T_V$ is very ample (in particular, V is a Fano variety). We look only at the height relative to this anticanonical divisor $\phi^*(\mathcal{O}_{\mathbf{P}^N}(1)) = \omega_V^{-1}$, and we assume that $V(K)$ is Zariski dense. The following question is a variant of the conjecture C' in [BM]:

Question 3.1. — *Does there exist a dense open subset $U \subset V$ and a constant $C > 0$ such that*

$$N_{U,H}(B) \sim CB(\log B)^{t-1}$$

as $B \rightarrow \infty$, where t is the rank of the Picard group of V . (Since V is Fano, $\text{Pic } V$ is a free \mathbf{Z} -module of finite rank.)

In fact, it is even possible to give a conjectural interpretation of C , but to describe this conjectural constant, we first need to express the height in terms of metrics.

Notation 3.2. — Let V be a geometrically integral smooth projective variety and H be the height corresponding to an embedding $\phi: V \rightarrow \mathbf{P}_K^N$. Let L be $\phi^*(\mathcal{O}_{\mathbf{P}^N}(1))$. We denote by s_0, \dots, s_N the pull-backs in $\Gamma(V, L)$ of the sections X_0, \dots, X_N of $\mathcal{O}_{\mathbf{P}^N}(1)$. We view L as a line bundle over V and define for any place v of K a v -adically continuous metric $\|\cdot\|_v: L(K_v) \rightarrow \mathbf{R}$ by the condition:

$$\forall x \in V(K_v), \quad \forall s \in \Gamma(V, L), \quad \|s(x)\|_v = \inf_{\substack{0 \leq i \leq N \\ s_i(x) \neq 0}} \left| \frac{s(x)}{s_i(x)} \right|_v.$$

Then the height H may be characterized by

$$\forall x \in V(K), \quad \forall s \in \Gamma(V, L), \quad s(x) \neq 0 \Rightarrow H(x) = \prod_{v \in \Omega_K} |s(x)|_v^{-1}.$$

From now on we assume that the above line bundle L is the anticanonical line bundle ω_V^{-1} . We now define a measure on the adelic space $V(\mathbf{A}_K)$ which coincides with the product $\prod_{v \in \Omega_K} V(K_v)$, since V is projective.

Definition 3.3. — For any place v of K , we normalize the Haar measure dx_v on K_v by the conditions

- $\int_{\mathcal{O}_v} dx_v = 1$ if v is finite,
- $dx_v([0, 1]) = 1$ if K_v is isomorphic to \mathbf{R} ,
- $dx_v = i dz d\bar{z} = 2 dx dy$ if K_v is isomorphic to \mathbf{C} .

The measure ω_v on $V(F_v)$ is defined locally by the formula

$$\omega_v = \left\| \frac{\partial}{\partial x_1} \wedge \cdots \wedge \frac{\partial}{\partial x_n} \right\|_v dx_{1,v} \cdots dx_{n,v}$$

if (x_1, \dots, x_n) is a local system of coordinates on $V(K_v)$ in the v -adic topology and where $\frac{\partial}{\partial x_1} \wedge \cdots \wedge \frac{\partial}{\partial x_n}$ is viewed as a section of ω_V^{-1} . The fact that these expressions glue together follows from the chosen normalization of the absolute value. Indeed the formula for a change of variables is given by

$$dy_{1,v} \cdots dy_{n,v} = \left| \det \left(\frac{\partial y_i}{\partial x_j} \right)_{\substack{1 \leq i \leq n \\ 1 \leq j \leq n}} \right|_v dx_{1,v} \cdots dx_{n,v}$$

(see [We, §2.2.1]).

Remark 3.4. — At any real place this construction amounts to the classical recipe for producing a measure on a differential variety from a continuous section of its canonical line bundle. At almost all finite places, using ideas of Tamagawa and Weil, one may prove the following proposition:

Proposition 3.5. — *For almost all finite \mathfrak{p} in Ω_K ,*

$$\omega_{\mathfrak{p}}(V(K_{\mathfrak{p}})) = \frac{\#V(\mathbf{F}_{\mathfrak{p}})}{(\#\mathbf{F}_{\mathfrak{p}})^{\dim V}}$$

where $\mathbf{F}_{\mathfrak{p}}$ is the residue field at \mathfrak{p} .

In particular, this implies that the product $\prod_{\mathfrak{p}} \omega_{\mathfrak{p}}(V(K_{\mathfrak{p}}))$ diverges. Therefore we have to introduce convergence factors. These factors are suggested by the Grothendieck-Lefschetz formula.

Definition 3.6. — We fix a finite set S of bad places containing all archimedean places and all places of bad reduction. Let \overline{K} be an algebraic closure of K and put $\overline{V} = V \times_K \overline{K}$. Then for $\mathfrak{p} \in \Omega_K - S$ one defines

$$L_{\mathfrak{p}}(s, \text{Pic}(\overline{V})) = \frac{1}{\det(1 - q^{-s} \text{Frob}_q | \text{Pic}(V_{\overline{\mathbf{F}}_{\mathfrak{p}}}) \otimes \mathbf{Q})}$$

where $q = \#\mathbf{F}_{\mathfrak{p}}$ and Frob_q is the q -power Frobenius automorphism of the field $\overline{\mathbf{F}}_{\mathfrak{p}}$, which induces a linear endomorphism of the \mathbf{Q} -vector space $\text{Pic}(V_{\overline{\mathbf{F}}_{\mathfrak{p}}}) \otimes \mathbf{Q}$.

The global L -function is given by the Euler product

$$L_S(s, \text{Pic}(\overline{V})) = \prod_{\mathfrak{p} \in \Omega_K - S} L_{\mathfrak{p}}(s, \text{Pic}(\overline{V}))$$

which converges for $\text{Re}(s) > 1$ and admits a meromorphic continuation to \mathbf{C} . We define the convergence factors by

$$\lambda_v = \begin{cases} L_v(1, \text{Pic}(\overline{V})), & \text{if } v \in \Omega_K - S, \\ 1, & \text{otherwise.} \end{cases}$$

The adelic measure on $V(\mathbf{A}_K)$ is then defined by the formula

$$\omega_H = \lim_{s \rightarrow 1} (s-1)^t L_S(s, \text{Pic}(\overline{V})) \frac{1}{\sqrt{d_K}^{\dim V}} \prod_{v \in \Omega_K} \lambda_v^{-1} \omega_v$$

where d_K is the absolute value of the discriminant of K .

Remarks 3.7. —

(i) The convergence of the product $\prod_{v \in \Omega_K} \lambda_v^{-1} \omega_v$ follows from the Lefschetz trace formula and Weil's conjecture about the absolute value of the eigenvalues of the Frobenius operator which was proven by Deligne [De].

(ii) By definition, the measure ω_H does not depend on S .

(iii) Note that $\sqrt{d_K}$ is the volume of \mathbf{A}_K/K for the measure $\prod_{v \in \Omega_K} dx_v$.

To define the conjectural constant it remains to multiply by two rational factors which are the object of the next definition.

Definition 3.8. — Let $C_{\text{eff}}^1(V)$ be the cone in $\text{Pic}(V) \otimes_{\mathbf{Z}} \mathbf{C}$ generated by the classes of effective divisors and $C_{\text{eff}}^1(V)^{\vee}$ the dual cone defined by

$$C_{\text{eff}}^1(V)^{\vee} = \{y \in \text{Pic}(V) \otimes_{\mathbf{Z}} \mathbf{R}^{\vee} \mid \forall x \in C_{\text{eff}}^1(V), \langle x, y \rangle \geq 0\}.$$

Then

$$\alpha(V) = \frac{1}{(t-1)!} \int_{C_{\text{eff}}^1(V)^{\vee}} e^{\langle \omega_V^{-1}, y \rangle} dy$$

where the measure on $\text{Pic}(V) \otimes_{\mathbf{Z}} \mathbf{R}^{\vee}$ is normalized so that the covolume of the dual lattice $\text{Pic}(V)^{\vee}$ is one. We also consider the integer

$$\beta(V) = \#H^1(K, \text{Pic}(\overline{V})).$$

Remarks 3.9. —

(i) The constant $\alpha(V)$ may also be defined as the volume of the domain

$$\{y \in C_{\text{eff}}^1(V)^\vee \mid \langle y, \omega_V^{-1} \rangle = 1\}$$

for a suitable measure on the affine hyperplane $\langle y, \omega_V^{-1} \rangle = 1$ (see [Pe1, §2.2.5]). Therefore, if there exists a finite family $(D_i)_{1 \leq i \leq r}$ of effective divisors on V such that

$$C_{\text{eff}}^1(V) = \sum_{i=1}^r \mathbf{R}_{\geq 0}[D_i]$$

then the constant $\alpha(V)$ is rational.

(ii) The constant $\beta(V)$ was introduced by Batyrev and Tschinkel in [BT1].

The conjectural constant is then defined as follows

Definition 3.10. — We define

$$\theta_H(V) = \alpha(V)\beta(V)\omega_H(\overline{V(K)}),$$

where $\overline{V(K)}$ denotes the closure of the rational points in the adelic space $V(\mathbf{A}_K)$.

We can now give a refined version of Question 3.1:

Empirical formula 3.11. — *With notation as in Question 3.1, there often exists a dense open subset $U_0 \subset V$ such that for any nonempty subset U of U_0 , one has*

$$(F) \quad N_{U,H}(B) \sim \theta_H(V)B(\log B)^{t-1}$$

4. Results

The formula (F) is true in the following cases:

- $V = G/P$, where G is a reductive algebraic group over K and P is a parabolic subgroup of G defined over K . It follows from the work of Langlands on Eisenstein series [Lan] (see Franke, Manin, Tschinkel [FMT] and [Pe1, §6]). We may take $U_0 = V$. In particular, it is true for any quadric.
- V is a smooth projective toric variety, that is an equivariant compactification of an algebraic torus (see [Pe1, §8–11] for particular cases, Batyrev and Tschinkel [BT1], [BT2], and [BT4], Salberger [Sal], and de la Bretèche [Bre1]). One may take the open orbit as U_0 . This case includes the plane blown up in 1, 2, or 3 points, and Hirzebruch surfaces.
- V is an equivariant compactification of an affine space for the action of the corresponding vector space (see Chambert-Loir, Tschinkel [CLT1], [CLT2], and [CLT3]).

— $V = \mathbf{P}_{\mathbf{Q}}^2$ blown up at $(1 : 0 : 0)$, $(0 : 1 : 0)$, $(0 : 0 : 1)$, $(1 : 1 : 1)$ (see Salberger for an upper bound, and de la Bretèche [Bre2]).

The formula (F) is compatible with:

- the circle method. In particular, it is true if $V \subset \mathbf{P}^n(\mathbf{Q})$ is a smooth hypersurface of degree d , if $n > 2^d(d-1)$ (see Birch [Bir]);
- products of varieties (see [FMT], [Pe1, §4]);
- numerical tests for some diagonal cubic surfaces [PT1], [PT2];
- lower bounds for some cubic surfaces (see Slater and Swinnerton-Dyer [SSD]). The problem of finding an optimal upper bound for cubic surfaces is still open.

All these examples support the empirical formula. However, there are also counterexamples, which will be discussed in the next section.

5. The counterexample of Batyrev and Tschinkel

Take $V \subset \mathbf{P}^3 \times \mathbf{P}^3$ defined by

$$x_0y_0^3 + x_1y_1^3 + x_2y_2^3 + x_3y_3^3 = 0.$$

We have

$$\text{Pic}(V) \simeq \text{Pic}(\mathbf{P}^3 \times \mathbf{P}^3) = \mathbf{Z} \times \mathbf{Z}$$

and $\omega_V^{-1} = \mathcal{O}_V(3, 1)$. In particular, V is a Fano variety. We may use the height

$$H: V(\mathbf{Q}) \rightarrow \mathbf{R}_{>0}$$

$$((x_0 : \dots : x_3), (y_0 : \dots : y_3)) \mapsto H_3(x)^3 H_3(y).$$

If (F) is true for V then there is an open subset U and a constant C such that

$$N_{U,H}(B) \sim CB \log B$$

as $B \rightarrow \infty$. There is a projection onto the first coordinate $\pi_1 : V \rightarrow \mathbf{P}^3$. If $(x_0 : \dots : x_3) \in \mathbf{P}^3$ is such that $\prod_{i=0}^3 x_i \neq 0$, then $\pi_1^{-1}(x)$ is a smooth cubic surface; if $x_1/x_0, x_2/x_0, x_3/x_0$ are cubes, then $\text{rk Pic}(\pi_1^{-1}(x)) = 4$. If (F) is true for the fiber, then

$$N_{\pi_1^{-1}(x),H}(B) \sim C_x B (\log B)^3$$

as $B \rightarrow \infty$, but these fibers are Zariski dense, so the answer to Question 3.1 can not be positive for both V and the fibers. In fact, Batyrev and Tschinkel prove the following more precise result:

Theorem 5.1 (Batyrev and Tschinkel [BT3]). — *If K contains a nontrivial cube root of unity, then for all nonempty $U \subset V$, (F) does not hold for U .*

6. Methods of counting

We now return to the methods used to prove the results given in Section 4.

Harmonic analysis: Assume that there exists a dense open subset U of V which is of the form G/H , where G is a reductive algebraic group, look at the height zeta function

$$\zeta_{U,H}(s) = \sum_{x \in U(K)} H(x)^{-s}$$

which converges when $\operatorname{Re}(s) \gg 0$.

The asymptotic behavior of $N_{U,H}(B)$ is given by the meromorphic properties of $\zeta_{U,H}(s)$. If $U = G$, one may use a Poisson formula. If $V = G/P$, $\zeta_{U,H}(s)$ is an Eisenstein series and we may apply the work of Langlands. In both cases the problem may be handled using harmonic analysis.

These methods do not apply when the variety is not an equivariant compactification of a homogeneous space. All other cases appearing in the list of section 4 have one preliminary step in common: they all use a lift to universal torsors.

Universal torsors: implicit in the case of a hypersurface in $\mathbf{P}^n(\mathbf{Q})$, it was made explicit by Salberger in [Sal] in his alternative treatment of split toric varieties over \mathbf{Q} ; it was then used by Salberger and de la Bretèche in the case of the plane blown up in 4 points. The end of this survey is devoted to the description of this preliminary step in those cases.

7. A basic example

In the case of hypersurfaces of large dimension and small degree, the principle of Manin follows from the following deep theorem which is based upon the Hardy-Littlewood circle method.

Theorem 7.1 (Birch [Bir]). — *Let $f \in \mathbf{Z}[x_0, \dots, x_N]$ be homogeneous of degree d , and let $W \subset \mathbf{A}^{N+1} - \{0\}$ be the cone defined by $f = 0$. Assume that:*

- (i) W is smooth,
- (ii) $W(\mathbf{R}) \neq \emptyset$, and for all primes p , $W(\mathbf{Q}_p) \neq \emptyset$,
- (iii) $N > 2^d(d-1)$.

Let

$$M_W(B) = \#\left\{x \in \mathbf{Z}^{N+1} - \{0\} \mid f(x) = 0 \text{ and } \sup_{0 \leq i \leq N} |x_i| \leq B\right\}.$$

Then there exist explicit $C > 0$ and $\delta > 0$ such that

$$M_W(B) = CB^{N+1-d} + O(B^{N+1-d-\delta}).$$

Let $\pi : \mathbf{A}^{N+1} - \{0\} \rightarrow \mathbf{P}^N$, and let $V = \pi(W)$ be the corresponding projective hypersurface. Then $\omega_V^{-1} = \mathcal{O}_V(N+1-d)$, so we may take the height

$$H(x) = H_N(x)^{N+1-d}$$

where H_N was defined in Section 2. Then

$$N_{V,H}(B) = \frac{1}{2} \# \left\{ x \in \mathbf{Z}^{N+1} - \{0\} \left| \begin{array}{l} f(x) = 0, \\ \sup |x_i|^{N+1-d} \leq B, \\ \gcd(x_i) = 1. \end{array} \right. \right\}.$$

Using Möbius inversion, we get

$$N_{V,H}(B) = \frac{1}{2} \sum_k \mu(k) \# \left\{ x \in (k\mathbf{Z})^{N+1} - \{0\} \left| \begin{array}{l} f(x) = 0, \\ \sup |x_i|^{N+1-d} \leq B \end{array} \right. \right\},$$

where $\mu : \mathbf{Z}_{>0} \rightarrow \{-1, 0, 1\}$ is the Möbius function. Then

$$\begin{aligned} N_{V,H}(B) &= \frac{1}{2} \sum_k \mu(k) M_W \left(\frac{B^{1/(N+1-d)}}{k} \right) \sim \frac{C}{2} \sum_k \frac{\mu(k)}{k^{N+1-d}} B \\ &= \frac{C}{2\zeta(N+1-d)} B. \quad \square \end{aligned}$$

The motivation behind the introduction of universal torsors was to generalize this simple descent argument to other varieties.

8. Universal torsors

Let V be a smooth, geometrically integral projective variety over K , where $\text{char } K = 0$. Assume (for simplicity) that V is Fano, which means that ω_V^{-1} is ample. Thus, if \overline{K} is an algebraic closure of K , and $\overline{V} = V \times_K \overline{K}$, then $\text{Pic}(\overline{V})$ is a free abelian group of finite rank.

Assume $K = \overline{K}$ first. Let L_1, \dots, L_t be line bundles on V such that $[L_1], \dots, [L_t]$ form a basis of $\text{Pic}(V) = \text{Pic}(\overline{V})$. Let $L_i^\times = L_i - \text{zero section}$. Consider

$$\pi : L_1^\times \times_V L_2^\times \times_V \cdots \times_V L_t^\times \rightarrow V.$$

On the left we have an action of \mathbf{G}_m^t , this is ‘the’ *universal torsor* of V .

Proposition 8.1. — *If $K = \overline{K}$, the universal torsor constructed above does not depend, up to isomorphism, on the chosen basis of the Picard group.*

Proof. — Let L'_1, \dots, L'_t be line bundles on V whose classes $[L'_1], \dots, [L'_t]$ form another basis of the Picard group of V . Let $M = (m_{i,j})$ in $\mathrm{GL}_n(\mathbf{Z})$ be the matrix such that

$$[L'_i] = \sum_{j=1}^t m_{j,i} [L_j].$$

In other words, for each i in $\{1, \dots, t\}$ we may fix an isomorphism

$$\psi_i : \bigotimes_{j=1}^t L_j^{\otimes m_{j,i}} \xrightarrow{\sim} L'_i.$$

But if E_1, \dots, E_m are one-dimensional vector spaces and k_1, \dots, k_m integers there is a canonical map

$$\begin{aligned} \times_{i=1}^m (E_i - \{0\}) &\rightarrow \bigotimes_{i=1}^m E_i^{\otimes k_i} \\ (y_1, \dots, y_m) &\mapsto \bigotimes_{i=1}^m y_i^{\otimes k_i} \end{aligned}$$

where for any vector space E of dimension one, and any non-zero y in E , $y^{\otimes -1}$ is the unique element of the dual E^\vee of E such that $y^{\otimes -1}(y) = 1$. In that way, composing with ψ_i , we get maps

$$\rho_i : \times_{j=1}^t L_j^\times \rightarrow L'_i{}^\times.$$

This map is equivariant for the action of \mathbf{G}_m^t in the following sense:

$$\forall (z_1, \dots, z_t) \in \mathbf{G}_m^t(K), \forall y \in \times_{j=1}^t L_j^\times(K), \rho_i((z_1, \dots, z_t) \cdot y) = \prod_{j=1}^t z_j^{m_{j,i}} \cdot \rho_i(y).$$

Note that if ρ'_i is another map from $\times_{j=1}^t L_j^\times$ to $L'_i{}^\times$ with the same equivariance property, then there is a section $s \in \Gamma(V, \mathbf{G}_m)$ such that

$$\forall y \in \times_{j=1}^t L_j^\times(K), \quad \rho'_i(y) = s(\pi(y)) \cdot \rho_i(y).$$

But, since V is projective, $\Gamma(V, \mathbf{G}_m) = K^*$ and ρ_i is unique up to multiplication by a constant. The maps ρ_i yield a map

$$\rho : \times_{i=1}^t L_i^\times \rightarrow \times_{i=1}^t L'_i{}^\times.$$

The matrix M defines a morphism of algebraic groups

$$\begin{aligned} \widetilde{M} : \mathbf{G}_m^t &\rightarrow \mathbf{G}_m^t \\ (z_1, \dots, z_t) &\mapsto (\prod_{j=1}^t z_j^{m_{j,i}})_{1 \leq i \leq t} \end{aligned}$$

and the map ρ is equivariant with respect to \widetilde{M} :

$$\forall z \in \mathbf{G}_m^t(K), \quad \forall y \in \prod_{i=1}^t L_i^\times(K), \quad \rho(z \cdot y) = \widetilde{M}(z) \cdot \rho(y).$$

Moreover, if ρ' is another map with the same equivariance property, then there is $z \in \mathbf{G}_m^t(K)$ such that $\rho' = z \cdot \rho$. Similarly we may define a map

$$\tau : \prod_{i=1}^t L_i'^\times \rightarrow \prod_{i=1}^t L_i^\times$$

which is equivariant with respect to \widetilde{M}^{-1} . Thus the composite map

$$\tau \circ \rho : \prod_{i=1}^t L_i'^\times \rightarrow \prod_{i=1}^t L_i^\times$$

is equivariant with respect to the identity map and therefore coincides with the action of an element of $\mathbf{G}_m^t(K)$. Thus $\tau \circ \rho$ and $\rho \circ \tau$ are isomorphisms. \square

For arbitrary fields, a universal torsor may be described as a K -structure on the above torsor. Let us define this notion more precisely:

Recall that there is a contravariant equivalence of categories between the category of algebraic tori, that is, algebraic groups T such that \overline{T} is isomorphic to $\mathbf{G}_{m, \overline{K}}^{\dim T}$, and the category of $\text{Gal}(\overline{K}/K)$ -lattices, that is, $\text{Gal}(\overline{K}/K)$ -modules which are free abelian groups of finite rank. The functors giving the equivalence are

$$T \mapsto X^*(T) = \text{Hom}_{\text{alg. gp.}}(\overline{T}, \mathbf{G}_m) \quad \text{and} \quad M \mapsto \text{Spec}(\overline{K}[M])^{\text{Gal}(\overline{K}/K)}.$$

Definition 8.2. — Let the Néron-Severi torus, T_{NS} , be the torus corresponding to the $\text{Gal}(\overline{K}/K)$ -lattice $\text{Pic } \overline{V}$.

If G is an algebraic group over K , then a G -torsor is a faithfully flat map $\pi : \mathcal{T} \rightarrow V$ with an action of G on \mathcal{T} such that locally for the faithfully flat topology, $\mathcal{T} \times_V U \simeq G \times U$, where the isomorphism is compatible with the action of G . (In another language, these are principal homogeneous spaces.)

A T_{NS} -torsor $\mathcal{T} \rightarrow V$ is said to be *universal* if $\overline{\mathcal{T}} \rightarrow \overline{V}$ is isomorphic as a torsor to $L_1^* \times_V \cdots \times_V L_t^* \rightarrow V$.

Why are these universal torsors interesting? The following facts are due to Colliot-Thélène and Sansuc, who introduced this notion.

- Proposition 8.3 (Colliot-Thélène, Sansuc).** — *With notations as above,*
- *For all $x \in V(K)$, there exists a unique (up to isomorphism) universal torsor $\pi : \mathcal{T} \rightarrow V$ such that $x \in \pi(\mathcal{T}(K))$.*
 - *If K is a number field, there exist up to isomorphism only finitely many universal torsors $\pi : \mathcal{T} \rightarrow V$ such that $\mathcal{T}(K) \neq \emptyset$.*

This proposition gives us a nice decomposition of the set of rational points

$$V(K) = \bigsqcup_{1 \leq i \leq m} \pi_i(\mathcal{T}_i(K)).$$

Heuristic 8.4. — *From the arithmetical point of view, universal torsors should be much simpler than the variety V .*

This heuristic can be justified by the following statement:

Proposition 8.5 (Colliot-Thélène, Sansuc). — *If \mathcal{T}^c is a smooth projective compactification of a universal torsor $\mathcal{T} \rightarrow V$, then $\mathcal{T}^c(A_K)^{\text{Br}} = \mathcal{T}^c(A_K)$. In other words, there is no Brauer-Manin obstruction to the Hasse principle or weak approximation.*

Example 8.6. — Let $V \subset \mathbf{P}^N$ be a hypersurface over \mathbf{Q} with $\dim V \geq 3$, $\deg V = d$, and $N + 1 - d > 0$. Then the cone $W \subset \mathbf{A}^{N+1} - \{0\}$ above V is, up to isomorphism, the unique universal torsor over V .

9. Toric varieties

The following construction is due to Cox. Let T be an algebraic torus and V a smooth projective equivariant compactification of T . This means that there is an action of T on V , an open subset $U \subset V$, and an isomorphism from U to T compatible with the actions of T .

Denote by $\Sigma(1)$ the set of orbits of codimension 1 in \overline{V} . Then there is an exact sequence of $\text{Gal}(\overline{K}/K)$ -modules

$$0 \rightarrow X^*(T) \xrightarrow{\text{div}} \mathbf{Z}^{\Sigma(1)} \xrightarrow{\rho} \text{Pic}(\overline{V}) \rightarrow 0$$

$$e_\sigma \mapsto [D_\sigma]$$

where D_σ is the closure of the orbit σ in \overline{V} , which is an irreducible divisor of \overline{V} . Moreover, we have

$$\omega_V^{-1} = \sum_{\sigma \in \Sigma(1)} [D_\sigma].$$

By duality, we get an exact sequence of tori

$$1 \rightarrow T_{\text{NS}} \rightarrow T_{\Sigma(1)} \xrightarrow{\pi} T \rightarrow 1.$$

But $T \subset V$ and we want to extend the map π to get a torsor over V . We do this in the following way: We consider the affine space

$$\mathbf{A}_{\Sigma(1)} = \text{Spec}\left(\overline{K}[X_\sigma]_{\sigma \in \Sigma(1)}^{\text{Gal}(\overline{K}/K)}\right)$$

and a closed subset $F \subset \mathbf{A}_{\Sigma(1)}$, defined over \overline{K} as a union of affine subspaces,

$$\overline{F} = \bigcup_{\substack{I \subset \Sigma(1) \\ \bigcap_{\sigma \in I} D_\sigma = \emptyset}} \left(\bigcap_{\sigma \in I} (X_\sigma = 0) \right).$$

Note that \overline{F} is stable under the action of the Galois group, so it is defined over K . We take $\mathcal{F} = \mathbf{A}_{\Sigma(1)} - F$.

Claim 9.1. — *For all $x \in T(K)$, the map*

$$\begin{aligned} T_{\Sigma(1)} &\rightarrow T \\ t &\mapsto \pi(t) \cdot x \end{aligned}$$

extends to a map $\mathcal{F} \rightarrow V$ sending $1 \in T_{\Sigma(1)}$ to x .

Theorem 9.2 (Colliot-Thélène, Sansuc, Salberger, Madore)

The above construction gives a bijection between $T(K)/T_{\Sigma(1)}(K)$ and isomorphism classes of universal torsors over V .

We return now to the problem of counting points. We assume that $K = \mathbf{Q}$, that the action of $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ on $X^*(T)$ and $\Sigma(1)$ are trivial, and that ω_V^{-1} is generated by global sections.

Then we consider

$$\mathcal{M} = \{m \in \mathbf{Z}^{\Sigma(1)} \mid \forall \sigma \in \Sigma(1), m_\sigma \geq 0 \text{ and } \rho(m) = \omega_V^{-1} \in \text{Pic } V\}.$$

For all $m \in \mathcal{M}$, let $X^m \in \mathbf{Q}[X_\sigma]_{\sigma \in \Sigma(1)}$ be the corresponding monomial. We lift the height to the universal torsor by

$$\widetilde{H}((y_\sigma)_{\sigma \in \Sigma(1)}) = \sup_{m \in \mathcal{M}} |X^m((y_\sigma)_{\sigma \in \Sigma(1)})|.$$

Theorem 9.3 (Salberger [Sal]). — *There exists a height H relative to ω_V^{-1} such that $N_{U,H}(B) = \widetilde{N}(B)/2^{\dim T_{\text{NS}}}$, where $\widetilde{N}(B)$ is the number of $(y_\sigma)_{\sigma \in \Sigma(1)}$ in $\mathbf{Z}^{\Sigma(1)}$ such that*

$$\begin{cases} \widetilde{H}(y) \leq B, \\ \forall I \subset \Sigma(1), \bigcap_{\sigma \in I} D_\sigma = \emptyset \Rightarrow \gcd(y_\sigma)_{\sigma \in I} = 1. \end{cases}$$

To prove Manin's conjecture in that case one may then proceed as follows: By use of a Möbius inversion formula, reduce to give an estimate

$$\#\left\{ (y_\sigma)_{\sigma \in \Sigma(1)} \in (\mathbf{Z} - \{0\})^{\Sigma(1)} \mid \widetilde{H}(y) \leq B \right\}.$$

and prove that when B goes to $+\infty$ this is asymptotic to

$$\text{vol}\left(\left\{ (y_\sigma)_{\sigma \in \Sigma(1)} \in \mathbf{R}^{\Sigma(1)} \mid \widetilde{H}(y) \leq B \right\}\right) \sim CB(\log B)^{\text{rk Pic } V - 1},$$

which proves the conjecture in this case.

10. The plane blown up in 4 points

The construction is due to Salberger and Skorobogatov. We consider in this section the blowup $\pi : V \rightarrow \mathbf{P}^2$ of $P_1 = (1 : 0 : 0)$, $P_2 = (0 : 1 : 0)$, $P_3 = (0 : 0 : 1)$, and $P_4 = (1 : 1 : 1)$. The exceptional divisors on V are $E_{i,5} = \pi^{-1}(P_i)$ and $E_{i,j}$, the strict pullback of the line through P_k and P_l if $\{i,j,k,l\} = \{1,2,3,4\}$. Then $E_{i,j} \cap E_{k,l} = \emptyset$ if and only if $\{i,j\} \cap \{k,l\} \neq \emptyset$. Then we consider the Grassmannian variety $\text{Gr}(2,5)$ of the planes in \mathbf{Q}^5 ; we may embed it into $\mathbf{P}(\Lambda^2 \mathbf{Q}^5)$. The cone above it, $W \subset \Lambda^2 \mathbf{Q}^5$ is given by the Plücker relations:

$$\begin{cases} X_{1,2}X_{3,4} - X_{1,3}X_{2,4} + X_{1,4}X_{2,3} = 0, \\ X_{1,2}X_{3,5} - X_{1,3}X_{2,5} + X_{1,5}X_{2,3} = 0, \\ X_{1,2}X_{4,5} - X_{1,4}X_{2,5} + X_{1,5}X_{2,4} = 0, \\ X_{1,3}X_{4,5} - X_{1,4}X_{3,5} + X_{1,5}X_{3,4} = 0, \\ X_{2,3}X_{4,5} - X_{2,4}X_{3,5} + X_{2,5}X_{3,4} = 0. \end{cases}$$

Indeed, the vector space $\Lambda^2 \mathbf{Q}^5$ has dimension 10, and we take $X_{i,j}$ as coordinates corresponding to the basis elements $e_i \wedge e_j$ for $i \neq j$. We consider the closed subset $F \subset W$ given by

$$F = \bigcup_{\{i,j\} \cap \{k,\ell\} \neq \emptyset} ((X_{i,j} = 0) \cap (X_{k,\ell} = 0)),$$

and define $\mathcal{T} = W - F$. There is an action $\mathbf{G}_m^5 \subset GL_5(\mathbf{Q})$ on \mathcal{T} , and $\mathcal{T}/\mathbf{G}_m^5$ is isomorphic to V and $\mathcal{T} \rightarrow V$ is up to isomorphism the only universal torsor.

We put

$$\mathcal{M} = \left\{ (m_{i,j})_{i < j} \in \mathbf{Z}^{10} \mid \sum m_{i,j} E_{i,j} = \omega_V^{-1} \right\}$$

and for $m \in \mathcal{M}$, $X^m \in \mathbf{Q}[X_{i,j}, i < j]$. Then we may lift the height by using

$$\widetilde{H}((y_{i,j})) = \sup_{m \in \mathcal{M}} |X^m(y)|.$$

Proposition 10.1 (Salberger). — *There is a height H relative to ω_V^{-1} such that $N_{U,H}(B) = \widetilde{N}(B)/2^{\dim T_{\text{NS}}}$ where $\widetilde{N}(B)$ is the number of $(y_{i,j})$ in $W(\mathbf{Z})$ such that*

$$\begin{cases} \widetilde{H}(y_{i,j}) \leq B, \\ \{i,j\} \cap \{k,l\} \neq \emptyset \Rightarrow \gcd(y_{i,j}, y_{k,l}) = 1. \end{cases}$$

This description, which was made by Salberger in order to get an upper bound for the number of points of bounded height, was also the first step in the proof of the empirical formula (F), given by de la Bretèche in [Bre2].

11. Generalization

The papers [Pe2] and [Pe3], showed how to generalize the lifting described in Sections 7, 9, and 10. The first fact enabling this generalization is the existence of a gauge form on each universal torsor:

Proposition 11.1. — *If V is a Fano variety over K , and \mathcal{T} a universal torsor over V , then*

- *the canonical bundle $\omega_{\mathcal{T}}$ is trivial,*
- $\Gamma(\mathcal{T}, \mathbf{G}_m) = K^*$.

As usual, a gauge form yields a measure:

Definition 11.2. — *Up to a constant, there exists a unique non-vanishing section of the canonical line bundle. Let $\check{\omega}_{\mathcal{T}}$ be such a section. This section defines for any place v of K a measure $\omega_{\mathcal{T},v}$ on $\mathcal{T}(K_v)$ which is locally given by*

$$\omega_{\mathcal{T},v} = \left| \left\langle \frac{\partial}{\partial x_1} \wedge \cdots \wedge \frac{\partial}{\partial x_N}, \check{\omega}_{\mathcal{T}} \right\rangle \right|_v dx_{1,v} \cdots dx_{N,v}$$

where (x_1, \dots, x_N) is an analytic local system of coordinates on $\mathcal{T}(K_v)$ for v -adic topology. We then define a canonical measure on $\prod_{v \in \Omega_K} \mathcal{T}(K_v)$ by

$$\omega_{\mathcal{T}} = \frac{1}{\sqrt{d_K}^{\dim \mathcal{T}}} \prod_{v \in \Omega_K} \omega_{\mathcal{T},v}$$

Remarks 11.3. —

(i) If V is a hypersurface in \mathbf{P}^n , the measure $\omega_{\mathcal{T},v}$ coincides with the classical Leray measure on $\mathcal{T}(K_v)$.

(ii) The measure $\omega_{\mathcal{T}}$ does not depend on the choice of the section $\check{\omega}_{\mathcal{T}}$.

(iii) The volume $\omega_{\mathcal{T}}(\prod_{v \in \Omega_K} \mathcal{T}(K_v))$ is infinite, but if $S \subset \Omega_K$ is a finite set of places containing the archimedean ones and \mathcal{T} a model of \mathcal{T} over the ring of S -integers \mathcal{O}_S , then the product $\prod_{v \in \Omega_K - S} \omega_{\mathcal{T},v}(\mathcal{T}(\mathcal{O}_v))$ converges.

Let $\mathcal{T}_1, \dots, \mathcal{T}_r$ be torsors representing all isomorphism classes of universal torsors over V having a rational point. It is then possible to construct families of integrable functions $\Psi_{i,j,B} : \prod_{v \in \Omega_K} \mathcal{T}(K_v) \rightarrow \mathbf{R}_{>0}$ such that upper bounds for the difference

$$\left| \sum_{y \in \mathcal{T}_i(K)} \Psi_{i,j,B}(y) - \int \prod_{v \in \Omega_K} \mathcal{T}(K_v) \Psi_{i,j,B}(y) \omega_{\mathcal{T}} \right|$$

yield an upper bound for the difference

$$|N_{U,H}(B) - \theta_H(V)B(\log B)^{t-1}|.$$

The liftings described in previous sections may be seen as particular cases of this descent argument.

References

- [BM] V. V. Batyrev and Y. I. Manin, *Sur le nombre des points rationnels de hauteur bornée des variétés algébriques*, Math. Ann. **286** (1990), 27–43.
- [BT1] V. V. Batyrev and Y. Tschinkel, *Rational points of bounded height on compactifications of anisotropic tori*, Internat. Math. Res. Notices **12** (1995), 591–635.
- [BT2] ———, *Height zeta functions of toric varieties*, J. Math. Sci. **82** (1996), n° 1, 3220–3239.
- [BT3] ———, *Rational points on some Fano cubic bundles*, C. R. Acad. Sci. Paris Sér. I Math. **323** (1996), n° 1, 41–46.

- [BT4] ———, *Manin's conjecture for toric varieties*, J. Algebraic Geom. **7** (1998), n° 1, 15–53.
- [Bir] B. J. Birch, *Forms in many variables*, Proc. Roy. Soc. London **265A** (1962), 245–263.
- [Bre1] R. de la Bretèche, *Compter des points d'une variété torique*, J. of Number theory **87** (2001), 315–331.
- [Bre2] ———, *Nombre de points de hauteur bornée sur les surfaces de Del Pezzo de degré 5*, Duke Math. J. **113** (2002), n° 3, 421–464.
- [CLT1] A. Chambert-Loir and Y. Tschinkel, *Points of bounded height on equivariant compactifications of vector groups, I*, Compositio Math. **124** (2000), n° 1, 65–93.
- [CLT2] ———, *Points of bounded height on equivariant compactifications of vector groups, II*, J. of Number Theory **85** (2000), n° 2, 172–188.
- [CLT3] ———, *On the distribution of points of bounded height on equivariant compactifications of vector groups*, Invent. Math. **148** (2002), n° 2, 421–452.
- [CTS1] J.-L. Colliot-Thélène et J.-J. Sansuc, *Torseurs sous des groupes de type multiplicatif; applications à l'étude des points rationnels de certaines variétés algébriques*, C. R. Acad. Sci. Paris Sér. A **282** (1976), 1113–1116.
- [CTS2] ———, *La descente sur une variété rationnelle définie sur un corps de nombres*, C. R. Acad. Sci. Paris Sér. A **284** (1977), 1215–1218.
- [CTS3] ———, *La descente sur les variétés rationnelles*, Journées de géométrie algébrique d'Angers (1979) (A. Beauville, ed.), Sijthoff & Noordhoff, Alphen aan den Rijn, 1980, pp. 223–237.
- [CTS4] ———, *La descente sur les variétés rationnelles, II*, Duke Math. J. **54** (1987), n° 2, 375–492.
- [De] P. Deligne, *La conjecture de Weil I.*, Publ. Math. I.H.E.S. **43** (1974), 273–307.
- [FMT] J. Franke, Y. I. Manin, and Y. Tschinkel, *Rational points of bounded height on Fano varieties*, Invent. Math. **95** (1989), 421–435.
- [Lan] R. P. Langlands, *On the functional equations satisfied by Eisenstein series*, Lecture Notes in Math., vol. 544, Springer-Verlag, Berlin, Heidelberg and New York, 1976.
- [Pe1] E. Peyre, *Hauteurs et mesures de Tamagawa sur les variétés de Fano*, Duke Math. J. **79** (1995), n° 1, 101–218.
- [Pe2] ———, *Terme principal de la fonction zêta des hauteurs et toseurs universels*, Nombre et répartition de points de hauteur bornée, Astérisque, vol. 251, SMF, Paris, 1998, pp. 259–298.
- [Pe3] ———, *Torseurs universels et méthode du cercle*, Rational points on algebraic varieties, Progress in Math., vol. 199, Birkhäuser, Basel, 2001, pp. 221–274.

- [PT1] E. Peyre and Y. Tschinkel, *Tamagawa numbers of diagonal cubic surfaces, numerical evidence*, Math. Comp. **70** (2000), n° 233, 367–387.
- [PT2] ———, *Tamagawa numbers of diagonal cubic surfaces of higher rank*, Rational points on algebraic varieties, Progress in Math., vol. 199, Birkhäuser, Basel, 2001, pp. 275–305.
- [Sal] P. Salberger, *Tamagawa measures on universal torsors and points of bounded height on Fano varieties*, Nombre et répartition de points de hauteur bornée, Astérisque, vol. 251, SMF, Paris, 1998, pp. 91–258.
- [Sc] S. H. Schanuel, *Heights in number fields*, Bull. Soc. Math. France **107** (1979), 433–449.
- [SSD] J. B. Slater and H. P. F. Swinnerton-Dyer, *Counting points on cubic surfaces, I*, Nombre et répartition de points de hauteur bornée, Astérisque, vol. 251, SMF, Paris, 1998, pp. 1–12.
- [ST1] M. Strauch and Y. Tschinkel, *Height zeta functions of twisted products*, Math. Res. Lett. **4** (1997), 273–282.
- [ST2] ———, *Height zeta functions of toric bundles over flag varieties*, Selecta Math. (N.S.) **5** (1999), n° 3, 325–396.
- [We] A. Weil, *Adèles and algebraic groups*, Progress in Mathematics, vol. 23, Birkhäuser, Boston, Basel, Stuttgart, 1982.

October 5, 2022

EMMANUEL PEYRE, Institut Fourier, UMR 5582 du CNRS,
Université de Grenoble I, BP 74, 38402 Saint-
Martin d'Hères, France • *E-mail*: Emmanuel.Peyre@ujf-grenoble.fr