

RESEARCH STATEMENT: BOULANGER JULIEN

1. INTRODUCTION

My research focuses on several geometric and dynamical questions related to translation surfaces. Translation surfaces lie at the crossroad between (Riemannian and algebraic) geometry, low-dimensional dynamics, ergodic theory, geometric group theory, Teichmüller theory, and number theory. In the first part of my thesis, I was specifically interested in the algebraic intersection of closed curves on these surfaces, and more precisely the question of knowing *how many times two closed curves of a given length can intersect*. The second part of my thesis is devoted to the study of *Veech groups* of translation surfaces, which somehow encode the symmetries of each given surface and which are Fuchsian groups. A fundamental question is to know *which Fuchsian groups can be realized as Veech groups of translation surfaces*.

Informally, a translation surface can be seen as a collection of polygons in the plane with parallel sides identified by translations. Such a surface inherits a flat metric except on a finite number of conical singularities whose angles are integer multiples of 2π , and which correspond to certain vertices of the polygons. The set of all translation surfaces¹ (of genus $g \geq 1$) has a natural geometric structure and can be interpreted as the Hodge bundle $\Omega\mathcal{M}_g$ of the holomorphic 1-forms above the moduli space \mathcal{M}_g of complex structures on a (compact orientable) genus g surface. It is naturally stratified by the number and the order of the singularities; namely $\Omega\mathcal{M}_g = \sqcup \mathcal{H}_g(k_1, \dots, k_n)$, where the k_i are integers representing the order of the singularities satisfying $\sum_{i=1}^n k_i = 2g - 2$.




FIGURE 1. On the left, the double heptagon translation surface. It has genus three and a single singularity of angle 10π . On the right, a fundamental domain in the Poincaré half-plane \mathbb{H}^2 for the orbit of the double heptagon under the action of $SL_2(\mathbb{R})$. Each point corresponds to an element of the orbit and the double heptagon lies at the two identified corners. The Veech group of the double heptagon is conjugated to the Hecke group of level seven.

This structure proves to be particularly rich, and in particular it carries a natural action of $GL_2^+(\mathbb{R})$ which comes from the action on the polygons. As shown by Veech, the stabilizer of a given $X \in \Omega\mathcal{M}_g$ is a Fuchsian group, which is now referred to as the Veech group of X . Every element of the Veech group induces an² *affine diffeomorphism* of the surface. The work of several authors, including Veech, has shown that this group encodes many geometric properties of the surface. The translation surfaces whose Veech group is a lattice (called Veech surfaces) form a

¹modulo the action of the mapping class group.

²In some cases, it may also induce several different affine diffeomorphisms.

class of particularly interesting surfaces, both from the geometrical and dynamical point of view. The orbit of these surfaces under the action of $SL_2(\mathbb{R}) \subset GL_2^+(\mathbb{R})$ projects in \mathcal{M}_g to a complex geodesic, which is called a *Teichmüller curve*. The classification of Teichmüller curves, and more generally of orbit closures under the $SL_2(\mathbb{R})$ -action, is a central problem in Teichmüller dynamics where recent progress lead to the developpement of many new and revolutionnary techniques, including the work of award winning mathematicians like M. Mirzakhani, C. McMullen and A. Eskin, to name a few.

Organization. In the sequel, I will describe the following.

- In Section 2, I adress the question of the classification of Teichmüller curves by explaining a result joint with Sam Freedman [BF22] which shows that there is no geometrically primitive Veech surface in the locus $Prym(2, 2) \subset \mathcal{H}(2, 2)$ of surfaces having a *Prym involution*.
- In Section 3, I focus on translation surfaces with infinitely generated Veech groups, and more specifically on the study of *connection points* that allow to build such surfaces. In [Bou22], I prove that the central points of the double heptagon are not connection points, which gives a negative answer to a question of P. Hubert and T. Schmidt. This question is related to the distribution of *saddle connections* as well as the notion of *Hecke continued fractions*, and exhibits a deep link with a problem from number theory.
- In Section 4, I elaborate upon the notion of (*algebraic*) *interaction strength* which quantifies the maximum number of intersection of two curves of given length on a Riemannian surface with conical singularities. I will specifically address the case of flat surfaces and the behavior of the interaction strength by deformation of a given surface, which I studied in [BLM22], [Bou23a] and [Bou23b] as well as in an ongoing work with I. Pasquinelli.

Finally, in Section 5, I describe a few other questions I am interested in. I am also up to working on other projects that are not especially related to the problems presented here.

2. VEECH SURFACES WITH A PRYM INVOLUTION

One method for classifying Veech surfaces may be to first search for those that cannot be constructed from another Veech surface by a covering construction. Such surfaces are called *geometrically primitive*. Despite several recent major advances such as the results of [EFW18] and [EMMW20], this question of the classification of primitive Veech surfaces remains largely open. In genus two, primitive Veech surfaces have been classified in a series of articles by C. McMullen [McM05b, McM05a, McM06a]. In genus three, the classification is not complete but C. McMullen [McM06b] exhibited an infinite family of primitive Veech surfaces. The work of several authors has since shown that, apart from this family, there is only a finite number of primitive Veech surfaces (up to the action of $GL_2^+(\mathbb{R})$), see [McM21, Theorem 5.5]. This infinite family consists of surfaces having a *Prym involution* and a single singularity. A translation surface with a Prym involution can be constructed as a double cover of a *half-translation surface* (branched at the singularities), see [McM06b] or [LN13]. To search for other primitive Veech surfaces, a natural idea is to look for translation surfaces with a Prym involution, but having several singularities. In this context, E. Lanneau and M. Möller [LM19] searched for primitive Veech surfaces in the loci $Prym(2, 1, 1)$ and $Prym(2, 2)$, which is made up of translation surfaces having a Prym involution and respective singularities of order $(2, 1, 1)$ and $(2, 2)$. They prove that there is no primitive Veech surface in $Prym(2, 1, 1)$ and identify 92 candidates in $Prym(2, 2)$. In a joint work with Sam Freedman [BF22] accepted for publication at *Comptes-Rendus de l'académie des sciences*, we show:

Theorem 2.1. *There are no geometrically primitive Veech surfaces in $Prym(2, 2)$.*

Our method relies on a computer program which uses the package *Flatsurf* of *Sage* and where we compute the orbit closure of the 92 candidates. A remaining open question is then:

<p>Question 1. Are there geometrically primitive Veech surfaces in $Prym(1, 1, 1)$?</p>

3. CONNECTION POINTS AND HECKE CONTINUED FRACTIONS

In another direction, we can look for translation surfaces whose Veech group is infinitely generated. A way of building such surfaces has been exhibited by P. Hubert and T. Schmidt [HS04], who define and use the notion of *connection point*. A connection point is a (non-singular) point of the surface such that any geodesic starting at a singularity and passing through this point again meets a singularity. A vertex-to-vertex geodesic trajectory is a *saddle connection*. A (non-singular) point of the surface is said to be *periodic* if its orbit under the action of the affine diffeomorphisms of the surface is finite. A periodic point is automatically a connection point. In this context, we can ask:

Question. *Is it possible to characterize the connection points of a given translation surface?*

By a result of Boshernitzan, when the field generated by the traces of the matrices of the Veech group (called the *trace field*), is either \mathbb{Q} or quadratic over \mathbb{Q} , the connection points are exactly (after a natural normalization) the points with coordinates in the trace field. However, as soon as the degree over \mathbb{Q} of the trace field is three or more, no example of a non-periodic connection point is known. In [Bou22], published at the *Bulletin de la SMF*, we are specifically interested in the double heptagon, built from two copies of a regular heptagon whose sides are glued together, and more generally the double n -gon for $n \geq 7$ odd. These surfaces are, with the regular n -gons for even n , the original surfaces studied by Veech in his founding article [Vee89]. Their trace fields have degree at least three over \mathbb{Q} for odd $n \geq 7$. They are therefore natural examples on which one can start looking for non periodic connection points. In particular, the centers of the n -gons, which are not periodic points, are candidates to be connections points. For $n = 7$ and $n = 9$, we show that in fact this is not the case:

Theorem 3.1. *The central points of the double heptagon (as well as the double nonagon) are not connection points.*

In light of the above result, we may ask:

Question 2. Are the central points of the double n -gon connection points for $n \geq 11$?

At the moment we have an algorithm which is able to tell that the answer is no for any $n \leq 201$ (we did not conducted the tests for greater n), but a general argument remains to be found.

To prove Theorem 3.1, we use the notion of Hecke continued fractions, introduced by D. Rosen in [Ros54]. It follows from a result of T. Schmidt and M. Sheingorn [SS95] (as well as Veech dichotomy [Vee89]) that these continued fractions characterize the periodic³ directions on the double n -gon: a direction on the double n -gon is periodic if and only if the Hecke continued fraction expansion of its slope (obtained from the "next-integer" continued fraction algorithm) is finite. In [Bou22], we exhibit geodesic trajectories from starting at the singularity and passing through one of the central points whose slope has an infinite periodic Hecke continued fraction expansion. In particular, such geodesic trajectories are not closed. It is interesting to note that such examples give vertex-to-vertex billiard trajectories on the $(\frac{\pi}{2}, \frac{\pi}{7}, \frac{3\pi}{14})$ -triangle such that all other trajectories in this direction are uniquely ergodic.

In light of Theorem 3.1, we can ask:

Question 3. Do there exist non-periodic connection points on the double heptagon?

Several computer tests allow to conjecture that the answer is *yes*. Showing such a result would require to obtain an algebraic description of the periodic directions on the double heptagon. This is equivalent to have a complete description of real numbers having a finite Hecke continued fraction expansion, and in particular closely related to the following:

³A direction is periodic if any geodesic in this direction is either closed or a vertex-to-vertex trajectory.




FIGURE 2. A vertex to vertex trajectory on the $(\frac{\pi}{2}, \frac{\pi}{7}, \frac{3\pi}{14})$ -triangle in an uniquely ergodic direction.

Question 4. Does every $x \in \mathbb{Q}[2 \cos \frac{\pi}{7}]$ have either a finite or an eventually periodic Hecke continued fraction expansion?

4. INTERACTION STRENGTH ON THE MODULI SPACE OF TRANSLATION SURFACES

Finally, a significant part of my thesis deals with the algebraic intersection of curves on a Riemannian surface, defined by assigning a sign to each intersection point, and investigates more specifically the following:

Question. *How many times can two closed curves of a given length intersect?*

This question, which makes sense for any orientable surface X on which we can measure the lengths of the closed curves, can be quantified by defining:

$$\text{KVol}(X) := \text{Vol}(X) \cdot \sup_{\alpha, \beta} \frac{\text{Int}(\alpha, \beta)}{l(\alpha)l(\beta)}$$

where the supremum is taken over pairs of closed curves on X . Normalizing by the volume makes the quantity scalar invariant. According to the terminology of [Tor23], we can refer to $\text{KVol}(X)$ as the *algebraic interaction strength* of the surface X . The study of KVol goes back to D. Massart's work in [Mas97], where this KVol appears (indirectly) as a comparison constant between the stable norm and the Hodge norm in homology. The study of KVol has since been deepened by D. Massart and B. Muetzel [MM14], but many questions remain open, starting with the question of the explicit computation of KVol on simple examples of surfaces. In this context, S. Cheboui, A. Kessi and D. Massart [CKM21a, CKM21b] initiated the study of KVol on translation surfaces by explicitly calculating KVol on the Teichmüller disk of some square-tiled surfaces. In a work in progress with I. Pasquinelli, we exhibit a wide class of surfaces for which we can accurately estimate KVol :

Theorem 4.1 (B.-Pasquinelli). *Let X be a translation surface made from a collection of polygons $(P_i)_{i \in I}$ whose sides are identified by pairs. We assume:*

(H1) *Each polygon is convex with obtuse angles.*

(H2) *Sides of the same polygon are not identified.*

Then, for any pair of closed curves (α, β) on X , we have:

$$\frac{\text{Int}(\alpha, \beta)}{l(\alpha)l(\beta)} \leq \frac{1}{l_0^2}$$

where l_0 is the minimal length of a side of the polygons P_i .

This family of surfaces includes Bouw-Möller surfaces. In the case of Bouw-Möller surfaces with a single singularity, we show that the upper bound obtained is optimal (and is realized by two systoles that intersect once), thus obtaining the exact value of KVol . This result generalizes the work [BLM22], and is obtained by an *subdivision method*. Namely, we decompose closed geodesics in small pieces for which we can estimate both lengths and intersections.




FIGURE 3. On the left, a fundamental domain for the Teichmüller curve $\mathbb{H}^2/S_{m,n}$ associated to the Bouw-Möller surface $S_{m,n}$. The geodesics $\gamma_{\infty, \pm \cot(\pi/n)}$ corresponds to the surfaces having maximal KVol, and the angle θ measures the distance between X and the geodesic $\gamma_{\infty, \cot(\pi/n)}$, namely $\cosh(\text{dist}_{\mathbb{H}^2}(X, \gamma_{\infty, \cot(\pi/n)})) = \frac{1}{\sin \theta}$. On the right, the hyperbolic surface $\mathbb{H}^2/S_{m,n}$.

When the translation surface has a single singularity and all the vertices of the polygons are identified with this singularity, this result allows to compare KVol and the systolic volume, defined as the volume multiplied by the inverse of the systole squared, and it improves, for this class of surfaces, a result of Massart and Muetzel. It is interesting to wonder:

Question 5. What is the optimal constant $C > 0$ such that for every translation surface X (resp. $CAT(0)$ surface X), we have

$$\text{KVol}(X) \leq C \text{Volsys}(X)?$$

We already know that $\frac{2}{\sqrt{3}} \leq C \leq 9$, and in fact we believe that $\frac{2}{\sqrt{3}}$ is the optimal constant.

Behavior of KVol under deformation. Next, we study the behavior of KVol on translation surfaces under deformation. For this, we prove and use a sharp result of hyperbolic geometry that allows to compare the angles between two pairs of saddle connections on a translation surface. This method applies particularly well to Bouw-Möller surfaces with a single singularity, for which we have:

Theorem 4.2. *Let $m, n \geq 2$ coprime. Given $d, d' \in \mathbb{R} \cup \{\infty\} \simeq \partial\mathbb{H}^2$, let $\gamma_{d,d'}$ the geodesic of the Poincaré half plane \mathbb{H}^2 whose endpoints are d and d' . Let $X = M \cdot S_{m,n}$ be a surface in the Teichmüller disk of the Bouw-Möller surface $S_{m,n}$ which is obtained from $S_{m,n}$ by applying a matrix $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{R})$. Then, we have:*

$$\text{KVol}(X) = K_0 \cdot \frac{1}{\cosh(\text{dist}_{\mathbb{H}^2}(\frac{di+b}{ci+a}, \Gamma_{m,n} \cdot \gamma_{\infty, \pm \cot \frac{\pi}{n}}))}$$

where $K_0 > 0$ is an explicit constant which only depends on m and n , $\Gamma_{m,n}$ is the Veech group of $S_{m,n}$ and $\text{dist}_{\mathbb{H}^2}$ is the hyperbolic distance in the Poincaré half plane.

A geometric interpretation of this result is given in Figure 3. More generally, this method allows to give a criterion to know whether KVol is bounded or not as a function on the Teichmüller disk of any Veech surface. This criterion uses the notion of separatrix diagram and makes it possible to show for example that:

Theorem 4.3. *KVol is bounded on the Teichmüller disk of any algebraically primitive Veech surface.*

The algebraically primitive Veech surfaces are those whose trace field has degree equal to their genus. Algebraically primitive Veech surfaces are automatically geometrically primitive.

It would be interesting to know whether we could use this method to estimate KVol on the Teichmüller disk of any primitive Veech surface of genus two. This would allow to answer the following question for $g = 2$:

Question 6. Find the optimal constant $C(g) > 0$ such that for any translation surface of genus g with a single singularity, we have $\text{KVol}(X) > C(g)$.

We already know from [MM14] that $C(g) \geq 1$ and I have proven in [Bou23b] that $C(g) \leq g$. In fact, we conjecture from computer experiments that $C(g) = g$.

5. FUTURE DIRECTIONS

Apart from the questions described in the previous sections, there are several other questions that I am interested in investigating. Namely:

- Besides the computation of KVol on specific surfaces, it would be interesting to use probabilistic methods to relate intersection and lengths on average. More specifically, the work of Athreya–Fairchild–Masur [AFM22] counts the growth (as the length goes to infinity) of pairs of saddle connections over wedge product constraint. It would be interesting to make a similar estimate under intersection constraints. The problem of counting closed geodesics under (geometric self-)intersections constraints has already been developed in different settings by Mirzakhani [Mir08], Lalley [Lal14], or Chaubet [Cha21].
- Further, apart from the case of translation surfaces, it would be interesting to study KVol for other surfaces (for example, hyperbolic surfaces or more generally non-positively curved Riemannian surfaces), and extend Questions 5 and 6. Concerning the computation of KVol , it should be noted that Theorem 4.1 extends to any surface made from flat polygons, and the same result should also hold for surfaces made from hyperbolic polygons (however, in the case of hyperbolic polygons the angle condition is less likely to be satisfied).
- Next, D. Massart and B. Muetzel [MM14] have shown that the quantity KVol appears as a comparison constant between the stable norm and the hodge norm in homology. In a recent work, J. Kahn and A. Wright [KW22] compared the hodge norm and the Teichmüller norm. It would be interesting to relate their work to KVol .
- Finally, one can wonder to what extent Theorem 4.2 generalizes to other families of Veech surfaces. Answering this question would require to better understand the fundamental domains for Teichmüller disks of Veech surfaces. There are already several known algorithms to compute fundamental domains, see [Bow], [Ron17] and [ESS22].

Apart from the questions presented here, I am also eager to work on other projects. I especially participated in several conferences and workshops of the ANR Adyct on spectral analysis on surfaces, as well as a reading group on Teichmüller theory and pseudo-Anosov diffeomorphisms. I also enjoy working with Sagemath to test my questions on simple examples and develop intuition. For example, I recently contributed to the implementation of the algebraic intersection of saddle connections on the package Flatsurf developed by V. Delecroix, J. Rùth and P. Hooper.

REFERENCES

- [AFM22] Jayadev S. Athreya, , S. Fairchild, and Howard A. Masur. Counting pairs of saddle connections. 2022.
- [BF22] Julien Boulanger and Sam Freedman. There are no primitive Teichmüller curves in Prym(2, 2). *accepted for publication at the Comptes rendus de l'académie des sciences*, 2022.
- [BLM22] Julien Boulanger, Erwan Lanneau, and Daniel Massart. Algebraic intersection in regular polygons. *arXiv:2110.14235*, 2022.
- [Bou22] Julien Boulanger. Central points of the double heptagon are not connection points. *Bulletin de la SMF*, 150(2):459–472, 2022.
- [Bou23a] Julien Boulanger. Algebraic intersection, lengths, and veech groups. *arXiv:2309.17165*, 2023.
- [Bou23b] Julien Boulanger. Lower bound for KVol on the minimal stratum of translation surfaces. *arXiv:2310.00130*, 2023.
- [Bow] J. Bowman. Teichmüller geodesics, Delaunay triangulations, and Veech groups. *Teichmüller theory and moduli problems. Ramanujan Math. Soc.*
- [Cha21] Yann Chaubet. Closed geodesics with prescribed intersection numbers. 2021.
- [CKM21a] Smail Cheboui, Arezki Kessi, and Daniel Massart. Algebraic intersection for translation surfaces in a family of Teichmüller disks. *Bull. Soc. Math. France*, 149(4):613–640, 2021.
- [CKM21b] Smail Cheboui, Arezki Kessi, and Daniel Massart. Algebraic intersection for translation surfaces in the stratum $\mathcal{H}(2)$. *C. R. Math. Acad. Sci. Paris*, 359:65–70, 2021.
- [EFW18] Alex Eskin, Simion Filip, and Alex Wright. The algebraic hull of the Kontsevich-Zorich cocycle. *Ann. of Math.*, 2018.
- [EMMW20] Alex Eskin, Curtis McMullen, Ronen Mukamel, and Alex Wright. Billiards, quadrilaterals and moduli spaces. *J. Amer. Math. Soc.*, 2020.
- [ESS22] B. Edwards, S. Sanderson, and T. A. Schmidt. Canonical translation surfaces for computing Veech groups. *Geom. Dedicata*, 216(5), 2022.
- [HS04] Pascal Hubert and Thomas Schmidt. Infinitely generated Veech groups. *Duke mathematical journal*, 123, pages 49–69, 2004.
- [KW22] Jeremy A. Kahn and Alex Wright. Hodge and teichmüller. *Journal of Modern Dynamics*, 2022.
- [Lal14] Steven P. Lalley. Statistical regularities of self-intersection counts for geodesics on negatively curved surfaces. *Duke Mathematical Journal*, 163(6):1191 – 1261, 2014.
- [LM19] Erwan Lanneau and Martin Moeller. Non-existence and finiteness results for Teichmüller curves in prym loci. *Experimental Mathematics*, 31:621 – 636, 2019.
- [LN13] Erwan Lanneau and Duc-Manh Nguyen. $GL_2^+(\mathbb{R})$ -orbits in Prym eigenform loci. *Geometry & Topology*, 20:1359–1426, 2013.
- [Mas97] Daniel Massart. Normes stables des surfaces. *Comptes Rendus de l'Académie des Sciences - Series I - Mathematics*, 324(2):221–224, 1997.
- [McM05a] Curt McMullen. Teichmüller curves in genus two : the decagon and beyond. *Journal für die reine und angewandte Mathematik*, 2005.
- [McM05b] Curtis T. McMullen. Teichmüller curves in genus two: Discriminant and spin. *Math. Ann.*, 333(1):87–130, 2005.
- [McM06a] Curt McMullen. Teichmüller curves in genus two : torsion divisors and the ratio of sines. *Inventiones Mathematicae*, 165, pages 651–672, 2006.
- [McM06b] Curtis T. McMullen. Prym varieties and Teichmüller curves. *Duke Mathematical Journal*, 133:569–590, 2006.
- [McM21] Curtis T. McMullen. Billiards and Teichmüller curves. 2021.
- [Mir08] Maryam Mirzakhani. Growth of the number of simple closed geodesics on hyperbolic surfaces. *Annals of Mathematics*, 168:97–125, 2008.
- [MM14] Daniel Massart and Bjoern Muetzel. On the intersection form of surfaces. *Manuscr. Math.*, 143(1-2):19–49, 2014.
- [Ron17] Mukamel Ronen. Fundamental domains and generators for lattice Veech groups. *Commentarii Mathematici Helvetici*, 2017.
- [Ros54] David Rosen. A class of continued fractions associated with certain properly discontinuous groups. *Duke Mathematical Journal*, 21:549–563, 1954.
- [SS95] Thomas Schmidt and Mark Sheingorn. Length spectra of the Hecke triangle group. *Mathematische Zeitschrift*, 220:369–398, 1995.
- [Tor23] Tina Torkaman. Intersection number, length, and systole on compact hyperbolic surfaces. 2023.
- [Vee89] W. A. Veech. Teichmüller curves in moduli space, Eisenstein series and an application to triangular billiards. *Invent. Math.*, 97(3):553–583, 1989.