

Structure theorems for compact Kähler manifolds

Jean-Pierre Demailly
joint work with Frédéric Campana & Thomas Peternell

Institut Fourier, Université de Grenoble I, France
& Académie des Sciences de Paris

KSCV10 Conference, August 7–11, 2014

Goals

- Analyze the geometric structure of **projective** or **compact Kähler manifolds**

Goals

- Analyze the geometric structure of **projective** or **compact Kähler manifolds**
- As is well known since the beginning of the XXth century at least, the geometry depends on the sign of the curvature of the canonical line bundle

$$K_X = \Lambda^n T_X^*, \quad n = \dim_{\mathbb{C}} X.$$

- Analyze the geometric structure of **projective** or **compact Kähler manifolds**
- As is well known since the beginning of the XXth century at least, the geometry depends on the sign of the curvature of the canonical line bundle

$$K_X = \Lambda^n T_X^*, \quad n = \dim_{\mathbb{C}} X.$$

- $L \rightarrow X$ is pseudoeffective (**psef**) if $\exists h = e^{-\varphi}$, $\varphi \in L^1_{\text{loc}}$, (possibly singular) such that $\Theta_{L,h} = -dd^c \log h \geq 0$ on X , in the sense of currents [for X projective: $c_1(L) \in \overline{\text{Eff}}$].

- Analyze the geometric structure of **projective** or **compact Kähler manifolds**
- As is well known since the beginning of the XXth century at least, the geometry depends on the sign of the curvature of the canonical line bundle

$$K_X = \Lambda^n T_X^*, \quad n = \dim_{\mathbb{C}} X.$$


- $L \rightarrow X$ is pseudoeffective (**psef**) if $\exists h = e^{-\varphi}$, $\varphi \in L^1_{\text{loc}}$, (possibly singular) such that $\Theta_{L,h} = -dd^c \log h \geq 0$ on X , in the sense of currents [for X projective: $c_1(L) \in \overline{\text{Eff}}$].
- $L \rightarrow X$ is **positive (semipositive)** if $\exists h = e^{-\varphi}$ smooth s.t. $\Theta_{L,h} = -dd^c \log h > 0$ (≥ 0) on X .

- Analyze the geometric structure of **projective** or **compact Kähler manifolds**
- As is well known since the beginning of the XXth century at least, the geometry depends on the sign of the curvature of the canonical line bundle


$$K_X = \Lambda^n T_X^*, \quad n = \dim_{\mathbb{C}} X.$$

- $L \rightarrow X$ is pseudoeffective (**psef**) if $\exists h = e^{-\varphi}$, $\varphi \in L^1_{\text{loc}}$, (possibly singular) such that $\Theta_{L,h} = -dd^c \log h \geq 0$ on X , in the sense of currents [for X projective: $c_1(L) \in \overline{\text{Eff}}$].
- $L \rightarrow X$ is **positive (semipositive)** if $\exists h = e^{-\varphi}$ smooth s.t. $\Theta_{L,h} = -dd^c \log h > 0$ (≥ 0) on X .
- L is **nef** if $\forall \varepsilon > 0$, $\exists h_\varepsilon = e^{-\varphi_\varepsilon}$ smooth such that $\Theta_{L,h_\varepsilon} = -dd^c \log h_\varepsilon \geq -\varepsilon \omega$ on X
[for X projective: $L \cdot C \geq 0$, $\forall C$ alg. curve].


Complex curves ($n = 1$) : genus and curvature



$g = 0, K_X < 0$
(positive curvature)



$g = 1, K_X = 0$
(zero curvature)



$g > 1, K_X > 0$
(negative curvature)

$$K_X = \Lambda^n T_X^*, \quad \deg(K_X) = 2g - 2$$

Comparison of positivity concepts

Recall that for a line bundle

$$> 0 \Leftrightarrow \text{ample} \Rightarrow \text{semiample} \Rightarrow \text{semipositive} \Rightarrow \text{nef} \Rightarrow \text{psef}$$

but none of the reverse implications in red holds true.

Comparison of positivity concepts

Recall that for a line bundle

$$> 0 \Leftrightarrow \text{ample} \Rightarrow \text{semiample} \Rightarrow \text{semipositive} \Rightarrow \text{nef} \Rightarrow \text{psef}$$

but none of the reverse implications in red holds true.

Example

Let X be the rational surface obtained by blowing up \mathbb{P}^2 in 9 distinct points $\{p_i\}$ on a smooth (cubic) elliptic curve $C \subset \mathbb{P}^2$, $\mu : X \rightarrow \mathbb{P}^2$ and \hat{C} the strict transform of C .

Comparison of positivity concepts

Recall that for a line bundle

$$> 0 \Leftrightarrow \text{ample} \Rightarrow \text{semiample} \Rightarrow \text{semipositive} \Rightarrow \text{nef} \Rightarrow \text{psef}$$

but none of the reverse implications in red holds true.

Example

Let X be the rational surface obtained by blowing up \mathbb{P}^2 in 9 distinct points $\{p_i\}$ on a smooth (cubic) elliptic curve $C \subset \mathbb{P}^2$, $\mu : X \rightarrow \mathbb{P}^2$ and \hat{C} the strict transform of C . Then

$K_X = \mu^* K_{\mathbb{P}^2} \otimes \mathcal{O}(\sum E_i) \Rightarrow -K_X = \mu^* \mathcal{O}_{\mathbb{P}^2}(3) \otimes \mathcal{O}(-\sum E_i)$,
thus

$$-K_X = \mu^* \mathcal{O}_{\mathbb{P}^2}(C) \otimes \mathcal{O}(-\sum E_i) = \mathcal{O}_X(\hat{C}).$$

Comparison of positivity concepts

Recall that for a line bundle

$$> 0 \Leftrightarrow \text{ample} \Rightarrow \text{semiample} \Rightarrow \text{semipositive} \Rightarrow \text{nef} \Rightarrow \text{psef}$$

but none of the reverse implications in red holds true.

Example

Let X be the rational surface obtained by blowing up \mathbb{P}^2 in 9 distinct points $\{p_i\}$ on a smooth (cubic) elliptic curve $C \subset \mathbb{P}^2$, $\mu : X \rightarrow \mathbb{P}^2$ and \hat{C} the strict transform of C . Then

$K_X = \mu^* K_{\mathbb{P}^2} \otimes \mathcal{O}(\sum E_i) \Rightarrow -K_X = \mu^* \mathcal{O}_{\mathbb{P}^2}(3) \otimes \mathcal{O}(-\sum E_i)$,
thus

$$-K_X = \mu^* \mathcal{O}_{\mathbb{P}^2}(C) \otimes \mathcal{O}(-\sum E_i) = \mathcal{O}_X(\hat{C}).$$

One has

$$-K_X \cdot \Gamma = \hat{C} \cdot \Gamma \geq 0 \quad \text{if } \Gamma \neq \hat{C},$$

$$-K_X \cdot \hat{C} = (-K_X)^2 = (\hat{C})^2 = C^2 - 9 = 0 \Rightarrow -K_X \text{ nef.}$$

Rationally connected manifolds

In fact

$$G := (-K_X)|_{\hat{C}} \simeq \mathcal{O}_{\mathbb{P}^2|_C}(3) \otimes \mathcal{O}_C(-\sum p_i) \in \text{Pic}^0(C)$$

Rationally connected manifolds

In fact

$$G := (-K_X)|_{\hat{C}} \simeq \mathcal{O}_{\mathbb{P}^2|_C}(3) \otimes \mathcal{O}_C(-\sum p_i) \in \text{Pic}^0(C)$$

If G is a **torsion point** in $\text{Pic}^0(C)$, then one can show that $-K_X$ is semi-ample, but otherwise **it is not semi-ample**.

Rationally connected manifolds

In fact

$$G := (-K_X)|_{\hat{C}} \simeq \mathcal{O}_{\mathbb{P}^2|_C}(3) \otimes \mathcal{O}_C(-\sum p_i) \in \text{Pic}^0(C)$$

If G is a **torsion point** in $\text{Pic}^0(C)$, then one can show that $-K_X$ is semi-ample, but otherwise **it is not semi-ample**.

Brunella has shown that $-K_X$ is C^∞ semipositive if $c_1(G)$ satisfies a diophantine condition found by T. Ueda, but that otherwise it may not be semipositive (although nef).

Rationally connected manifolds

In fact

$$G := (-K_X)|_{\hat{C}} \simeq \mathcal{O}_{\mathbb{P}^2|_C}(3) \otimes \mathcal{O}_C(-\sum p_i) \in \text{Pic}^0(C)$$

If G is a **torsion point** in $\text{Pic}^0(C)$, then one can show that $-K_X$ is semi-ample, but otherwise **it is not semi-ample**.

Brunella has shown that $-K_X$ is C^∞ semipositive if $c_1(G)$ satisfies a diophantine condition found by T. Ueda, but that otherwise it may not be semipositive (although nef).

$\mathbb{P}^2 \# 9$ points is an example of rationally connected manifold:

Rationally connected manifolds

In fact

$$G := (-K_X)|_{\hat{C}} \simeq \mathcal{O}_{\mathbb{P}^2|_C}(3) \otimes \mathcal{O}_C(-\sum p_i) \in \text{Pic}^0(C)$$

If G is a **torsion point** in $\text{Pic}^0(C)$, then one can show that $-K_X$ is semi-ample, but otherwise **it is not semi-ample**.

Brunella has shown that $-K_X$ is C^∞ semipositive if $c_1(G)$ satisfies a diophantine condition found by T. Ueda, but that otherwise it may not be semipositive (although nef).

$\mathbb{P}^2 \# 9$ points is an example of rationally connected manifold:

Definition

Recall that a compact complex manifold is said to be **rationally connected** (or RC for short) if any 2 points can be joined by a chain of rational curves

Rationally connected manifolds

In fact

$$G := (-K_X)|_{\hat{C}} \simeq \mathcal{O}_{\mathbb{P}^2|_C}(3) \otimes \mathcal{O}_C(-\sum p_i) \in \text{Pic}^0(C)$$

If G is a **torsion point** in $\text{Pic}^0(C)$, then one can show that $-K_X$ is semi-ample, but otherwise **it is not semi-ample**.

Brunella has shown that $-K_X$ is C^∞ semipositive if $c_1(G)$ satisfies a diophantine condition found by T. Ueda, but that otherwise it may not be semipositive (although nef).

$\mathbb{P}^2 \# 9$ points is an example of rationally connected manifold:

Definition

Recall that a compact complex manifold is said to be **rationally connected** (or RC for short) if any 2 points can be joined by a chain of rational curves

Remark. $X = \mathbb{P}^2$ blown-up in ≥ 10 points is RC but $-K_X$ not nef.

Ex. of compact Kähler manifolds with $-K_X \geq 0$

(Recall: By Yau, $-K_X \geq 0 \Leftrightarrow \exists \omega$ Kähler with $\text{Ricci}(\omega) \geq 0$.)

Ex. of compact Kähler manifolds with $-K_X \geq 0$

(Recall: By Yau, $-K_X \geq 0 \Leftrightarrow \exists \omega$ Kähler with $\text{Ricci}(\omega) \geq 0$.)

- Ricci flat manifolds
 - Complex tori $T = \mathbb{C}^g / \Lambda$

Ex. of compact Kähler manifolds with $-K_X \geq 0$

(Recall: By Yau, $-K_X \geq 0 \Leftrightarrow \exists \omega$ Kähler with $\text{Ricci}(\omega) \geq 0$.)

- Ricci flat manifolds
 - Complex tori $T = \mathbb{C}^g / \Lambda$
 - Holomorphic symplectic manifolds S (also called hyperkähler): $\exists \sigma \in H^0(S, \Omega_S^2)$ symplectic

Ex. of compact Kähler manifolds with $-K_X \geq 0$

(Recall: By Yau, $-K_X \geq 0 \Leftrightarrow \exists \omega$ Kähler with $\text{Ricci}(\omega) \geq 0$.)

- Ricci flat manifolds

- Complex tori $T = \mathbb{C}^g / \Lambda$

- Holomorphic symplectic manifolds S (also called hyperkähler): $\exists \sigma \in H^0(S, \Omega_S^2)$ symplectic

- Calabi-Yau manifolds Y : $\pi_1(Y)$ finite and some multiple of K_Y is trivial (may assume $\pi_1(Y) = 1$ and K_Y trivial by passing to some finite étale cover

Ex. of compact Kähler manifolds with $-K_X \geq 0$

(Recall: By Yau, $-K_X \geq 0 \Leftrightarrow \exists \omega$ Kähler with $\text{Ricci}(\omega) \geq 0$.)

- Ricci flat manifolds
 - Complex tori $T = \mathbb{C}^g / \Lambda$
 - Holomorphic symplectic manifolds S (also called hyperkähler): $\exists \sigma \in H^0(S, \Omega_S^2)$ symplectic
 - Calabi-Yau manifolds Y : $\pi_1(Y)$ finite and some multiple of K_Y is trivial (may assume $\pi_1(Y) = 1$ and K_Y trivial by passing to some finite étale cover)
- the rather large class of rationally connected manifolds Z with $-K_Z \geq 0$

Ex. of compact Kähler manifolds with $-K_X \geq 0$

(Recall: By Yau, $-K_X \geq 0 \Leftrightarrow \exists \omega$ Kähler with $\text{Ricci}(\omega) \geq 0$.)

- Ricci flat manifolds
 - Complex tori $T = \mathbb{C}^g / \Lambda$
 - Holomorphic symplectic manifolds S (also called hyperkähler): $\exists \sigma \in H^0(S, \Omega_S^2)$ symplectic
 - Calabi-Yau manifolds Y : $\pi_1(Y)$ finite and some multiple of K_Y is trivial (may assume $\pi_1(Y) = 1$ and K_Y trivial by passing to some finite étale cover)
- the rather large class of rationally connected manifolds Z with $-K_Z \geq 0$
- all products $T \times \prod S_j \times \prod Y_k \times \prod Z_\ell$.

Main result. Essentially, this is a complete list !

Structure theorem for manifolds with $-K_X \geq 0$

Theorem

[Campana, D, Peternell, 2012] Let X be a compact Kähler manifold with $-K_X \geq 0$. Then

(a) \exists holomorphic and isometric splitting

$$\tilde{X} \simeq \mathbb{C}^q \times \prod Y_j \times \prod S_k \times \prod Z_\ell$$

where $Y_j =$ Calabi-Yau (holonomy $SU(n_j)$),

$S_k =$ holomorphic symplectic (holonomy $Sp(n'_k/2)$), and

$Z_\ell =$ RC with $-K_{Z_\ell} \geq 0$ (holonomy $U(n''_\ell)$).

Structure theorem for manifolds with $-K_X \geq 0$

Theorem

[Campana, D, Peternell, 2012] Let X be a compact Kähler manifold with $-K_X \geq 0$. Then

- (a) \exists holomorphic and isometric splitting

$$\tilde{X} \simeq \mathbb{C}^q \times \prod Y_j \times \prod S_k \times \prod Z_\ell$$

where $Y_j =$ Calabi-Yau (holonomy $SU(n_j)$),

$S_k =$ holomorphic symplectic (holonomy $Sp(n'_k/2)$), and

$Z_\ell =$ RC with $-K_{Z_\ell} \geq 0$ (holonomy $U(n''_\ell)$).

- (b) There exists a finite étale Galois cover $\hat{X} \rightarrow X$ such that the Albanese map $\alpha : \hat{X} \rightarrow \text{Alb}(\hat{X})$ is an (isometrically) locally trivial holomorphic fiber bundle whose fibers are products $\prod Y_j \times \prod S_k \times \prod Z_\ell$, as described in (a).

Structure theorem for manifolds with $-K_X \geq 0$

Theorem

[Campana, D, Peternell, 2012] Let X be a compact Kähler manifold with $-K_X \geq 0$. Then

- (a) \exists holomorphic and isometric splitting

$$\tilde{X} \simeq \mathbb{C}^q \times \prod Y_j \times \prod S_k \times \prod Z_\ell$$

where $Y_j =$ Calabi-Yau (holonomy $SU(n_j)$),

$S_k =$ holomorphic symplectic (holonomy $Sp(n'_k/2)$), and

$Z_\ell =$ RC with $-K_{Z_\ell} \geq 0$ (holonomy $U(n''_\ell)$).

- (b) There exists a finite étale Galois cover $\hat{X} \rightarrow X$ such that the Albanese map $\alpha : \hat{X} \rightarrow \text{Alb}(\hat{X})$ is an (isometrically) locally trivial holomorphic fiber bundle whose fibers are products $\prod Y_j \times \prod S_k \times \prod Z_\ell$, as described in (a).
- (c) $\pi_1(\hat{X}) \simeq \mathbb{Z}^{2q} \rtimes \Gamma$, Γ finite (“almost abelian” group).

Criterion for rational connectedness

Criterion

Let X be a projective algebraic n -dimensional manifold. The following properties are equivalent.

- (a) X is **rationally connected**.

Criterion for rational connectedness

Criterion

Let X be a projective algebraic n -dimensional manifold. The following properties are equivalent.

- (a) X is **rationally connected**.
- (b) For every invertible subsheaf $\mathcal{F} \subset \Omega_X^p := \mathcal{O}(\Lambda^p T_X^*)$, $1 \leq p \leq n$, \mathcal{F} is **not psef**.

Criterion for rational connectedness

Criterion

Let X be a projective algebraic n -dimensional manifold. The following properties are equivalent.

- (a) X is **rationally connected**.
- (b) For every invertible subsheaf $\mathcal{F} \subset \Omega_X^p := \mathcal{O}(\Lambda^p T_X^*)$, $1 \leq p \leq n$, \mathcal{F} is **not psef**.
- (c) For every invertible subsheaf $\mathcal{F} \subset \mathcal{O}((T_X^*)^{\otimes p})$, $p \geq 1$, \mathcal{F} is **not psef**.

Criterion for rational connectedness

Criterion

Let X be a projective algebraic n -dimensional manifold. The following properties are equivalent.

- (a) X is **rationally connected**.
- (b) For every invertible subsheaf $\mathcal{F} \subset \Omega_X^p := \mathcal{O}(\Lambda^p T_X^*)$, $1 \leq p \leq n$, \mathcal{F} is **not psef**.
- (c) For every invertible subsheaf $\mathcal{F} \subset \mathcal{O}((T_X^*)^{\otimes p})$, $p \geq 1$, \mathcal{F} is **not psef**.
- (d) For some (resp. for any) ample line bundle A on X , there exists a constant $C_A > 0$ such that

$$H^0(X, (T_X^*)^{\otimes m} \otimes A^{\otimes k}) = 0 \quad \forall m, k \in \mathbb{N}^* \text{ with } m \geq C_A k.$$

Proof of the RC criterion

Proof (essentially from Peternell 2006)

(a) \Rightarrow (d) is easy (RC implies there are many rational curves on which T_X , so $T_X^* < 0$), (d) \Rightarrow (c) and (c) \Rightarrow (b) are trivial.

Proof of the RC criterion

Proof (essentially from Peternell 2006)

(a) \Rightarrow (d) is easy (RC implies there are many rational curves on which T_X , so $T_X^* < 0$), (d) \Rightarrow (c) and (c) \Rightarrow (b) are trivial.

Thus the only thing left to complete the proof is **(b) \Rightarrow (a)**.

Proof of the RC criterion

Proof (essentially from Peternell 2006)

(a) \Rightarrow (d) is easy (RC implies there are many rational curves on which T_X , so $T_X^* < 0$), (d) \Rightarrow (c) and (c) \Rightarrow (b) are trivial.

Thus the only thing left to complete the proof is **(b) \Rightarrow (a)**.

Consider the **MRC quotient** $\pi : X \rightarrow Y$, given by the “equivalence relation $x \sim y$ if x and y can be joined by a chain of rational curves.

Proof of the RC criterion

Proof (essentially from Peternell 2006)

(a) \Rightarrow (d) is easy (RC implies there are many rational curves on which T_X , so $T_X^* < 0$), (d) \Rightarrow (c) and (c) \Rightarrow (b) are trivial.

Thus the only thing left to complete the proof is **(b) \Rightarrow (a)**.

Consider the **MRC quotient** $\pi : X \rightarrow Y$, given by the “equivalence relation $x \sim y$ if x and y can be joined by a chain of rational curves.

Then (by definition) the fibers are RC, maximal, and a result of Graber-Harris-Starr (2002) implies that **Y is not uniruled**.

Proof of the RC criterion

Proof (essentially from Peternell 2006)

(a) \Rightarrow (d) is easy (RC implies there are many rational curves on which T_X , so $T_X^* < 0$), (d) \Rightarrow (c) and (c) \Rightarrow (b) are trivial.

Thus the only thing left to complete the proof is **(b) \Rightarrow (a)**.

Consider the **MRC quotient** $\pi : X \rightarrow Y$, given by the “equivalence relation $x \sim y$ if x and y can be joined by a chain of rational curves.

Then (by definition) the fibers are RC, maximal, and a result of Graber-Harris-Starr (2002) implies that **Y is not uniruled**.

By BDPP (2004), **Y not uniruled $\Rightarrow K_Y$ psef**. Then $\pi^* K_Y \hookrightarrow \Omega_X^p$ where $p = \dim Y$, which is a contradiction unless $p = 0$, and therefore X is RC.

Generalized holonomy principle

Generalized holonomy principle

Let $(E, h) \rightarrow X$ be a hermitian holomorphic vector bundle of rank r over X compact/ \mathbb{C} . Assume that

$$\Theta_{E,h} \wedge \frac{\omega^{n-1}}{(n-1)!} = B \frac{\omega^n}{n!}, \quad B \in \text{Herm}(E, E), \quad B \geq 0 \text{ on } X.$$

Generalized holonomy principle

Generalized holonomy principle

Let $(E, h) \rightarrow X$ be a hermitian holomorphic vector bundle of rank r over X compact/ \mathbb{C} . Assume that

$$\Theta_{E,h} \wedge \frac{\omega^{n-1}}{(n-1)!} = B \frac{\omega^n}{n!}, \quad B \in \text{Herm}(E, E), \quad B \geq 0 \text{ on } X.$$

Let H the restricted holonomy group of (E, h) . Then

- (a) If there exists a psef invertible sheaf $\mathcal{L} \subset \mathcal{O}((E^*)^{\otimes m})$, then \mathcal{L} is flat and invariant under parallel transport by the connection of $(E^*)^{\otimes m}$ induced by the Chern connection ∇ of (E, h) ; moreover, H acts trivially on \mathcal{L} .

Generalized holonomy principle

Generalized holonomy principle

Let $(E, h) \rightarrow X$ be a hermitian holomorphic vector bundle of rank r over X compact/ \mathbb{C} . Assume that

$$\Theta_{E,h} \wedge \frac{\omega^{n-1}}{(n-1)!} = B \frac{\omega^n}{n!}, \quad B \in \text{Herm}(E, E), \quad B \geq 0 \text{ on } X.$$

Let H the restricted holonomy group of (E, h) . Then

- (a) If there exists a psef invertible sheaf $\mathcal{L} \subset \mathcal{O}((E^*)^{\otimes m})$, then \mathcal{L} is flat and invariant under parallel transport by the connection of $(E^*)^{\otimes m}$ induced by the Chern connection ∇ of (E, h) ; moreover, H acts trivially on \mathcal{L} .
- (b) If H satisfies $H = \text{U}(r)$, then none of the invertible sheaves $\mathcal{L} \subset \mathcal{O}((E^*)^{\otimes m})$ can be psef for $m \geq 1$.

Generalized holonomy principle

Generalized holonomy principle

Let $(E, h) \rightarrow X$ be a hermitian holomorphic vector bundle of rank r over X compact/ \mathbb{C} . Assume that

$$\Theta_{E,h} \wedge \frac{\omega^{n-1}}{(n-1)!} = B \frac{\omega^n}{n!}, \quad B \in \text{Herm}(E, E), \quad B \geq 0 \text{ on } X.$$

Let H the restricted holonomy group of (E, h) . Then

- (a) If there exists a psec invertible sheaf $\mathcal{L} \subset \mathcal{O}((E^*)^{\otimes m})$, then \mathcal{L} is flat and invariant under parallel transport by the connection of $(E^*)^{\otimes m}$ induced by the Chern connection ∇ of (E, h) ; moreover, H acts trivially on \mathcal{L} .
- (b) If H satisfies $H = \text{U}(r)$, then none of the invertible sheaves $\mathcal{L} \subset \mathcal{O}((E^*)^{\otimes m})$ can be psec for $m \geq 1$.

Proof. $\mathcal{L} \subset \mathcal{O}((E^*)^{\otimes m})$ which has trace of curvature ≤ 0 while $\Theta_{\mathcal{L}} \geq 0$, use Bochner formula.



Generically nef vector bundles

Definition

Let X compact Kähler manifold, $\mathcal{E} \rightarrow X$ torsion free sheaf.

(a) \mathcal{E} is **generically nef with respect to the Kähler class ω** if

$$\mu_{\omega}(\mathcal{S}) \geq 0$$

for all torsion free quotients $\mathcal{E} \rightarrow \mathcal{S} \rightarrow 0$.

If \mathcal{E} is ω -generically nef for all ω , we simply say that \mathcal{E} is **generically nef**.

Generically nef vector bundles

Definition

Let X compact Kähler manifold, $\mathcal{E} \rightarrow X$ torsion free sheaf.

- (a) \mathcal{E} is **generically nef with respect to the Kähler class ω** if

$$\mu_{\omega}(\mathcal{S}) \geq 0$$

for all torsion free quotients $\mathcal{E} \rightarrow \mathcal{S} \rightarrow 0$.

If \mathcal{E} is ω -generically nef for all ω , we simply say that \mathcal{E} is **generically nef**.

- (b) Let

$$0 = \mathcal{E}_0 \subset \mathcal{E}_1 \subset \dots \subset \mathcal{E}_s = \mathcal{E}$$

be a filtration of \mathcal{E} by torsion free coherent subsheaves such that the quotients $\mathcal{E}_{i+1}/\mathcal{E}_i$ are ω -stable subsheaves of $\mathcal{E}/\mathcal{E}_i$ of maximal rank. We call such a sequence a **refined Harder-Narasimhan (HN) filtration w.r.t. ω** .

Characterization of generically nef vector bundles

It is a standard fact that refined HN-filtrations always exist, moreover

$$\mu_\omega(\mathcal{E}_i/\mathcal{E}_{i-1}) \geq \nu_\omega(\mathcal{E}_{i+1}/\mathcal{E}_i)$$

for all i .

Proposition

Let (X, ω) be a compact Kähler manifold and \mathcal{E} a torsion free sheaf on X . Then \mathcal{E} is ω -generically nef if and only if

$$\mu_\omega(\mathcal{E}_{i+1}/\mathcal{E}_i) \geq 0$$

for some refined HN-filtration.

Proof. Easy arguments on filtrations. □

Theorem

(Junyan Cao, 2013) Let X be a compact Kähler manifold with $-K_X$ nef. Then the tangent bundle T_X is ω -generically nef for all Kähler classes ω .

Proof. use the fact that $\forall \varepsilon > 0, \exists$ Kähler metric with $\text{Ricci}(\omega_\varepsilon) \geq -\varepsilon \omega_\varepsilon$ (Yau, DPS 1995).

Theorem

(Junyan Cao, 2013) Let X be a compact Kähler manifold with $-K_X$ nef. Then the tangent bundle T_X is ω -generically nef for all Kähler classes ω .

Proof. use the fact that $\forall \varepsilon > 0, \exists$ Kähler metric with $\text{Ricci}(\omega_\varepsilon) \geq -\varepsilon \omega_\varepsilon$ (Yau, DPS 1995).

From this, one can deduce

Theorem

Let X be a compact Kähler manifold with nef anticanonical bundle. Then the bundles $T_X^{\otimes m}$ are ω -generically nef for all Kähler classes ω and all positive integers m . In particular, the bundles $S^k T_X$ and $\bigwedge^p T_X$ are ω -generically nef.

A lemma on sections of contravariant tensors

Lemma

Let (X, ω) be a compact Kähler manifold with $-K_X$ nef and

$$\eta \in H^0(X, (\Omega_X^1)^{\otimes m} \otimes \mathcal{L})$$

where \mathcal{L} is a **numerically trivial** line bundle on X .

A lemma on sections of contravariant tensors

Lemma

Let (X, ω) be a compact Kähler manifold with $-K_X$ nef and

$$\eta \in H^0(X, (\Omega_X^1)^{\otimes m} \otimes \mathcal{L})$$

where \mathcal{L} is a **numerically trivial** line bundle on X .

Then the filtered parts of η w.r.t. the refined HN filtration are **parallel** w.r.t. the Bando-Siu metric in the 0 slope parts, and the < 0 slope parts vanish.

A lemma on sections of contravariant tensors

Lemma

Let (X, ω) be a compact Kähler manifold with $-K_X$ nef and

$$\eta \in H^0(X, (\Omega_X^1)^{\otimes m} \otimes \mathcal{L})$$

where \mathcal{L} is a **numerically trivial** line bundle on X .

Then the filtered parts of η w.r.t. the refined HN filtration are **parallel** w.r.t. the Bando-Siu metric in the 0 slope parts, and the < 0 slope parts vanish.

Proof. By Cao's theorem there exists a refined HN-filtration

$$0 = \mathcal{E}_0 \subset \mathcal{E}_1 \subset \dots \subset \mathcal{E}_s = T_X^{\otimes m}$$

with ω -stable quotients $\mathcal{E}_{i+1}/\mathcal{E}_i$ such that $\mu_\omega(\mathcal{E}_{i+1}/\mathcal{E}_i) \geq 0$ for all i . Then we use the fact that any section in a (semi-)negative slope reflexive sheaf $\mathcal{E}_{i+1}/\mathcal{E}_i \otimes \mathcal{L}$ is parallel w.r.t. its Bando-Siu metric (resp. vanishes).



Smoothness of the Albanese morphism (after Cao)

Theorem (J.Cao 2013, D-Peternell, 2014)

Non-zero holomorphic p -forms on a compact Kähler manifold X with $-K_X$ nef **vanish only on the singular locus of the refined HN filtration of T^*X .**

Smoothness of the Albanese morphism (after Cao)

Theorem (J.Cao 2013, D-Peternell, 2014)

Non-zero holomorphic p -forms on a compact Kähler manifold X with $-K_X$ nef **vanish only on the singular locus of the refined HN filtration of T^*X .**

This implies the following result essentially due to J.Cao.

Corollary

Let X be a compact Kähler manifold with nef anticanonical bundle. Then the Albanese map $\alpha : X \rightarrow \text{Alb}(X)$ is a **submersion** on the complement of the HN filtration singular locus in X [$\Rightarrow \alpha$ surjects onto $\text{Alb}(X)$ (Paun 2012)].

Proof. The differential $d\alpha$ is given by $(d\eta_1, \dots, d\eta_q)$ where (η_1, \dots, η_q) is a basis of 1-forms, $q = \dim H^0(X, \Omega_X^1)$.

Smoothness of the Albanese morphism (after Cao)

Theorem (J.Cao 2013, D-Peternell, 2014)

Non-zero holomorphic p -forms on a compact Kähler manifold X with $-K_X$ nef **vanish only on the singular locus of the refined HN filtration of T^*X .**

This implies the following result essentially due to J.Cao.

Corollary

Let X be a compact Kähler manifold with nef anticanonical bundle. Then the Albanese map $\alpha : X \rightarrow \text{Alb}(X)$ is a **submersion** on the complement of the HN filtration singular locus in X [$\Rightarrow \alpha$ surjects onto $\text{Alb}(X)$ (Paun 2012)].

Proof. The differential $d\alpha$ is given by $(d\eta_1, \dots, d\eta_q)$ where (η_1, \dots, η_q) is a basis of 1-forms, $q = \dim H^0(X, \Omega_X^1)$.

Cao's thm \Rightarrow rank of $(d\eta_1, \dots, d\eta_q)$ is $= q$ generically.



Emerging general picture

Conjecture (known for X projective!)

Let X be compact Kähler, and let $X \rightarrow Y$ be the MRC fibration (after taking suitable blow-ups to make it a genuine morphism). Then K_Y is psef.

Emerging general picture

Conjecture (known for X projective!)

Let X be compact Kähler, and let $X \rightarrow Y$ be the MRC fibration (after taking suitable blow-ups to make it a genuine morphism). Then K_Y is psef.

Proof ? Take the part of slope > 0 in the HN filtration of T_X w.r.t. to classes in the dual of the psef cone, and apply duality.

Emerging general picture

Conjecture (known for X projective!)

Let X be compact Kähler, and let $X \rightarrow Y$ be the MRC fibration (after taking suitable blow-ups to make it a genuine morphism). Then K_Y is psef.

Proof ? Take the part of slope > 0 in the HN filtration of T_X w.r.t. to classes in the dual of the psef cone, and apply duality.

Remaining problems

- Develop the theory of singular Calabi-Yau and singular holomorphic symplectic manifolds.

Emerging general picture

Conjecture (known for X projective!)

Let X be compact Kähler, and let $X \rightarrow Y$ be the MRC fibration (after taking suitable blow-ups to make it a genuine morphism). Then K_Y is psef.

Proof ? Take the part of slope > 0 in the HN filtration of T_X w.r.t. to classes in the dual of the psef cone, and apply duality.

Remaining problems

- Develop the theory of singular Calabi-Yau and singular holomorphic symplectic manifolds.
- Show that the “slope 0” part corresponds to blown-up tori, singular Calabi-Yau and singular holomorphic symplectic manifolds (as fibers and targets).

Emerging general picture

Conjecture (known for X projective!)

Let X be compact Kähler, and let $X \rightarrow Y$ be the MRC fibration (after taking suitable blow-ups to make it a genuine morphism). Then K_Y is psef.

Proof ? Take the part of slope > 0 in the HN filtration of T_X w.r.t. to classes in the dual of the psef cone, and apply duality.

Remaining problems

- Develop the theory of singular Calabi-Yau and singular holomorphic symplectic manifolds.
- Show that the “slope 0” part corresponds to blown-up tori, singular Calabi-Yau and singular holomorphic symplectic manifolds (as fibers and targets).
- The rest of T_X (slope < 0) yields a general type quotient.

Thank you for your attention!

