

TWO QUESTIONS ON MAPPING CLASS GROUPS

LOUIS FUNAR

ABSTRACT. We show that central extensions of the mapping class group M_g of the closed orientable surface of genus g by \mathbb{Z} are residually finite. Further we give rough estimates of the largest $N = N_g$ such that homomorphisms from M_g to $SU(N)$ have finite image. In particular, homomorphisms of M_g into $SL(\lfloor \sqrt{g+1} \rfloor, \mathbb{C})$ have finite image. Both results come from properties of quantum representations of mapping class groups.

2000 MSC Classification: 57 M 07, 20 F 36, 20 F 38, 57 N 05.

Keywords: Mapping class group, central extension, quantum representation.

1. INTRODUCTION AND STATEMENTS

Set Σ_g^r for the orientable surface of genus g with r punctures. We denote by M_g^r the mapping class group of Σ_g^r , namely the group of isotopy classes of homeomorphisms that fix the punctures.

The following answers Question 6.4 of Farb (see Chapter 2 of [7]).

Proposition 1.1. *The central extensions of the mapping class group M_g (or the punctured mapping class group M_g^1 , for $g \geq 4$) by \mathbb{Z} are residually finite.*

Remark 1.1. The universal central extension $\widetilde{M}_g(1)$ surjects onto the universal central extension $\widetilde{Sp}(2g, \mathbb{Z})$ of the (integral) symplectic group, whose class is the Maslov class (generating $H^2(Sp(2g, \mathbb{Z}))$). It is known that $\widetilde{Sp}(2g, \mathbb{Z})$ is the pull-back of $Sp(2g, \mathbb{Z})$ into the universal covering $\widetilde{Sp}(2g, \mathbb{R})$ of the real symplectic group.

By a result of Deligne (see [6]) the extension $\widetilde{Sp}(2g, \mathbb{Z})$, for $g \geq 2$, is *not* residually finite since any finite index subgroup of it contains $2\mathbb{Z}$, where \mathbb{Z} is the central kernel $\widetilde{Sp}(2g, \mathbb{Z}) \rightarrow Sp(2g, \mathbb{Z})$. The same holds, more generally, for some other arithmetic groups having the congruence subgroup property.

The method of proof uses quantum representations of mapping class groups.

Definition 1.1. *The group Γ has property F_n if all homomorphisms $\Gamma \rightarrow PU(n)$ have finite image. Moreover, the group Γ has property F if it has property F for every n .*

Observe that the property F is inherited by finite index subgroups.

Remark 1.2. Let G be a connected, semi-simple, almost \mathbb{Q} -simple algebraic \mathbb{Q} -group and Γ an arithmetic lattice in G . If $G_{\mathbb{R}}$ has real rank at least 2 and $G^{ad}(\mathbb{R})$ has no compact factor then Γ has property F . This follows from ([17], Chap. VIII, Thm.B) for $K = \mathbb{Q}, l = \mathbb{R}, S$ containing only the Archimedean place of \mathbb{Q} and $\mathbf{H} = PO(n)$. In particular, any discrete group Γ commensurable with $Sp(2g, \mathbb{Z})$ for $g \geq 2$ or to $SL(2, \mathcal{O})$, where \mathcal{O} is the ring of integers in a totally real number field of degree at least 2, has property F .

Mapping class groups have not property F . It is therefore interesting to understand whether they have property F_n for some n . This is related to a question of Farb in [7] concerning linear representations in small degree. The previous remark shows that we cannot use unitary representations of M_g that factor

through $Sp(2g, \mathbb{Z})$, as the later group has no finite dimensional unitary representations with infinite image. Our second result states as follows:

Proposition 1.2. *The maximal number N_g for which M_g has property F_{N_g} satisfies:*

$$\sqrt{g+1} \leq N_g < \begin{cases} 5^{g/2} F_{g-1}, & \text{if } g \text{ is even,} \\ 5^{(g-1)/2} (F_g + F_{g-2}), & \text{if } g \text{ is odd,} \end{cases}$$

where F_j is the Fibonacci sequence, defined by $F_0 = 0, F_1 = 1$ and the recurrence $F_{n+1} = F_n + F_{n-1}$, for $n \geq 1$. Moreover, the upper bounds are valid for any finite index subgroup of M_g .

Corollary 1.1. *Every homomorphism $M_g \rightarrow SL([\sqrt{g+1}], \mathbb{C})$ has finite image, if $g \geq 1$.*

It is likely that N_g behaves like an exponential for large g . This seems difficult to check because very few unitary representations of M_g are known. On the other hand one might expect that the maximal n with the property that every homomorphism $M_g \rightarrow SL(n, \mathbb{C})$ has finite image is a linear function on g .

Notice that groups having homomorphisms with infinite image into $SL(2, \mathbb{C})$ have not the property T of Kazhdan. However, M_g has no such representations, if $g \geq 3$, by the Corollary above.

Results of similar flavor were proved in [10] where it is shown that representations $M_g \rightarrow GL(2\sqrt{g-1}, \mathbb{C})$ cannot be faithful and in [3] where it is shown that the image of an element of M_g under a representation into $GL(g, \mathbb{C})$ should have algebraic eigenvalues.

One inequality above is an immediate consequence of a theorem of Bridson ([3]) concerning the property FA_n , which was introduced by Farb in [9]. The second inequality comes from the existence of quantum representations of M_g with infinite image ([13]).

2. PROOF OF PROPOSITION 1.1

We prove the claim for the universal central extension first. This is known when $g = 1$ since the universal central extension is isomorphic to the braid group B_3 .

An important result due independently to Andersen ([1]) and to Freedman, Walker and Zhang ([11]) states that the $SU(2)$ TQFT representation of the mapping class group is *asymptotically faithful*. Specifically, there is a sequence of representations ρ_k (indexed by an integer k , called the level) $\rho_k : M_g \rightarrow PU(N(k, g))$ into the projective unitary group of dimension $N(k, g)$ (for some $N(k, g)$ depending exponentially on k) such that the intersection of the kernels $\bigcap_{k \geq 2} \ker \rho_k$ is trivial for $g \geq 3$, and respectively the center of the mapping class group M_2 (which is a group of order two generated by hyperelliptic involution), when $g = 2$. Moreover, for $g = 2$ we can use the $SU(n)$ TQFT representation, with $n \geq 3$, for which the intersection of the kernels above is trivial (see [1]). When using this result we will say that we make use of the asymptotic faithfulness (of the quantum representations).

Each quantum representation is a projective representation which lifts to a linear representation $\tilde{\rho}_k : \widetilde{M}_g(12) \rightarrow U(N(k, g))$ of the central extension $\widetilde{M}_g(12)$ of the mapping class group M_g by \mathbb{Z} . The later representation corresponds to invariants of 3-manifolds with a p_1 -structure. Masbaum, Roberts ([18]) and Gervais ([15]) gave a precise description of this extension. Namely, the cohomology class $c_{\widetilde{M}_g(12)} \in H^2(M_g, \mathbb{Z})$ associated to this extension equals 12 times the signature class χ . It is known (see [16]) that the group $H^2(M_g)$ is generated by χ , when $g \geq 2$. Recall that χ is the class of one fourth the Meyer signature cocycle.

Observe that the ρ_k action of the center of $\widetilde{M}_g(12)$ is by roots of unity of order $2k$ (see [18] for the explicit formula). In fact, this action corresponds to the change of the p_1 -structure of a 3-manifold and it is well-known that the quantum invariant changes by a root of unity of order $2k$. Thus every element of the center acts non-trivially via $\tilde{\rho}_k$, for large enough k , so that the representations of $\widetilde{M}_g(12)$ are also asymptotically faithful. This implies that $\widetilde{M}_g(12)$ is residually finite. In fact, let $a \in \widetilde{M}_g(12)$ be any element $a \neq 1$. By the asymptotic faithfulness there exists some level k so that $\tilde{\rho}_k(a) \in U(N(k, g))$ is non-trivial. The subgroup $\tilde{\rho}_k(\widetilde{M}_g(12)) \subset U(N(k, g))$ is a discrete linear group and thus, by a classical theorem of Malcev, it is residually finite. In particular, there exists a homomorphism of $\tilde{\rho}_k(\widetilde{M}_g(12))$ onto

some finite group sending $\tilde{\rho}_k(a)$ into a non-trivial element. This shows that every non-trivial element of $\widetilde{M}_g(12)$ is detected by some homomorphism into some finite group.

The universal central extension is $\widetilde{M}_g(1)$, where $\widetilde{M}_g(n)$ denotes the central extension by \mathbb{Z} whose class is $c_{\widetilde{M}_g(n)} = n\chi$. It is immediate from their explicit presentations (see [15]) that $\widetilde{M}_g(d)$ embeds into $\widetilde{M}_g(n)$ if d divides $n \neq 0$. Such an embedding sends the generator z of the center into $z^{n/d}$. In particular, $\widetilde{M}_g(1)$ embeds in $\widetilde{M}_g(12)$ and thus the universal central extension is residually finite.

Now, an arbitrary central extension of M_g by \mathbb{Z} is either trivial and hence residually finite, or else isomorphic to $\widetilde{M}_g(n)$, for some $n \in \mathbb{Z} \setminus \{0\}$. We observed above that there is an injective homomorphism $\widetilde{M}_g(1) \rightarrow \widetilde{M}_g(n)$, which sends the central element z into z^n . Moreover, the image is a normal subgroup of $\widetilde{M}_g(n)$. In particular, we have $\widetilde{M}_g(n)/\widetilde{M}_g(1) = \mathbb{Z}/n\mathbb{Z}$. This implies that $\widetilde{M}_g(n)$ is residually finite. In fact, any element of $\widetilde{M}_g(n)$ which is not detected by the homomorphism onto $\mathbb{Z}/n\mathbb{Z}$ belongs to $\widetilde{M}_g(1)$. Inducting finite group representations from $\widetilde{M}_g(1)$ to $\widetilde{M}_g(n)$ we obtain finite group representations of the later detecting every non-trivial element of $\widetilde{M}_g(1)$. This proves the claim.

Remark 2.1. Freedman, Walker and Zhang already observed in [11] that a simple consequence of the asymptotic faithfulness is that M_g is residually finite.

Remark 2.2. This proof works more generally for the punctured mapping class group M_g^1 and for those extensions whose cohomology classes are of the form $n\chi + e$, for some $n \in \mathbb{Z}$. Recall that $H^2(M_g^1) = \mathbb{Z}\chi \oplus \mathbb{Z}e$, where χ is the signature class and e is the class associated to the puncture, for $g \geq 4$ (see [16]).

Remark 2.3. Notice that there exist quantum type representations of $Sp(2g, \mathbb{Z})$, for instance those associated to the monodromy of level k theta functions in the $U(1)$ gauge theory (see e.g. [12, 14]). Again these are only projective unitary representations which lift to unitary representations of some central extension $\rho_{Sp,k} : Sp(2g, \mathbb{Z})(4) \rightarrow U(k^g)$. Here $Sp(2g, \mathbb{Z})(4)$ is the central extension of $Sp(2g, \mathbb{Z})$ by \mathbb{Z} whose class is 4 times the Maslov class. However, these representations factor through the integer metaplectic group. Further the generator of the kernel of $Sp(2g, \mathbb{Z})(4) \rightarrow Sp(2g, \mathbb{Z})$ acts as the multiplication by a root of unity of order 8, for any level k . Thus the intersection of $\cap_{k \geq 2} \rho_{Sp,k}$ is $2\mathbb{Z}$, and the result of Deligne cited above shows that this is sharp.

3. PROOF OF PROPOSITION 1.2

We consider first the following notion introduced by Farb in [9]:

Definition 3.1. *Let $n \geq 1$. A group Γ has property FA_n if any isometric action on any n -dimensional CAT(0) cell complex X has a fixed point.*

Observe that property FA_1 corresponds to Serre's property FA , which asks that any action without inversions of Γ on a real tree should fix a vertex. Notice that Kazhdan groups have property FA . Moreover if a group has property FA_n then it has property FA_k for all $k < n$. It is known (see [9]) that a group Γ with property FA_{n-1} has n -integral representation type, namely the eigenvalues of matrices in $\rho(\Gamma)$, for a homomorphism $\rho : \Gamma \rightarrow GL(n, K)$ with K a field, are algebraic integers if $\text{char}(K) = 0$. Moreover, there are only finitely many conjugacy classes of irreducible representations of Γ into $GL(n, K)$, for an algebraically closed field K .

Culler and Vogtmann proved that M_g has property FA_1 in [5]. In [7] one asks to estimate the maximal $n = n(g)$ for which M_g has property FA_n .

There is a version of FA_n , namely the strong FA_n (which implies FA_n), in which one considers complete CAT(0) spaces and semi-simple actions. It is proved by Bridson in ([3], see also [2]) that M_g has strong FA_g . Moreover it is known that M_g acts (faithfully if $g > 2$) by semi-simple isometries on the completion of the Teichmüller space with the Weil-Petersson metric, which has dimension $6g - 6$. Thus $g \leq n(g) \leq 6g - 7$.

The key point is to relate the property FA_n to the finiteness of unitary representations. Specifically, we have the following:

Proposition 3.1. *If Γ is a finitely generated group with property FA_{n^2-1} then the representations $\Gamma \rightarrow SL(n, \mathbb{C})$ have finite image.*

Proof. Let $\bar{\Gamma}$ be the image of Γ under some homomorphism into $SL(n, \mathbb{C})$. A finitely generated subgroup $\bar{\Gamma}$ of $SL(n, \mathbb{C})$ lies into some $SL(n, A)$, where A is a finitely generated \mathbb{Q} -algebra contained in \mathbb{C} . Let $\varphi : A \rightarrow \bar{\mathbb{Q}}$ be a specialization of A , which induces a morphism $\varphi : SL(n, A) \rightarrow SL(n, \bar{\mathbb{Q}})$. The image $\varphi(\bar{\Gamma})$ belongs then to some $SL(n, K)$, where K is a finite extension of $\bar{\mathbb{Q}}$.

Lemma 3.1. *If all specializations $\varphi(\bar{\Gamma})$ are finite then $\bar{\Gamma}$ is finite.*

Proof. Jordan's theorem says that there is some $f(n)$ such that any finite subgroup of $GL(n, K)$ has a normal abelian subgroup of index at most $f(n)$. The intersection of all subgroups of $\bar{\Gamma}$ of index at most $f(n)$ is then a finite index subgroup $U \subset \bar{\Gamma}$ such that $\varphi([U, U]) = 1$ for every specialization φ . Since specializations φ separate the points of A we have $[U, U] = 1$. Therefore there exists a finite index normal abelian subgroup U of $\bar{\Gamma}$. If U is finite then $\bar{\Gamma}$ will be finite and we are done.

Let us assume from now on that U is infinite. Since U is finitely generated abelian there is an infinite order element $Z \in U$. The following lemma will show that there exists a specialization φ such that $\varphi(Z)$ is of infinite order, contradicting our assumptions. Thus U should be finite abelian and the result will follow.

Lemma 3.2. *Let Z be a matrix with entries in a finitely generated \mathbb{Q} -algebra A contained in \mathbb{C} . Suppose that for every number field K and any ring homomorphism $\varphi : A \rightarrow K$ the image $\varphi(Z)$ is a matrix of finite order. Then Z has finite order.*

Proof. By Noether's normalization lemma (see [19], p.63) there exist algebraically independent elements $\xi_1, \xi_2, \dots, \xi_p \in A$ such that A is an integral extension of the purely transcendental extension $B = \mathbb{Q}[\xi_1, \dots, \xi_p]$. Moreover ξ_1, \dots, ξ_p form a transcendence basis for the field of fractions of A over \mathbb{Q} .

Let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of the matrix Z . We will prove that λ_j are roots of unity. First λ_j are integral over A because they are the roots of the characteristic polynomial of Z , which is a monic polynomial with coefficients in A . The integrality is transitive and hence λ_i are integral over B . Thus λ_j satisfies an algebraic equation $P_j(\lambda_j) = 0$, where $P_j \in B[X]$ is the minimal polynomial of λ_j over B . The polynomial P_j is monic and irreducible because B is a unique factorization domain. This implies that, if we consider P_j as a polynomial from $\mathbb{Q}[\xi_1, \dots, \xi_p, X]$ then it is still an irreducible polynomial in the $p+1$ variables ξ_1, \dots, ξ_p, X .

If $p=0$ then the fractions field of A is a number field and thus λ_i should be roots of unity.

Let us assume henceforth that $p \geq 1$. Observe that any specialization $\varphi : B \rightarrow \bar{\mathbb{Q}}$ can be lifted (not uniquely) to a specialization $\varphi : D \rightarrow \bar{\mathbb{Q}}$ of a finite extension D of B . First, specializations of B can be extended to possibly infinite specializations (see [20], Thm.6, p.31) of any extension D of B . Moreover, the extended specialization is finite on any element of D which is integral over B (see [20], Prop.22, p.41). In particular, any specialization of B extends to $A[\lambda_1, \dots, \lambda_n]$. On the other hand, observe that any specialization φ of B corresponds to prescribing the values of $\varphi(\xi_j) \in \bar{\mathbb{Q}}$ arbitrarily.

Hilbert's irreducibility theorem states that there exist infinitely many (actually a Zariski dense set of) specializations $\varphi : B \rightarrow \mathbb{Q}$ such that the polynomials $\varphi(P_j) \in \mathbb{Q}[X]$ are still irreducible. Since $\varphi(Z)$ is of finite order, each $\varphi(\lambda_j)$ is a root of unity so that $\varphi(P_j)$ should be a cyclotomic polynomial. The degree of $\varphi(P_j)$ is the degree d_j of P_j , since these are monic polynomials. But there are only finitely many cyclotomic polynomials of given degree. Let S be the finite family of coefficients of all cyclotomic polynomials of degree smaller or equal to $\max(d_1, \dots, d_n)$. It suffices then to choose some specialization φ of B for which one coefficient of some $\varphi(P_j)$ does not belong to S . For instance it suffices to choose a specialization for which some coefficient of $\varphi(P_j)$ is not in \mathbb{Z} , because cyclotomic polynomials have coefficients in \mathbb{Z} . This is possible unless all coefficients of the polynomials P_j are independent on the ξ_1, \dots, ξ_p . This might happen only if $P_j \in \mathbb{Q}[X]$, namely if all its coefficients, which are elements of

$\mathbb{Q}[\xi_1, \dots, \xi_p]$, are actually constant. But in this case all λ_j are algebraic integers. This contradicts the fact that the transcendence degree of the fractions field of A was supposed to be $p \geq 1$. Therefore all eigenvalues λ_j of Z are roots of unity.

An alternative argument is as follows. The set of \mathbb{C} -valued specializations $\varphi : A[\lambda_1, \dots, \lambda_n] \rightarrow \mathbb{C}$ is an irreducible affine algebraic variety of dimension p and λ_j is a rational function on it. If λ_j is a root of unity for any $\overline{\mathbb{Q}}$ -valued specialization then $|\lambda_j|$ is identically 1. But a bounded regular function on a irreducible complex algebraic variety should be constant. This implies that all λ_j are algebraic integers and we conclude as above.

Eventually, it suffices to show that Z is diagonalizable. Consider the Jordan-Chevalley decomposition $Z = D + N$, where D is semi-simple, N is nilpotent and $DN = ND$. The entries of the matrices D and N belong to the field of fractions of A (see [4], Thm.7, p.71-72). Let $a \in A$ be the least common multiple of denominators arising in the entries of D and N . Every specialization φ of A with the property that $\varphi(a) \neq 0$ extends uniquely to a specialization, still denoted φ , of the localization of A at a . In particular, it makes sense to consider $\varphi(D)$ and $\varphi(N)$. Therefore $\varphi(Z) = \varphi(D) + \varphi(N)$ is the Jordan-Chevalley decomposition of $\varphi(Z)$. But the minimal polynomial of $\varphi(Z)$ divides $X^s - 1$, where s is the order of $\varphi(Z)$. This implies that the minimal polynomial has distinct roots and so $\varphi(Z)$ is semi-simple. The uniqueness of the Jordan-Chevalley decomposition yields then $\varphi(N) = 0$. Since this holds for any specialization φ such that $\varphi(a) \neq 0$ and such specializations separate the points of A we derive that $N = 0$. Thus Z is diagonalizable and hence of finite order, as claimed. \square

Remark 3.1. We could also use ([3], Prop. 6.1) which says that the image in $GL(g, \mathbb{C})$ of an element of a finitely generated group with strong property FA_g has algebraic eigenvalues. However, lemma 3.2 can be applied to more general situations, since there is no assumption on Z . \square

It suffices now to show that, for any specialization φ , the image $G = \varphi(\overline{\Gamma})$ is finite. Observe that, if Γ has property FA_{n-1} , then $G = \varphi(\overline{\Gamma})$ has also property FA_{n-1} . We will show that:

Lemma 3.3. *Let K be a number field. Then a finitely generated subgroup $G \subset SL(n, K)$ with property FA_{n^2-1} should be finite.*

Proof. We prove that, for any embedding of K into a local field K_v the image of G in $SL(n, K_v)$ is precompact.

If $G \subset SL(n, K)$ is a finitely generated subgroup with property FA_{n-1} then its image in $SL(n, K_v)$ is precompact, for each non-Archimedean valuation v of K . In fact, G acts on the Bruhat-Tits building associated to $SL(n, K_v)$, which is a $(n-1)$ -dimensional CAT(0) cell complex. The G -action has a fixed point because G has property FA_{n-1} and hence G is contained in the stabilizer of a vertex, which is a compact subgroup.

In what concerns the Archimedean valuations it suffices to consider the complex ones. But $SL(n, \mathbb{C})$ acts on the symmetric space $SL(n, \mathbb{C})/SU(n)$ of non-compact type and real dimension $n^2 - 1$. Since this space is CAT(0) and G has property FA_{n^2-1} it follows that the image of G into $SL(n, \mathbb{C})$ is contained in the stabilizer $U(n)$ for any complex valuation inducing an embedding $K \rightarrow \mathbb{C}$.

Eventually recall that $SL(n, K)$ embeds as a discrete subgroup of the special linear group $SL(n, A_K)$ over the adèles ring A_K of K . By above G is discrete and precompact into $SL(n, A_K)$ and hence finite. \square

This proves Proposition 3.1. \square

Remark 3.2. If G is a subgroup of $U(n) \cap SL(n, \mathbb{Q})$ with property FA_{n-1} then G is finite. This follows from above by using the fact that there is an unique complex Archimedean valuation on \mathbb{Q} , and one knows already that G is contained in the compact group $U(n)$. In particular, if Γ has property FA_{n-1} then the image of any homomorphism $\Gamma \rightarrow U(n) \cap SL(n, \mathbb{Q})$ is finite.

End of the proof of Proposition 1.2. The result of proposition 3.1 holds also for representations into $PSL(n, \mathbb{C})$ and a fortiori for representations into $PU(n)$. Since M_g has property FA_g we derive the lower bound inequality.

Consider now the smallest (projective) quantum representation $M_g \rightarrow PU(d_g)$ with infinite image, for $g \geq 2$. This is the $SO(3)$ quantum representation in level 5 (see e.g. [13]), whose dimension d_g is given by the Verlinde formula:

$$d_g = \left(\frac{5}{4}\right)^{g-1} \sum_{j=1}^4 \left(\sin \frac{2\pi j}{5}\right)^{2-2g} = \begin{cases} 5^{g/2} F_{g-1}, & \text{if } g \text{ is even,} \\ 5^{(g-1)/2} (F_g + F_{g-2}) & \end{cases}$$

where F_j is the Fibonacci sequence $F_0 = 0, F_1 = 1, F_{n+1} = F_n + F_{n-1}$. For instance $d_2 = 5$. These mapping class group representations come from the so-called Fibonacci TQFT.

Moreover, it is clear that the upper bound holds for any finite index subgroup of M_g . In fact the image of a finite index subgroup of the mapping class group by the quantum representation is still infinite. This proves the claim.

Remark 3.3. The property FA_{n-1} is not inherited by the finite index subgroups. Actually M_2 has a finite index subgroup which surjects onto a free non-abelian group and hence it has not property FA_1 . The situation is subtler for $g \geq 3$ and it seems unknown whether finite index subgroups of M_g have property FA_1 . Bridson proved in [2] that for any normal subgroup H of index n in M_g , for $g \geq 3$, and any homomorphism $\phi : H \rightarrow G$ to a group G acting by hyperbolic isometries on some complete CAT(0) space – in particular, to $G = \mathbb{Z}$ – the n -th powers of Dehn twists (which belong to H) lie in the kernel of ϕ . Such homomorphisms ϕ have therefore striking similarities with the quantum representations.

Corollary 1.1 follows from Proposition 3.1 above and Bridson’s result from [3] saying that M_g has strong FA_g .

Acknowledgements. We are indebted to Jean-Benoît Bost, Benson Farb, Eric Gaudron and Bertrand Remy for helpful discussions and to the referees for pointing out an incomplete argument in the first version. The author was partially supported by the ANR Reperf: ANR-06-BLAN-0311.

REFERENCES

- [1] J.E.Andersen, *Asymptotic faithfulness of the quantum $SU(n)$ representations of the mapping class groups*, Ann. of Math. (2) 163(2006), 347–368.
- [2] M.R.Bridson, *Semisimple actions of mapping class groups on $CAT(0)$ spaces*, arXiv:0908.0685, "The Geometry of Riemann Surfaces", London Mathematical Society Lecture Notes 368, Dedicated to Bill Harvey on his 65th birthday, to appear.
- [3] M.R.Bridson, *On the dimension of $CAT(0)$ spaces where mapping class groups act*, arXiv:0908.0690.
- [4] C.Chevalley, *Théorie des groupes de Lie II. Groupes algébriques*, *Actualités Sci. Ind.* 1152, Hermann & Cie., Paris, 1951.
- [5] M.Culler and K.Vogtmann, *A group-theoretic criterion for property FA*, Proc. Amer. Math. Soc. 124(1996), 677–683.
- [6] P.Deligne, *Extensions centrales non résiduellement finies de groupes arithmétiques*, C. R. Acad. Sci. Paris Sér. A-B 287(1978), no. 4, A203–A208.
- [7] B.Farb, *Some problems on mapping class groups and moduli space*, in Problems on mapping class groups and related topics, 11–55, Proc. Sympos. Pure Math., 74, Amer. Math. Soc., Providence, RI, 2006.
- [8] B.Farb and D.Margalit, *A primer of mapping class groups*, preprint.
- [9] B.Farb, *Group actions and Helly’s theorem*. Advances Math. 222(2009), 1574–1588.
- [10] B.Farb, A.Lubotzky and Y.Minsky, *Rank one phenomena in mapping class groups*, Duke Math. J. 106(2001), 581–597.
- [11] M.H.Freedman, K.Walker and Zhenghan Wang, *Quantum $SU(2)$ faithfully detects mapping class groups modulo center*, Geom. Topol. 6(2002), 523–539.
- [12] L.Funar, *Représentations du groupe symplectique et variétés de dimension 3*, C. R. Acad. Sci. Paris Sér. I Math. 316(1993), no. 10, 1067–1072.
- [13] L. Funar, *On the TQFT representations of the mapping class groups*, Pacific J. Math. 188(1999), 251–274.
- [14] L.Funar, *Some abelian invariants of 3-manifolds*, Rev. Roumaine Math. Pures Appl. 45(2000), 825–861.
- [15] S.Gervais, *Presentation and central extensions of mapping class groups*, Trans. Amer. Math. Soc. 348(1996), 3097–3132.

- [16] M. Korkmaz and A.I. Stipsicz, *The second homology groups of mapping class groups of oriented surfaces*, Math. Proc. Cambridge Philos. Soc. 134(2003), 479–489.
- [17] G.Margulis, Discrete subgroups of semisimple Lie groups, *Ergebnisse Math. Grenz.* 17, Springer Verlag, Berlin, 1991.
- [18] G.Masbaum and J.D.Roberts, *On central extensions of mapping class groups*, Math. Ann. 302(1995), 131–150.
- [19] M.Reid, Undergraduate commutative algebra, *London Mathematical Society Student Texts* 29, Cambridge University Press, Cambridge, 1995.
- [20] A.Weil, Foundations of Algebraic Geometry, *American Mathematical Society Colloquium Publications* 29, American Mathematical Society, New York, 1946.

INSTITUT FOURIER BP 74, UMR 5582, UNIVERSITY OF GRENOBLE I, 38402 SAINT-MARTIN-D'HÈRES CEDEX, FRANCE
E-mail address: funar@fourier.ujf-grenoble.fr