

# Determinant of Sturm-Liouville operators and semi-classical trace formulae (IHP, June 2005)

Yves Colin de Verdière

July 4, 2005

Institut Fourier, Unité mixte de recherche CNRS-UJF 5582  
BP 74, 38402-Saint Martin d'Hères Cedex (France)  
yves.colin-de-verdiere@ujf-grenoble.fr  
<http://www-fourier.ujf-grenoble.fr/~ycolver/>

## Introduction

These lectures will be devoted to *Semi-Classical Trace Formulae* (SCTF). The aim of SCTF is to derive properties of the eigenvalues of a *Schrödinger operator*  $\hat{H}_h$  in the *semi-classical regime* where  $h$ , the Planck constant, is very small. From the mathematical point of view,  $h$  will tend to  $0^+$ . The formal idea is to compute the trace of a suitable function of  $\hat{H}_h$  in two ways

- $\text{Trace}(f(\hat{H}_h)) = \sum_j f(E_j(h))$  where  $E_j(h)$  are the eigenvalues of  $\hat{H}_h$
- $\text{Trace}(f(\hat{H}_h)) = \int_X [f(\hat{H}_h)](x, x) |dx|$  where  $[A](x, y)$  denotes the Schwartz kernel of the operator  $A$ .

Then comes the hard part which is to find suitable formulae or approximations to  $[f(\hat{H}_h)](x, y)$  using methods of partial differential equations.

For example, if  $\Delta_g$  is the Laplace operator on a compact Riemannian manifold, we start with  $\hat{H} = \frac{h^2}{2} \Delta_g$ . There are several possibilities for  $f$ :

- $f(E) = e^{-tE}$  which corresponds to *heat equation*
- $f(E) = e^{-it\sqrt{2E}/h} = e^{-it\sqrt{\Delta_g}}$  which corresponds to the *wave equation*

- $f(E) = e^{-itE/h}$  which corresponds to the *Schrödinger equation*.

The trace of heat kernel  $\sum \exp(-th^2\lambda_j/2)$  where  $\lambda_j$  are the eigenvalues of  $\Delta_g$  admits as  $h \rightarrow 0$  (or  $t \rightarrow 0$  with  $h = 1$ ) an asymptotic expansion whose coefficients are related to locally computable quantities like curvature and see nothing about the classical dynamics [BGM71].

Wave equations give the more natural results for the Laplace operator, but are working only for the Laplace operator. Moreover they make a strong use of Schwartz's distributions [Du-Gu75].

We will use Schrödinger equation. Our goal is to derive a rather elementary proof which is not using the full symbolic calculus of Fourier integral operators, but only the *metaplectic representation*.

We will start with a more simpler type of operator which is the quantization of a *twist map* and then see how to reduce the Schrödinger case to the previous case.

There will be three main parts:

1. We will first write a trace formula for a Fourier Integral Operators  $U$  associated with a twist canonical transformation: the Schwartz kernel of such an operator admits a WKB form and the asymptotics of the traces  $t_N := \text{Trace}(U^N)$  can be derived from the stationnary phase approximation. The main result is a rather explicit asymptotic expansion involving the fixed points of  $\chi^N$ .
2. From the previous formula, we will derive a trace formula for *Schrödinger operators* in the semi-classical limit and the expression of the smoothed density of states known as *Gutzwiller formula*. Gutzwiller formula expresses the smoothed density of states in terms of periodic orbits of the classical limit which is an Hamiltonian system. We will need the short time asymptotic of the propagator which goes back to Lax and Hörmander [La57, Hör68].
3. One of the main difficulties is the computation of the stationnary phase expansions. We will show how the calculus of determinants can be achieved using the metaplectic representation. In particular, we will give formulae for some *determinants of discrete Sturm-Liouville operators* (Jacobi type matrices) which goes back to a formula of Levit and Smilansky [Lev-Smil77] in the continous case (see also [K-T91]).

## 1 About the history

SCTF has several origines : on one side, Selberg's trace formula [Sel56] is an exact summation formula concerning the case of locally symmetric spaces; this formula was interpreted by H. Huber [Hub59] as a formula relating eigenvalues

of the Laplace operator and lengths of closed geodesics (also called the “lengths spectrum”) on a closed surface of curvature  $-1$ .

On the other side, around 1970, 2 groups of physicists developed independently asymptotic trace formulae:

- M. Gutzwiller [Gut71] for the Schrödinger operator, using the quasi-classical approximation of the Green function; it is interesting to note that the word “trace formula” is not written, but Gutzwiller instead speaks of a new “quantization method” (the old one being “EBK” or “Bohr-Sommerfeld rules”).
- R. Balian and C. Bloch [BB70, BB71, BB72] for the eigenfrequencies of a cavity use what they call “multiple reflection expansions”. They asked about a possible application to Kac’s problem.

At the same time, under the influence of Mark Kac’s famous paper [Kac66] “Can one hear the shape of a drum?”, mathematicians became quite interested into inverse spectral problems, mainly using heat kernel expansions (for the state of the art around 1970, see [BGM71]).

The SCTF was put into its final mathematical form for the Laplace operator on compact Riemannian manifolds without boundary by 3 groups:

- In my thesis [CdV73a, CdV73b], I used the short time expansion of the Schrödinger kernel and an approximate Feynman path integral. I proved that the spectrum of the Laplace operator determines generically the lengths of closed geodesics
- J. Chazarain [Cha74] derived the qualitative form of the trace for the wave kernel using FIO’s
- Using the full power of the symbolic calculus of FIO’s, H. Duistermaat and V. Guillemin [Du-Gu75] were able to compute the main term of the singularity from the Poincaré map of the closed orbit. Their paper became, at least for mathematicians, the canonical reference on the subject.

After that, people were able to extend SCTF to:

- General semi-classical Hamiltonians [Br-Ur91, Mei92]
- Manifolds with boundary [Gu-Me79]
- Surfaces with conical singularities and polygonal billiards [Bo-Pa-Sc00, Hill02]
- Several operators commuting operators [Cha-Po99]

Recently, people [Gu96, Zel97, Zel98] became able to say something about the non principal terms in the singularities expansion which come from the semi-classical Birkhoff normal form.

## 2 The trace formula for a quantized twist map

### 2.1 Twist maps

**Definition 1** A twist map is a canonical diffeomorphism  $\chi : V \rightarrow W$  where  $V$  and  $W$  are open sets of the cotangent bundle  $T^*X$  of a smooth  $d$ -manifold  $X$ , which satisfies: the projection  $\pi : \Gamma_\chi \rightarrow X \times X$  where  $\Gamma_\chi$  is the graph of  $\chi$  is a diffeomorphism of  $\Gamma_\chi$  onto an open subset  $\Omega$  of  $X \times X$ .

The twist map  $\chi$  will be called exact if the closed form  $\alpha_\chi$ , which is the restriction of the Liouville 1-form  $\xi dx - \eta dy$  of  $T^*(X \times X)$  to  $\Gamma_\chi$  is exact.

We will call  $S : \Omega \rightarrow \mathbb{R}$  any function which satisfies  $dS = \alpha_\chi$ . Such a function  $S$  is called a generating function of  $\chi$ .

A locally twist map is a canonical diffeomorphism  $\chi : V \rightarrow W$  which is locally a twist map.

If  $\chi$  is a twist map with generating function  $S$ , we have

$$\chi\left(y, -\frac{\partial S}{\partial y}\right) = \left(x, \frac{\partial S}{\partial x}\right).$$

Also, if  $A := \chi'(z_0)$  is the derivative of  $\chi$  at the point  $z_0$  and if  $\chi$  is twist at  $z_0$ ,  $A$  is a twist linear map whose generating function is the Hessian of the generating function  $S$  of  $\chi$  at the point  $(x_0, y_0)$  corresponding to  $z_0$ .

#### Example 2.1 (Twist maps of the annulus)

This is a much studied example since Poincaré and Birkhoff (see [Be94]). We take  $X = \mathbb{R}/\mathbb{Z}$ :

- The *Poincaré map*  $P$  of an elliptic periodic orbit of an Hamiltonian system with 2 degrees of freedom: the map  $P$  admits in the generic case a so-called Birkhoff normal form

$$P : (\theta, \rho) \rightarrow (\theta + F(\rho) + o(\rho^N), \rho + o(\rho^N))$$

and hence is a twist map of the annulus for  $\rho$  close to 0 if  $F'(0) \neq 0$ .

- The *Frenkel-Kontorova map*  $\chi(y, \eta) = (y + \alpha(y) + \eta, \alpha(y) + \eta)$  is a twist canonical map which is exact iff  $\alpha(y)$  is the derivative of a periodic function  $f(y)$ . In the latter case, we can take

$$S(x, y) = \frac{1}{2}(x - y)^2 + f(y).$$

- The *billiard map*: if  $X$  is a smooth convex billiard plane table, the return map,  $T(s, \sin \theta) = (s', \sin \theta')$  where  $s$  is the arc length and  $\theta$  the incidence angle on the boundary at the point  $m(s)$ , contains the main part of the dynamics and is a twist map [K-T91]. It admits the distance  $S(s, s') = d(m(s), m(s'))$  as a generating function.

**Example 2.2 (Small time flow of a regular Lagrangian)** Let us start with a Lagrangian  $L(x, v) : TX \rightarrow \mathbb{R}$  and assume that the Legendre transform  $\text{Leg} : (x, v) \rightarrow (x, \frac{\partial L}{\partial v})$  is a global diffeomorphism from  $TX$  onto  $T^*X$ . Let  $\varphi_t$  be the associated Hamiltonian flow:  $\varphi_t : T^*X \rightarrow T^*X$ . If  $V \subset T^*X$  is an open *bounded* set, the flow  $\varphi_t$  restricted to  $V$  is a twist map for  $t \neq 0$  small enough.

The proof: *consider the expansion*

$$\varphi_t(y, \eta) = (y + t \frac{\partial H}{\partial \xi}(y, \eta) + O(t^2), \eta - t \frac{\partial H}{\partial x}(y, \eta) + O(t^2))$$

and apply inverse function Theorem using the fact that  $\frac{\partial^2 H}{\partial \xi^2}$  is invertible.

Let us consider the open set  $\Omega \subset X \times X$  as before, for any  $(x, y) \in \Omega$  there exists an unique extremal curve  $\gamma_{xy} : [0, t] \rightarrow X$  from  $y$  to  $x$  so that  $\text{Leg}(y, \dot{\gamma}(0)) \in V$ . The action integral  $S(x, y) := \int_0^t L(\gamma_{xy}(s), \dot{\gamma}_{xy}(s)) ds$  is a generating function for  $\varphi_t : V \rightarrow W$ .

A more specific example:

**Example 2.3**  $(X, g)$  is a Riemannian manifold,  $Q : X \rightarrow \mathbb{R}$ ,  $L = \frac{1}{2} \|v\|^2 - Q(x)$ ,  $H = \frac{1}{2} \|\xi\|^2 + Q(x)$  and  $Q_\infty = \liminf_{x \rightarrow \infty} Q(x)$ . We can take  $E < Q_\infty$  and  $V = \{H(y, \eta) < E\}$ .

If  $Q \equiv 0$ , we have  $S(x, y) = d^2(x, y)/2t$  where  $d$  is the Riemannian distance and we can take  $\Omega = \{(x, y) | d(x, y) < \rho\}$  where  $\rho$  is the injectivity radius of  $(X, g)$ . In this case, the map  $(y, \eta) \rightarrow x(\varphi_t(y, \eta))$  is the exponential map:  $x = \exp(y, t\eta)$ .  $\varphi_t$  is locally twist near  $(y, \eta)$  with  $x = \exp(y, t\eta)$  if  $x$  and  $y$  are not conjugate points along the geodesic  $\gamma(s) = \exp(y, st\eta)$ ,  $0 \leq s \leq 1$ .

## 2.2 Quantizations of a twist map

We will consider, for  $h > 0$ , an operator  $U_h : L^2(X) \rightarrow L^2(X)$  whose (Schwartz) kernel is

$$[U_h](x, y) = (2\pi i h)^{-d/2} a(x, y) s(x, y) e^{iS(x, y)/h} |dx dy|^{\frac{1}{2}}$$

with  $a \in C_o^\infty(\Omega)$  and  $s(x, y) = |\det(\partial_{x, y}^2 S)|^{\frac{1}{2}}$ . The fonction  $s$  is a normalization which makes  $U_h$  unitary in case  $a = 1$  and  $S$  is quadratic (see Section 4.2). The operator  $U$  is smoothing, with compactly supported kernel and is called a *quantization of  $\chi$* . It would be nice to know about its spectrum in the semi-classical limit!

It is interesting to note that in general  $U_h$  is not a *normal operator*. It would imply, by stationnary phase approximation, that, if  $\tilde{a}(y, \eta) = a(x, y)$  with  $(x, \xi) = \chi(y, \eta)$ , then  $|\tilde{a}|$  is constant along the trajectories of  $\chi$  which is impossible if they escape to infinity!

An important exception will be  $\psi(\hat{H}) \exp(-it\hat{H}/h)$  with  $\hat{H}$  a semi-classical Schrödinger operator and the support of  $\psi$  is contained in  $] -\infty, E_\infty[$  (see Section 3.1).

## 2.3 Discrete Feynman path integrals

One way to get some informations on the eigenvalues of  $U$  is to derive “trace formulae” expressing the asymptotic behaviour of  $t_N := \text{Trace}(U^N)$  as  $h \rightarrow 0$ .

We can formally write  $[U^N]$  and  $t_N$  as discrete Feynman integrals: a path  $\gamma$  of length  $N$  will be a sequence  $(y = x_0, \dots, x = x_N)$  with  $(x_k, x_{k+1}) \in \Omega$ . We will denote by  $P_{x,y}^N$  the set of such paths with the measure  $|d\gamma| = |dx_1 \cdots dx_{N-1}|$ . The action of the path  $\gamma = (y, \dots, x)$  will be  $S(\gamma) := S(x, x_{N-1}) + \cdots + S(x_1, y)$ . We have

$$[U^N](x, y) = (2\pi i h)^{-Nd/2} \int_{P_{x,y}^N} e^{\frac{i}{h} S(\gamma)} A(\gamma) |d\gamma|, \quad (1)$$

and

$$t_N = (2\pi i h)^{-Nd/2} \int_X |dx| \left( \int_{P_{x,x}^N} e^{\frac{i}{h} S(\gamma)} A(\gamma) |d\gamma| \right). \quad (2)$$

Using Fubini and defining  $P^N$  as the set of periodic sequences  $(x_0, x_1, \dots, x_{N-1}, x_0)$ , so that  $\forall i, (x_i, x_{i+1}) \in \Omega$ , we get

$$t_N = (2\pi i h)^{-Nd/2} \int_{P^N} e^{\frac{i}{h} S(\gamma)} A(\gamma) |d\gamma|.$$

We will evaluate the previous expressions by the stationary phase.

**Lemma 1** • *The critical points of  $S(\gamma) : P_{x,y}^N \rightarrow \mathbb{R}$  are sequences  $(y, x_1, \dots, x_{N-1}, x)$  which are the projections of orbits  $\chi^k(y, \eta)$ ,  $k = 0, \dots, N-1$ :  $\chi^k(y, \eta) = (x_k, \xi_k)$*

- *The critical points of  $S(x_0, x_1, \dots, x_{N-1}, x_0)$  are the projections of closed orbits sitting in  $V$ .*

*Proof.*–

Let us define  $\xi_0 = -\partial_2 S(x_1, y)$  and for  $k = 1, \dots, N$ ,  $\xi_k = \partial_1 S(x_k, x_{k-1})$ . The criticality condition is, for all  $k = 1, \dots, N-1$ ,  $\partial_2 S(x_{k+1}, x_k) = -\partial_1 S(x_k, x_{k-1})$  which is equivalent to say that, for  $k = 1, \dots, N-1$ , we have  $(x_{k+1}, \xi_{k+1}) = \chi(x_k, \xi_k)$ .

The same argument extends to the cyclic case.

□

## 2.4 Non degeneracy

There are several possible non degeneracy assumptions. They can be formulated “à la Morse-Bott” (critical point of action integrals) or purely symplectically.

**Definition 2** Two submanifolds  $Y$  and  $Z$  of  $X$  intersect cleanly iff  $Y \cap Z$  is a manifold whose tangent space is the intersection of the tangent spaces of  $Y$  and  $Z$ .

A diffeomorphism  $F$  admits a clean manifold of fixed points if the graph of  $F$  intersects cleanly the diagonal.

A manifold  $Z$  of critical points of a function  $f$  is Morse-Bott ND if the Hessian of  $f$  is ND transversally to  $Z$ .

We will need the

**Lemma 2** Using the previous notations,  $\chi^N$  admits a clean manifold  $W$  of fixed points if and only if the action  $S(x_1, x_2, \dots, x_{N-1}, x_1)$  admits a Morse-Bott ND critical manifold.

The following statement will be explained in Section 4.4:

**Lemma 3 (Canonical measures)** If  $W$  is a clean manifold of fixed points of a symplectic map,  $W$  is equipped with a canonical smooth measure  $d\mu_W$ .

## 2.5 Van Vleck's formula

**Theorem 1** Let us give  $x, y \in X$  so that  $\chi^N$  is a locally twist map near each  $(y, \eta)$  so that  $\chi^N(y, \eta) = (x, \cdot)$  (which is true for almost all pairs  $(x, y)$  thanks to Sard's Theorem), we have

$$[U^N](x, y) \sim (2\pi i h)^{-d/2} \left( \sum_{\alpha} |\det \partial_{xy}^2 S_{\alpha}|^{\frac{1}{2}} e^{-i\mu_{\alpha}\pi/2} e^{iS_{\alpha}/h} \sum_{j=0}^{\infty} B_j^{\alpha} h^j \right)$$

where

- $\alpha$  labels the solutions of  $\chi^N(y, \eta_{\alpha}) = (x, \cdot)$ . We will denote by  $\gamma_{\alpha} = (x_0^{\alpha} = y, x_1^{\alpha}, \dots, x_N = x)$  the projection of the trajectories:  $\chi^k(y, \eta_{\alpha}) = (x_k^{\alpha}, \xi_k^{\alpha})$
- $\mu_{\alpha}$  is the Morse index<sup>1</sup> of  $S(\gamma) : P_{xy}^N \rightarrow \mathbb{R}$  for the critical point  $\gamma_{\alpha}$ .
- $S_{\alpha}(x, y) = \sum_{k=0}^{N-1} S(x_{k+1}^{\alpha}, x_k^{\alpha})$ , with  $x_N^{\alpha} = x$ ,  $x_0^{\alpha} = y$  (a local generating function for  $\chi^N$ )
- $B_0^{\alpha} = \prod_{k=0}^{N-1} a(x_{k+1}^{\alpha}, x_k^{\alpha})$ .

*Proof.*–

This formula is a consequence of stationary phase (Theorem 9) and of the formula for the determinant of a Jacobi matrix (Theorem 11). □

---

<sup>1</sup>The Morse index of a critical point  $x$  of a smooth function  $F : X \rightarrow \mathbb{R}$  is the maximal dimension of a subspace of the tangent space  $T_x X$  on which the hessian of  $F$  is  $< 0$

## 2.6 Trace formulae

We get:

**Theorem 2** *Let us assume that  $\chi$  admits (a finite number of) clean connected manifolds  $W_\alpha$  of periodic points, i.e. points satisfying  $\chi^N(z) = z$ . We get the following asymptotic expansion of the trace  $t_N$ :*

$$t_N := \text{Trace}(U^N) \sim \sum_{\alpha} (2\pi i h)^{-\nu_\alpha/2} e^{-i\iota_\alpha \pi/2} e^{iS(W_\alpha)/h} \left( \sum_{j=0}^{\infty} A_j^\alpha h^j \right),$$

with

•

$$A_0^\alpha = \int_{W_\alpha} \prod_{j=1}^N a(x(\chi^j(z)), x(\chi^{j-1}(z))) d\mu_\alpha,$$

with  $d\mu_\alpha$  is the canonical measure on  $W_\alpha$

•  $\nu_\alpha = \dim W_\alpha$

•  $\iota_\alpha$  is the Morse index of the action  $S(\gamma)$  on the periodic path  $\gamma_\alpha = (x_0, x_1, \dots, x_N = x_0)$

*Proof.*–

The proof is by application of the stationary phase for ND critical manifolds (Section 4.1) and computation of the canonical measure (Section 4.5).

□

In general, it is difficult to extract precise informations on the asymptotic behaviour of the eigenvalues of  $U_h$  from the trace formula. We will now see that the situation is much better in case of a flow  $U(t)$ .

## 3 Trace formula for Schrödinger operators

The purpose of this section is to describe the so-called “semi-classical trace formula” (SCTF) relating the *spectrum* of a semi-classical Hamiltonian to the *periods of closed orbits* of its classical limit. We will mainly describe the case of the Schrödinger operator on a Riemannian manifold which contains the purely Riemannian case.

## 3.1 Schrödinger operators on Riemannian manifolds

### 3.1.1 The Laplace operator

Let  $(X, g)$  be a smooth compact connected Riemannian manifold of dimension  $d$ . Let us denote by  $|dx| = |g| |dx_1 \cdots dx_d|$ , with  $|g| = \sqrt{\det(g_{ij})}$ , the canonical volume element on  $(X, g)$  and  $L^2(X, |dx|)$  the associated Hilbert space. There exists a canonical symmetric differential operator of order 2, the *Laplace operator* on  $(X, g)$ , denoted  $\Delta_g$ , which is given in local coordinates by:

$$\Delta_g = -|g|^{-1} \partial_i |g| g^{ij} \partial_j .$$

Beause  $X$  is compact,  $\Delta_g$  admits an unique self-adjoint extension and  $L^2(X, |dx|_g)$  admits an orthonormal basis of (smooth) eigenfunctions. It will be convenient to label such a basis by integers  $(\varphi_j, j = 1, \dots)$  so that  $\Delta_g \varphi_j = \lambda_j \varphi_j$  and

$$\lambda_1 = 0 < \lambda_2 \leq \lambda_3 \leq \dots \leq \lambda_j \leq \dots$$

with  $\lambda_j \rightarrow \infty$ . Note that the sequence  $(\lambda_j)$  is uniquely defined, but the  $\varphi_j$  are not unique, only the eigenspaces are.

With the exception of a very few cases (compact rank one symmetric spaces, tori) the eigenvalues are not explicitly computable.

It is known (Weyl's law) that  $\lambda_j \sim c_d (\text{vol}(X))^{-2/d} j^{2/d}$  with  $c_d > 0$  an universal constant.

The goal of the SCTF is the description in terms of the geodesic flow of some asymptotic properties as  $j \rightarrow \infty$  (the classical limit) of the eigenvalues of  $\Delta_g$ .

### 3.1.2 Semi-classical Schrödinger operators

If  $(X, g)$  is a (possibly non compact) Riemannian manifold and  $Q : X \rightarrow \mathbb{R}$  a smooth function which satisfies  $\liminf_{x \rightarrow \infty} Q(x) = E_\infty > -\infty$ , the differential operator  $\hat{H} = \frac{1}{2} h^2 \Delta + Q$  is semi-bounded from below and admits self-adjoint extensions. For all those extensions the spectrum is discrete in the intervall  $] - \infty, E_\infty[$  and eigenfunctions  $\hat{H} \varphi_j = E_j \varphi_j$  are localized in the domain  $Q \leq E_j$ . If  $X$  is compact and  $Q = 0$ , we recover the case of the Laplace operator.

We will denote by

$$\inf Q < E_1(h) < E_2(h) \leq \dots \leq E_j(h) \leq \dots < E_\infty$$

this part of the spectrum.

The SCTF can also be derived the same way for Schrödinger operators with magnetic field ... One can even extend it to Hamiltonian systems which are not obtained by Legendre transform from a regular Lagrangian. In this case, Morse indices have to be replaced by the more general Maslov indices.

## 3.2 Classical dynamics

### 3.2.1 Geodesics

The geodesic flow  $G_t$  of  $(X, g)$  is defined as the following 1-parameter group of diffeomorphisms on the tangent bundle  $TX$ :  $G_t(x_0, v_0) = (x_t, v_t)$  where  $t \rightarrow x_t$  is the geodesic of  $X$  so that  $x(t=0) = x_0$  and  $\dot{x}(t=0) = v_0$  and  $v_t$  is the speed at time  $t$  of the previous geodesic. It is well known that the geodesic flow can also be described using the Hamiltonian formulation. On  $T^*X$ , one considers the Hamiltonian  $H_0(x, \xi) = \frac{1}{2}\|\xi\|_g^2$  and will still denote by  $G_t$  the geodesic flow on  $T^*X$ .

### 3.2.2 Newton flows

Euler-Lagrange equations for the Lagrangian  $L(x, v) := \frac{1}{2}\|v\|_g^2 - Q(x)$  admit a Hamiltonian formulation on  $T^*X$  whose energy is given by  $H = \frac{1}{2}\|\xi\|_g^2 + Q(x)$ . We will denote by  $X_H$  the Hamiltonian vector field

$$X_H := \sum_j \frac{\partial H}{\partial \xi_j} \partial_{x_j} - \frac{\partial H}{\partial x_j} \partial_{\xi_j} .$$

Preservation of  $H$  by the dynamics shows immediately that the Hamiltonian flow  $\Phi_t$  restricted to  $H < E_\infty$  is complete.

### 3.2.3 Closed orbits

**Definition 3** A closed orbit  $(\gamma, T)$  of the Hamiltonian  $H$  consists of an orbit  $\gamma$  of  $X_H$  which is homeomorphic to a circle and a nonzero real number  $T$  so that  $\Phi_T(z) = z$  for all  $z \in \gamma$ .  $T$  will be called the period of  $\gamma$ .

We will denote by  $T_0(\gamma) > 0$  (the primitive period) the smallest  $T > 0$  for which  $\Phi_T(z) = z$ .

- The (linear) Poincaré map  $\Pi_\gamma$  of a closed orbit  $(\gamma, T)$  with  $H(\gamma) = E$ : we restrict the flow to  $S_E := \{H = E\}$  and take an hypersurface  $\Sigma$  inside  $S_E$  transversal to  $\gamma$  at the point  $z_0$ . The associated return map  $P_\gamma$  is a local diffeomorphism fixing  $z_0$ . Its linearization  $\Pi_\gamma := P'_\gamma(z_0)$  is the linear Poincaré map, an invertible (symplectic) endomorphism of the tangent space  $T_{z_0}\Sigma$ .
- The Morse index  $\iota(\gamma)$ : closed orbits  $(\gamma, T)$  are critical points of the action integral  $\int_0^T L(\gamma(s), \dot{\gamma}(s)) ds$  on the manifold  $C^\infty(\mathbb{R}/T\mathbb{Z}, X)$ . They have always a finite Morse index (see [Mil67]) which is denoted by  $\iota(\gamma)$ . For general Hamiltonian systems, the Morse index is replaced by the Conley-Zehnder index [Mei94].

- The *nullity index*  $\nu(\gamma)$  is the dimension of the space of infinitesimal deformations of the closed orbit  $\gamma$  by closed orbits. It is the dimension of the kernel of the map  $(\delta t, \delta z) \rightarrow \Phi'_T(z_0)\delta z - \delta z + \delta t X_H$ . We have always  $\nu(\gamma) \geq 2$ .

**Example 3.1 (Geodesic flows)**

- Manifold with  $< 0$  sectional curvature: *in this case, we have for all closed geodesics*  $\iota(\gamma) = 0$ ,  $\nu(\gamma) = 2$
- Flat tori of dimension  $d$ : *we have then*  $\iota(\gamma) = 0$  and  $\nu(\gamma) = d + 1$ .
- Sphere of dimension 2 with constant curvature: *if  $\gamma_n$  is the  $n$ th iterate of the great circle, we have*  $\iota(\gamma_n) = 2|n|$  and  $\nu(\gamma_n) = 4$ .

It is a beautiful result of J.-P. Serre [Se50] that any pair of points  $(a, b)$  on a closed Riemannian manifold are end points of infinitely many geometrically distinct geodesics. Counting geometrically distinct closed geodesics is much harder especially for simple manifold like the spheres. It is now known that every closed Riemannian manifold admits infinitely many geometrically distinct closed geodesics (at least in some cases for a generic metric, [Be00] Chap. V). Concerning more general Hamiltonian systems a lot is known [Ho-Ze94].

**3.2.4 Non degeneracy and orbits cylinders**

**Definition 4** A closed orbit  $\gamma$  will called weakly non degenerate (WND) if 1 is not an eigenvalue of the Poincaré map  $\Pi_\gamma$ .

**Remark 1** If  $\gamma$  is WND, it does not imply that the iterates are WND, because some non trivial roots of unity could be eigenvalues of  $\Pi_\gamma$ .

**Lemma 4** If  $\gamma_{E_0}$  is a WND orbit with  $H(\gamma) = E_0$ , there exists for  $E$  close to  $E_0$  a closed orbit  $\gamma_E$  contained in  $H = E$  smoothly depending on  $E$ .

**Definition 5** The smooth family  $\gamma_E$  with  $E$  close to  $E_0$  will be called an orbit cylinder.

For an orbit cylinder  $\gamma_E$  we will denote by  $T(E)$  the period of  $\gamma_E$  which is a smooth function of  $E$ .

**Definition 6** A closed orbit  $\gamma$  will called strongly non degenerate (SND) if  $\gamma$  is weakly non degenerate and  $T'(E_0) \neq 0$ .

**Example 3.2** In the case of the Riemannian geodesics, both non degeneracy assumptions coincide. They are true if, for example, the metric has  $< 0$  curvature. If  $X$  is fixed, all closed geodesics are (SND) for a generic (in the Baire sense) Riemannian metric.

More general ND assumptions can be introduced in order to cover for example the case of completely integrable systems. In this case they recover the usual ND assumptions of KAM theories.

**Proposition 1** *If  $\gamma$  is a WND orbit of period  $T$ , there exists an (unique) symplectic splitting of the tangent space to any point  $z$  of  $\gamma$  so that the differential of the flow at time  $T$  is given by:*

$$\Phi'_T(z) = \begin{pmatrix} 1 & -dT/dE & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \Pi_\gamma \end{pmatrix} .$$

*The first piece of the splitting is the 2d tangent space to the orbit cylinder.*

### 3.2.5 Actions

**Definition 7** *If  $(\gamma, T)$  is a periodic orbit, we define the following quantities which are both called action of  $\gamma$ :*

$$A(\gamma) = \int_\gamma \xi dx, \quad B(\gamma) = \int_\gamma (\xi dx - H dt) = \int_0^T L(\gamma(t), \dot{\gamma}(t)) dt .$$

From the definition of the Legendre transform, we get:  $A(\gamma) - B(\gamma) = H(\gamma)T$ .

If  $\gamma$  is SND, we have 2 smooth functions  $A = A(E)$  and  $B = B(T)$  associated to any orbit cylinder. They satisfies

$$\frac{dA}{dE} = T(E), \quad \frac{dB}{dT} = -E(T)$$

We can summary the previous properties as follows:

**Theorem 3**  *$A(E)$  and  $-B(T)$  are Legendre transforms of each other in the symplectic space  $(\mathbb{R}^2, dE \wedge dT)$ .*

## 3.3 Playing with spectral densities: the pre-trace formula

We will define “regularized spectral densities”. The general idea is as follows: we want to study a sequence of real numbers  $E_j(h)$  (a spectrum) in some intervall  $[a, b]$  depending of a small parameter  $h$ . We introduce a non negative function  $\rho \in \mathcal{S}(\mathbb{R})$  which satisfy  $\int \rho(t) dt = 1$  and introduce

$$D_{\rho, \varepsilon, h}(E) = \sum \frac{1}{\varepsilon} \rho \left( \frac{E - E_j(h)}{\varepsilon} \right) .$$

It gives the analysis of the spectrum at the scale  $\varepsilon$ . Of course, we will adapt the scaling  $\varepsilon$  to the small parameter  $h$ . If the scaling is smaller than the size of the mean spacing of the spectrum, we will get a very precise resolution of the spectrum.

The general philosophy is:

- If  $h$  is the semi-classical parameter of a semi-classical Hamiltonian, the mean spacing  $\overline{\Delta E}$  is of order  $h^d$  (Weyl's law). The trace formula gives the asymptotic behaviour of  $D_{\rho,\varepsilon,h}(E)$  for  $\varepsilon \sim h$  (and hence  $\varepsilon \gg \overline{\Delta E}$  except if  $d = 1$ ). This behaviour is not universal and thus contains a lot of informations (in our case, on closed trajectories)
- Better resolution of the spectrum needs the use of the long time behaviour of the classical dynamics and is conjecturally “universal”

From now on, we choose 2 smooth functions:

- $\rho(\tau)$  which is equal to 1 near 0 and so that  $\hat{\rho}(t) = \int e^{-i\tau t} \rho(\tau) d\tau$  is compactly supported
- $\psi \in C_0^\infty(]-\infty, E_\infty[)$ .

We define

- The *regularized density of states*

$$D(E) := \sum_{j=1}^{\infty} \psi(E_j) \frac{1}{h} \rho\left(\frac{E - E_j(h)}{h}\right).$$

If  $I = [a, b] \subset ]-\infty, E_\infty[$ ,  $a < E < b$  and  $\psi \equiv 1$  near  $E$ , we have

$$D(E) = \sum_{a < E_j < b} \frac{1}{h} \rho\left(\frac{E - E_j(h)}{h}\right) + O(h^\infty).$$

- The *partition function* given by the (finite) sum:

$$Z(t) = \sum_{j=1}^{\infty} \psi(E_j) e^{-itE_j/h} = \text{Trace}(U(t)) \quad (3)$$

with  $U(t) = \psi(\hat{H}) \exp(-it\hat{H}/h)$ .

Duistermaat-Guillemin's trick relates the behaviour of  $D(E)$  to the behaviour of  $Z(t)$  thanks to the fundamental *pre-trace formula*:

$$D(E) = \frac{1}{2\pi h} \int_{\mathbb{R}} e^{itE/h} \hat{\rho}(t) Z(t) dt. \quad (4)$$

The idea is then to start from a semi-classical approximation of the propagator  $[U(t)](x, y)$  and to insert  $Z(t) = \int_X [U(t)](x, x) |dx|$  into Equation (4).

### 3.4 The Schrödinger trace

Our goal is now to describe the asymptotic behaviour of  $D(E)$  around  $E_0$ . The main results can be expressed as follows:

**Theorem 4** *Let us assume that  $E_0 < E_\infty$  is not a critical value of  $H$  and all periodic trajectories contained in  $H = E_0$  and of periods in the support of  $\hat{\rho}$  are WND, we have, for  $E$  close enough to  $E_0$ :*

- $D(E) = D_{\text{Weyl}}(E) + \sum_{\gamma} D_{\gamma}(E) + O(h^\infty)$  where  $\gamma$  is one of the (finitely many) periodic trajectories in  $H = E$  and periods in  $\text{Support}(\hat{\rho})$

- 

$$D_{\text{Weyl}}(E) = (2\pi h)^{-d} \left( \sum_{j=0}^{\infty} a_j(E) h^j \right), \quad (5)$$

with  $a_0(E) = \int_{H=E} |dx d\xi / dH|$ .

- $D_{\gamma}(E)$  is a WKB function whose principal part is, if  $\gamma$  is SND:

$$\frac{\varepsilon}{2\pi h} e^{-i u(\gamma)\pi/2} e^{i \int_{\gamma} \xi dx / h} a_{\gamma}(E) \quad (6)$$

with  $a_{\gamma}(E) = \rho(T(E)) T_0(E) / |\det(\text{Id} - \Pi_{\gamma})|^{\frac{1}{2}}$ ,  $T_0(E)$  the primitive period, and

$$\varepsilon = \begin{cases} -i & \text{if } T'(E) > 0 \\ 1 & \text{if } T'(E) < 0 \end{cases}$$

If  $\gamma$  is only WND, the formula is the same except that  $\varepsilon e^{-i u(\gamma)\pi/2}$  is now (the exponential of) a Maslov index.

### 3.5 The Weyl term

Choose  $a < E_\infty$  and let  $t_0(a) > 0$  be the smallest period of a non trivial closed trajectory  $\gamma$  with  $H(\gamma) \leq a$  (see [Yo69]). In this section, we fix  $\rho$  so that  $\text{Support}(\hat{\rho})$  is contained in  $] -t_0, t_0[$ . If  $E$  is not a critical value of  $H$ , and, for  $\psi \in C_0^\infty(]-\infty, a[)$ , we have, insisting on the dependence w.r. to  $\psi$ :

$$D_{\psi}(E) = D_{\text{Weyl}}(E) \sim (2\pi h)^{-d} \left( \sum_{j=0}^{\infty} a_j(E, \psi) h^j \right),$$

with  $a_0(E, \psi) = \int_{H=E} \psi(H) |dL/dH|$ .

If we define the distributions

$$A_j(\psi) = \int_{-\infty}^{+\infty} a_j(E, \psi) dE$$

we get:

$$\sum \psi(E_j) \sim (2\pi h)^{-d} \left( \sum_{j=0}^{\infty} A_j(\psi) h^j \right) ,$$

Moreover, the previous asymptotics works even if they are critical values of  $H$  inside the support of  $\psi$ . The  $A_j$ 's are, for  $j \geq 1$  of the following form:

$$A_j(\psi) = \int_{T^*X} \sum_{l \geq 2}^{N_j} \psi^{(l)}(H(x, \xi)) P_{j,l}(x, \xi) |dx d\xi|$$

and the  $P_{j,l}(x, \xi)$ 's are ‘‘locally computable’’ from the Hamiltonian  $H(x, \xi)$  and its derivatives [CdV05].

**Remark 2** *If  $H(x, -\xi) = H(x, \xi)$ , we have the ‘‘transmission property’’ and the  $A_{2j+1}$  vanish.*

In the case of the Laplace operator and  $\chi(E) = e^{-E}$ , the  $A_j$ 's are called the ‘‘heat invariants’’ [BGM71].

From that kind of formula, it is possible to deduce the following estimates on the remainder term in Weyl's law :

**Theorem 5** *If  $a, b$  are not critical values of  $H$ :*

$$\#\{j | a \leq E_j(h) \leq b\} = (2\pi h)^{-d} \text{volume}(\{a \leq H \leq b\}) (1 + O(h)) .$$

This remainder estimate is optimal and was first shown in rather great generality by Hörmander [Hör68].

### 3.6 The contributions of periodic orbits

The proof involves the study, for  $t \neq 0$ , of  $Z(t) = \text{Trace}(U(t))$  which is given by:

**Proposition 2** • *If  $t$  is not the period of a trajectory  $\gamma$  with  $H(\gamma) \subset \text{Support}(\psi)$ ,  $Z(t) = O(h^\infty)$*

- *If  $\gamma$  is an SND closed trajectory of period  $T$ ,  $Z_\gamma(t)$  is near  $T$  a WKB function whose principal symbol is:*

$$\frac{1}{(2\pi i h)^{1/2}} e^{-i u(\gamma) \pi / 2} e^{i B(T) / h} b_0(T) , \tag{7}$$

$$\text{with } b_0(T) = T_0 \left| \frac{dE}{dT} \right|^{1/2} |\det(\text{Id} - \Pi_\gamma)|^{-1/2} .$$

It is then enough to put the previous asymptotic formula in Equation (4) and to apply once more a stationary phase argument.

## 3.7 Formal derivation from the Feynman path integral

### 3.7.1 The Feynman integral

R. Feynman ([Fe-Hi65, Mo51]) found a *geometric* representation of the propagator, i.e. the kernel  $p(t, x, y)$ , with  $t \neq 0$ , of the unitary group  $\exp(-it\hat{H}/\hbar)$  using an *integral* (FPI:= Feynman path integral) on the manifold  $\Omega_{t,x,y}$  of paths from  $y$  to  $x$  in the time  $t$ ; if  $L(\gamma, \dot{\gamma})$  is the Lagrangien, we have, for  $t > 0$ :

$$p(t, x, y) = \int_{\Omega_{t,x,y}} e^{\frac{i}{\hbar} \int_0^t L(\gamma(s), \dot{\gamma}(s)) ds} |d\gamma| ,$$

where  $|d\gamma|$  is a “Riemannian measure” on the manifold  $\Omega_{t,x,y}$  with the natural Riemannian structure.

There is no justification FPI as a flexible tool, but as we will see FPI is a very efficient tool in semi-classics.

### 3.7.2 The trace and loop manifolds

Let us try a formal calculation of the partition function and its semi-classical limit. We get:

$$Z(t) = \int_X |dx| \int_{\Omega_{x,x,t}} e^{\frac{i}{\hbar} \int_0^t L(\gamma(s), \dot{\gamma}(s)) ds} |d\gamma| .$$

If we denote by  $\Omega_t$  the manifold of paths  $\gamma : \mathbb{R}/t\mathbb{Z} \rightarrow X$ , (loops) and we apply Fubini (sic !), we get:

$$Z(t) = \int_{\Omega_t} e^{\frac{i}{\hbar} \int_0^t L(\gamma(s), \dot{\gamma}(s)) ds} |d\gamma| .$$

### 3.7.3 The semi-classical limit

We want to apply stationary phase in order to get the asymptotic expansion of  $Z(t)$ ; critical points  $J_t : \Omega_t \rightarrow \mathbb{R}$  are the closed orbits of the Euler-Lagrange flow and hence of the Hamiltonian flow of period  $t$ . We need

1. *The ND assumption (Morse-Bott)*
2. *The Morse index*
3. *The determinant of the Hessian*

1. The ND assumption is the original Morse-Bott one in Morse theory: we have smooth manifolds of critical points and the Hessian is transversally ND.

2. The Morse index is the Morse index of the action fonctionnal on closed loops:  $A(\gamma) := \int_0^t L(\gamma(s), \dot{\gamma}(s)) ds$ .

3. The Hessian is associated to a periodic Sturm-Liouville operator for which many regularizations have already been proposed (Levit-Smilanski [Lev-Smil77] and CdV [CdV99], Zeta regularization, see [R-S] ).

We get that way a sum of contributions given by the components  $W_{j,t}$  of  $W_t$ :

$$Z_j(t) = (2\pi i h)^{-1/2} e^{\frac{i}{h} B(\gamma_j)} A_j(h)$$

with  $A_j(h) \sim \sum_{l=0}^{\infty} a_{j,l} h^l$  and

$$a_{j,0} = \frac{e^{-i\mu\frac{\pi}{2}}}{|\delta|^{\frac{1}{2}}}$$

where  $\mu$  is the Morse index and  $\delta$  is a regularized determinant.

### 3.8 Sketch of the proof of SCTF

The proof starts with an approximation of the propagator  $[U(t)](x, y)$ . This approximation splits into 3 parts:

- **Near  $t = 0$ ,** it is known (see [La57]) that, if  $f(x) = a(x)e^{iS_0(x)/h}$  is a rapidly oscillating function with  $a \in C_0^\infty(X)$ , the solution of the Cauchy problem

$$\begin{cases} \frac{h}{i} \partial_t u + \hat{H}u &= 0 \\ u(t=0) &= f \end{cases}$$

is given, for  $t$  small enough by:

$$u(x, t) = a_h(x, t) e^{iS(x,t)/h}$$

with

- $a_h(x, t) \equiv \sum_{j=0}^{\infty} a_j(x, t) h^j$
- $S(x, t)$  satisfies the Hamilton-jacobi equation  $\partial_t S + H(x, \partial_x S) = 0$
- $a_h(x, 0) = a(x)$ ,  $S(x, 0) = S_0(x)$

Using  $S_0(x) = \langle x | \xi \rangle$  and the Fourier transform we get

$$[U(t)](x, y) \equiv (2\pi h)^{-d} \int e^{iS(t,x,y,\xi)/h} \left( \sum_{j=0}^{\infty} A_j(x, \xi, t) h^j \right) |d\xi| \quad (8)$$

with

- $S(t, x, y, \xi) = \langle x - y | \xi \rangle - tH(x, \xi) + O(t^2)$
- $A_0(x, \xi, 0) = \psi(H(x, \xi))$

- **For  $t \neq 0$  small enough.** If  $F < E_\infty$  and  $V := \{(y, \eta) | H(y, \eta) < F\}$ , then, for  $t \neq 0$  and  $|t| \leq t_F$  small enough, the Hamiltonian flow of  $H$  is a twist map. If  $\psi \in C_o^\infty(]-\infty, F[)$  and  $t \in [-t_F, t_F] \setminus 0$ , the propagator  $[U(t)] = [\exp(-it\hat{H}/h)\psi(H)]$  admits a WKB expansion of the form introduced in Section 2.2. It can be proved by applying the stationary phase to the expression (8).
- **For  $t$  large.** We have

$$\psi^N(\hat{H})e^{-it\hat{H}/h} = \left(\psi(\hat{H})e^{-i\frac{t}{N}\frac{\hat{H}}{h}}\right)^N$$

and, for any  $t$ , we will choose  $N$  so that  $t/N < t_F$ . We get then for  $[\psi^N(\hat{H})e^{-it\hat{H}/h}](x, y)$  an expression of the form given in Equation (1).

### 3.9 The wave trace

Let  $(X, g)$  be a compact Riemannian manifold and  $\lambda_1 = \mu_1^2 \leq \dots \leq \lambda_j = \mu_j^2 \leq \dots$  the eigenvalues of  $\Delta_g$ . The solution of the Cauchy problem for the wave equation  $\partial_t^2 u + \Delta_g u = 0$  is given in terms of the unitary group  $U^W(t) = \exp(-it\sqrt{\Delta})$  whose trace  $Z^W(t) = \sum_j e^{-it\mu_j}$  is a Schwartz distribution which is called the *wave trace*. We can rewrite in a semi-classical way  $U^W(t) = \exp(-it\hat{K}/h)$  with  $\hat{K} := h\sqrt{\Delta}$  and the classical Hamiltonian is then  $K(x, \xi) = \|\xi\|$  which is homogeneous of degree 1. The associated flow is the geodesic flow with speed  $\equiv 1$ . A closed orbit for  $K$ , even if it satisfies WND, is never SND, because periods of  $K$ -closed orbits are independant of energy!

One can pass from the Schrödinger SCTF to the wave trace quite formally because the eigenvalues  $F_j = h\mu_j$  of  $h\sqrt{\Delta}$  are related to the eigenvalues  $E_j = h^2\lambda_j/2$  of  $h^2\Delta/2$  by  $E_j = \frac{1}{2}(E'_j)^2$ .

The wave trace expansion can be derived from Theorem 4 as applied to  $h\sqrt{\Delta}$ . One can also pass directly from the trace formula for  $\hat{H}$  to a trace formula for  $\Phi(\hat{H})$  where  $\Phi$  is a diffeomorphism near the energy  $E$ .

**Theorem 6** *Let us assume that  $\hat{\rho}(t) \equiv 1$  near  $T_0$  and  $\psi(E) \equiv 1$  near  $E_0$ . The expansion corresponding to the periodic orbit  $\gamma$  (for both Hamiltonians  $H$  and  $\Phi(H)$ ) of new period  $T'_0 = T_0/\phi'(E_0)$  and new energy  $E'_0 = \Phi(E_0)$  is obtained just by the change of variables  $E' = \Phi(E)$  in the smoothed measures  $D_{\rho, \psi}(E)|dE|$ .*

From the Schrödinger trace  $D(\mu) := D_{\rho, \psi}(h, E)$ , with  $h = \mu^{-1}$ ,  $E = \frac{1}{2}$  and  $\psi \in C_o^\infty(]0, 1[)$ , equal to 1 near  $\frac{1}{2}$ , we get the asymptotic expansion of  $\sum \rho(\mu - \mu_j)$  which is the content of the formula of Duistermaat-Guillemin [Du-Gu75]:

$$\sum \rho(\mu - \mu_j) \sim \mu^{-1}D(\mu) ,$$

where  $\sim$  means that the asymptotic expansions are the same.

### 3.10 Degenerate cases

The trace formula can be extended to much more degenerated cases. The non trivial contributions to  $D(E)$  will come from oscillating integrals with phases whose critical points are bijectively associated with closed orbits. It implies that general results on stationary phase expansions depending on the theory of singularities can be applied (see for example [AR86, Mal75]).

### 3.11 The integrable case

As observed by Berry-Tabor [Be-Ta76], the trace formula in this case comes from the Poisson summation formula. Asymptotics of the eigenvalues to any order can then be given in the so-called quantum integrable case by Bohr-Sommerfeld rules.

Using (semi-classical) action-angle coordinates, we start with the Hamiltonian on the torus  $\mathbb{R}^d/2\pi\mathbb{Z}^d$  defined by

$$\hat{H}\exp(i\langle\nu|x\rangle) = H(h\nu)\exp(i\langle\nu|x\rangle)$$

and compute the trace  $Z_a(t)$  of  $a(hD_x)\exp(-it\hat{H}/h)$  using the expression of the eigenvalues  $H(h\nu)$ :

$$Z_a(t) = \sum_{\nu\in\mathbb{Z}^d} a(h\nu)e^{-itH(h\nu)/h} .$$

We will apply Poisson summation formula as well as the approximation of the  $h$ -Fourier transform of  $a(\xi)\exp(-itH(\xi))$  given from stationary phase. The ND condition for stationary phase will be that  $\xi \rightarrow \nabla H(\xi)$  is a local diffeo. on the support of  $a$ . After some calculations we get, for  $t \neq 0$ :

$$Z(t) \sim \frac{1}{(2\pi th)^{d/2}} \sum_{\gamma\in\mathbb{Z}^d} e^{itA(\gamma)/h} (2\pi)^d |\det(H''_{\xi\xi}(\xi_\gamma))|^{-\frac{1}{2}}$$

with  $t\nabla H(\xi_\gamma) = \gamma$  and  $tA(\gamma)$  the action of the closed orbit  $\gamma$ .

### 3.12 The maximally degenerated case

Let us assume that  $(X, g)$  is a compact Riemannian manifold for which all geodesics have the same smallest period  $T_0 = 2\pi$ . Then we have the following clustering property [Wei77, CdV79a, Bo-Gu81]:

**Theorem 7** *There exists some constant  $C$  and some integer  $\alpha$  so that*

1. *the spectrum of  $\Delta$  is contained in the union of the intervalls*

$$I_k = [(k + \frac{\alpha}{4})^2 - C, (k + \frac{\alpha}{4})^2 + C]$$

2.  $N(k) = \#\text{Spectrum}(\Delta) \cap I_k$  is a polynomial function of  $k$  for  $k$  large enough.

The property 2 is consequence of the trace formula [CdV79a].

Another application is:

**Example 3.3 (Rational harmonic oscillators)**

Let us consider the harmonic oscillator

$$\hat{H} = \frac{h^2}{2}\Delta + \frac{1}{2}(x^2 + 4y^2)$$

in  $(\mathbb{R}^2, \text{Eucl})$  whose spectrum is  $E_N = h(N + \frac{3}{2})$  with multiplicities  $a_N = \#\{2k + l = N \mid k, l \geq 0\}$ . One can check that

$$a_N = \frac{1}{2}(N + \frac{3}{2}) + \frac{(-1)^N}{4} . \tag{9}$$

Let us consider the SCTF with a function  $\rho$  so that the compactly supported Fourier transform  $\hat{\rho}$  satisfies  $\sum_{l \in \mathbb{Z}} \hat{\rho}(t - 2\pi l) \equiv 1$ . We have then by applying Equation (4) with  $E = 1$ ,  $h = (N_0 + \frac{3}{2})^{-1}$ :

$$D(1) = \frac{N_0 + \frac{3}{2}}{2\pi} \sum_{N=0}^{\infty} a_N \int_{\mathbb{R}} \hat{\rho}(t) \psi\left(\frac{N + \frac{3}{2}}{N_0 + \frac{3}{2}}\right) e^{it(N_0 - N)} dt$$

which is equal to  $(N_0 + \frac{3}{2})a_{N_0}$  and hence

$$a_N = (N + \frac{3}{2})^{-1}(D_{\text{Weyl}} + D_{\gamma}) + o(1)$$

We easily compute, using formulas (5) and (6)

•

$$D_{\text{Weyl}} = \left(\frac{N + \frac{3}{2}}{2\pi}\right)^2 \left(\frac{d}{dE}\right)_{E=1} \text{Vol}\left\{\frac{1}{2}(\xi^2 + \eta^2) + \frac{1}{2}(x^2 + 4y^2) \leq E\right\} = \frac{(N + \frac{3}{2})^2}{2} + O(1)$$

- $D_{\gamma}$  for the “short” orbit  $\gamma(t) = (x = 0, \xi = 0; y = \cos 2t/\sqrt{2}, \eta = -2 \sin 2t/\sqrt{2})$ , with  $T_0 = \pi$ ,  $\Pi_{\gamma} = -\text{Id}$ ,  $A(\gamma) = 2\pi$ :  $D_{\gamma} = (N + \frac{3}{2})\frac{(-1)^N}{4}$

and deduce an equality

$$a_N = \frac{1}{2}(N + \frac{3}{2}) + \frac{(-1)^N}{4} + \varepsilon_N ,$$

with  $\varepsilon_N \rightarrow 0$ . Because  $\varepsilon_N$  is an integer,  $\varepsilon_N \equiv 0$  for  $N$  large. The previous type of result can be extended to any rational harmonic oscillator (see the contribution of B. Zhilinskii in [Mi01] pp. 126-136).

### 3.13 Applications to the inverse spectral problem

We will now restrict ourselves to the case of the Laplace operator on a compact Riemannian manifold  $(X, g)$ . The following result is a corollary of SCTF:

**Theorem 8** ([CdV73a, CdV73b]) *If  $X$  is given, there exists a generic subset  $\mathcal{G}_X$ , in the sense of Baire category, of the set of smooth Riemannian metrics on  $X$ , so that, if  $g \in \mathcal{G}_X$ , the length spectrum of  $(X, g)$  can be recovered from the Laplace spectrum. The set  $\mathcal{G}_X$  contains all metric with  $< 0$  sectional curvature and (conjecturally) all metrics with  $\leq 0$  sectional curvature.*

We can take the set of metrics for which all closed geodesics are non degenerate and the length spectrum is simple.

## 4 Sturm-Liouville determinants and the metaplectic representation

### 4.1 The stationary phase approximation

Let us consider the following integral

$$I(h) = (2\pi i h)^{-N/2} \int_{\mathbb{R}^N} e^{iS(x)/h} a(x) |dx|$$

where  $S : \mathbb{R}^N \rightarrow \mathbb{R}$  is smooth and  $a \in C_o^\infty(\mathbb{R}^N)$ .

**Theorem 9 (stationary phase)**

- If  $S$  has no critical point in the support of  $a$ ,  $I(h) = O(h^\infty)$
- If the critical points of  $S$  in the support of  $a$  belongs to a non degenerate connected critical manifold  $W$  of dimension  $n$ ,

$$I(h) = (2\pi i h)^{-n/2} e^{-i\nu\pi/2} e^{iS(W)/h} \left( \sum_{k=0}^{\infty} c_j h^j \right) + O(h^\infty)$$

with

$$c_0 = \int_W a(y) |dy|$$

where  $\nu$  is the Morse index of  $S$  along  $W$  and  $|dy|$  is the quotient of the measure  $|dx|$  by the ‘‘Riemannian measure’’ on the normal bundle to  $W$  associated to the Hessian of  $S$ :

$$|dy| := \frac{|dx|}{|\det(\partial_{\alpha\beta}^2 S)|^{\frac{1}{2}} |dz|}$$

where  $z = (z_\alpha)$  are coordinates on the normal bundle.

*Proof.*–

Using Morse lemma, the integral can be reduced to the case where there are local coordinates  $(y, z)$  so that  $S(y, z) = \frac{1}{2}Q(z)$  with  $Q$  a non degenerate quadratic form. The proof then works by first integrating w.r. to  $z$  and using an elegant argument due to Hörmander (see [Hör83] s. 7.7) for the case of a non degenerate critical point. □

**Remark 3** *As suggested by Don Zagier, we can reformulate the stationary phase formula as follows: let us consider the case of an ND manifold  $W$  of critical points of dimension  $n$  and the Schwartz distributions  $T_h = (2\pi i h)^{(n-N)/2} e^{iS(x)/h} |dx|$ . Then the weak limit of  $T_h$  is the Radon measure  $e^{-i\nu\pi/2} \mu_W$ .*

## 4.2 The metaplectic representation

Good references for the metaplectic representation are [Fo89] chap. 4 and [Hör85] Section 18.5.

In what follows,  $\varepsilon$  will denote a number in the set  $\{\pm 1, \pm i\}$ .

A symplectic linear map  $\chi : T^*\mathbb{R}^d \rightarrow T^*\mathbb{R}^d$  defined by

$$[\chi] = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$$

is a *linear* twist symplectic map if and only if  $\beta$  is invertible;  $\chi$  admits then an unique quadratic generating function  $q(x, y) = \frac{1}{2}\langle Ax|x\rangle + \langle By|x\rangle + \frac{1}{2}\langle Cy|y\rangle$  with  $A, C$  symmetric matrices and  $B$  an invertible matrix. We have:

$$[\chi] = \begin{pmatrix} -B^{-1}C & -B^{-1} \\ {}^tB - AB^{-1}C & -AB^{-1} \end{pmatrix} .$$

We will define four operators  $\hat{\chi}$  on  $L^2(\mathbb{R}^d)$  by their Schwartz kernels

$$[\hat{\chi}](x, y) = \varepsilon (2\pi i h)^{-d/2} e^{iq(x, y)/h} |\det(B)|^{\frac{1}{2}} .$$

Using unitarity of Fourier transforms, we see that  $\hat{\chi}$  is an unitary map of  $L^2(\mathbb{R}^d)$ . Moreover, if  $\chi_1, \chi_2, \chi_2 \circ \chi_1$  are twist maps, we have

$$\widehat{\chi_2 \chi_1} = \varepsilon \widehat{\chi_2 \circ \chi_1}$$

as follows from the calculus of Fresnel integrals. The closure of all  $\hat{\chi}$ 's, with  $\chi$  linear twist maps, is a Lie subgroup  $M(d)$  of the Hilbert unitary group  $U(L^2(\mathbb{R}^d))$ . The mapping  $\hat{\chi} \rightarrow \chi$  extends to a group morphism of  $M(d)$  onto the symplectic group  $Sp(d)$  whose kernel is  $\mathbb{Z}/4\mathbb{Z}$ . The connected component of the identity of  $M(d)$  is a two-fold covering of the symplectic groupe called the *metaplectic group*  $Mp(d)$  and the previous recipe gives a natural unitary representation of  $Mp(d)$  into  $L^2(\mathbb{R}^d)$  called the *metaplectic representation*.

### 4.3 Metaplectic traces

Metaplectic maps are not trace class, but they admit traces in the sense of distribution as follows:

let us consider, for  $\chi$  a linear symplectic map, the distribution  $I_\chi$  on  $\mathbb{R}^{2d}$  defined by  $I_\chi(p) = \text{Trace}(\hat{\chi}\text{Op}(p))$  where  $\text{Op}(p)$  is the Weyl quantization of  $p \in C_o^\infty(T^*\mathbb{R}^d)$  defined by

$$[\text{Op}(p)](x, y) = (2\pi h)^{-d} \int_{\mathbb{R}^d} e^{i\langle x-y|\xi\rangle/h} p\left(\frac{x+y}{2}, \xi\right) |d\xi| .$$

Let us denote by  $F_\chi = \ker(\chi - \text{Id})$  the space of fixed points of  $\chi$  and  $n = \dim F_\chi$ .

**Theorem 10** *The distributional trace admits the following asymptotic behaviour  $I_\chi(p) \sim \varepsilon(2\pi i h)^{-n/2} \int_{F_\chi} p d\mu_\chi$  where  $d\mu_\chi$  is a Lebesgue measure on  $F_\chi$ .*

*Moreover  $d\mu_\chi$  is a purely symplectic invariant of  $\chi$ : if  $\chi_2 = \chi^{-1}\chi_1\chi$ ,  $d\mu_{\chi_2} = \chi^*(d\mu_{\chi_1})$ .*

*Proof.*–

The first assertion comes from stationary phase approximation.

The second comes from *exact* Egorov theorem (see [Hör85] p. 158):  $\hat{\chi}\text{Op}(p)\hat{\chi}^{-1} = \text{Op}(p \circ \chi)$  and the circularity of the trace.

□

**Remark 4** *If  $\chi$  is twist, one can reduce to  $p = p(x)$  and use stationary phase on  $\mathbb{R}^d$  instead  $\mathbb{R}^{2d}$ .*

### 4.4 Measures on fixed point sets

Using the fact that  $d\mu_\chi$  is invariant by conjugacy, we can reduce the computations to suitable normal forms:

**Example 4.1** • *If  $F = 0$ ,  $d\mu_\chi = |\det(\text{Id} - \chi)|^{-\frac{1}{2}}\delta(0)$ .*

• *If  $[\chi] = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$  in a symplectic basis  $(\partial_x, \partial_\xi)$  with  $a \neq 0$ ,  $d\mu_\chi = |a|^{-\frac{1}{2}}|dx|$ .*

*Both examples can be proved using twist maps and evaluating the trace by stationary phase from a generating function.*

*For the first example, we can take:*

$$q(x, y) = \frac{1}{2}\langle Ax|x\rangle + \langle By|x\rangle + \frac{1}{2}\langle Cy|y\rangle$$

*with  $B$  invertible, and check the identity:*

$$\det(\text{Id} - \chi)\det(B) = \det(A + B + {}^t B + C) .$$

For the second one take

$$q(x, y) = \frac{1}{2a}(x - y)^2 .$$

It is easy to extend the previous construction to the case of a symplectic diffeomorphism with a clean manifold of fixed points  $W$  getting a measure  $\mu_W$  on  $W$ .

## 4.5 Applications to discrete Sturm-Liouville with Dirichlet or periodic boundary conditions

We assume that  $E$ , a  $d$ -dimensional real vector space, is equipped with a Lebesgue measure  $|dx|$ . We will consider a ‘‘Jacobi matrix’’  $[L]$  on  $E^{\oplus N+1}$  given by

$$[L] := \begin{pmatrix} A_0 & B_0 & 0 & \cdots & 0 & 0 \\ {}^tB_0 & A_1 & B_1 & \cdots & 0 & 0 \\ 0 & {}^tB_1 & A_2 & \cdots & 0 & 0 \\ 0 & 0 & \cdots & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & \cdots & A_{N-1} & B_{N-1} \\ 0 & 0 & \cdots & \cdots & {}^tB_{N-1} & 0 \end{pmatrix} ,$$

and denote by  $Q = Q_L$  the associated quadratic form

$$Q(x_0, \cdots, x_N) = \frac{1}{2} \langle Lx | x \rangle .$$

Our goal is to compute the determinants of  $L$ , of the restriction  $L_0$  of  $L$  to  $x_0 = x_N = 0$  and of the ‘‘restriction’’  $L_{\text{per}}$  of  $L$  to  $x_0 = x_N$ :

$$[L_{\text{per}}] := \begin{pmatrix} A_0 & B_0 & 0 & \cdots & 0 & B_{N-1} \\ {}^tB_0 & A_1 & B_1 & \cdots & 0 & 0 \\ 0 & {}^tB_1 & A_2 & \cdots & 0 & 0 \\ 0 & 0 & \cdots & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ {}^tB_{N-1} & 0 & \cdots & \cdots & {}^tB_{N-2} & A_{N-1} \end{pmatrix} .$$

Let us denote by  $b_i = \det(-B_i)$ , by  $\chi_i$  the canonical transformation generated by  $q_i(u, v) = \frac{1}{2} \langle A_i u | u \rangle + \langle B_i v | u \rangle$  and

$$\chi = \chi_{N-1} \circ \cdots \circ \chi_0 .$$

We have

$$\chi = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} .$$

**Theorem 11** *We have the following formulae:*

$$\det(L) = b_0 \cdots b_{N-1} \det(\gamma) , \quad (10)$$

$$\det(L_0) = b_0 \cdots b_{N-1} \det(\beta) , \quad (11)$$

and

$$\det(L_{\text{per}}) = (-1)^d b_0 \cdots b_{N-1} \det(\text{Id} - \chi) . \quad (12)$$

*Proof.*–

For the formula (11), we see that

$$[\widehat{\chi_{N-1}} \circ \cdots \circ \widehat{\chi_0}](x, y) = (2\pi i h)^{-Nd/2} \int e^{Q(y, x_1, \dots, x_{N-1}, y)/h} |b_0 \cdots b_{N-1}|^{\frac{1}{2}} |dx_1 \cdots dx_{N-1}|$$

which is equal to  $[\hat{\chi}]$  and can also be evaluated by stationary phase as  $|b_0 \cdots b_{N-1}|^{\frac{1}{2}} / |\det L|^{\frac{1}{2}}$ . It gives the formula up to a  $\pm$  sign.

The formula (12) is similar evaluating the trace of  $\hat{\chi}$  in 2 ways.

□

From the previous formulae, we can get the Van Vleck formula (Theorem 1) and the trace formula (Theorem 2) for a twist map with isolated periodic points.

## 4.6 Regularized Determinants of continuous Sturm-Liouville operators

Let us consider the scalar differential operator

$$L = -\frac{d^2}{dx^2} + q(x), \quad 0 \leq x \leq T$$

which we will discretize as follows: let  $\varepsilon = T/N$ , and  $L_\varepsilon : \mathbb{R}^{N+1} \rightarrow \mathbb{R}^{N-1}$  defined by

$$(L_\varepsilon x)_j = \frac{2x_j - x_{j-1} - x_{j+1}}{\varepsilon^2} + q(j\varepsilon)x_j, \quad 1 \leq j \leq N .$$

We will consider the quadratic form  $q_\varepsilon$  on  $\mathbb{R}^{N+1}$  defined by:

$$q_\varepsilon(x_0, \dots, x_N) = \frac{1}{2} \langle Lx | x \rangle_{\mathbb{R}^{N-1}} .$$

We introduce the following operators:

- $L_{\varepsilon, \text{Dir}}$  the restriction of  $L_\varepsilon$  to  $x_0 = x_{N+1} = 0$ ,
- $L_{\varepsilon, \text{Neu}}$  the operator on  $\mathbb{R}^{N+1}$  associated to  $q_\varepsilon$
- $L_{\varepsilon, \text{Per}}$  the operator on  $\mathbb{R}^N$  associated to the restriction of  $q_\varepsilon$  to  $x_0 = x_{N+1}$

and  $\chi_\varepsilon : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  the symplectic map defined by

$$\chi_\varepsilon(x_0, \frac{x_1 - x_0}{\varepsilon}) = (x_N, \frac{x_N - x_{N-1}}{\varepsilon}),$$

with  $L_\varepsilon x = 0$  and the matrix of  $\chi_\varepsilon$ :

$$[\chi_\varepsilon] = \begin{pmatrix} \alpha_\varepsilon & \beta_\varepsilon \\ \gamma_\varepsilon & \delta_\varepsilon \end{pmatrix}.$$

From Theorem 11, we get:

- $\det(L_{\varepsilon, \text{Dir}}) = \frac{1}{\varepsilon^N} \gamma_\varepsilon$
- $\det(L_{\varepsilon, \text{Neu}}) = \frac{1}{\varepsilon^N} \beta_\varepsilon$
- $\det(L_{\varepsilon, \text{Per}}) = -\frac{1}{\varepsilon^N} \det(\text{Id} - \chi_\varepsilon)$ .

As  $\varepsilon \rightarrow 0$ , we get

- For  $k = 1, \dots$ , the eigenvalue  $\lambda_k(\varepsilon)$  of  $L_{\varepsilon, \text{Dir}}$  (resp.  $L_{\varepsilon, \text{Neu}}$ , resp.  $L_{\varepsilon, \text{Per}}$ ) converges to the the eigenvalue  $\lambda_k$  of the corresponding boundary value operator associated to  $L$
- $\chi_\varepsilon$  converges to the map  $\chi : (x(0), x'(0)) \rightarrow (x(T), x'(T))$  with  $Lx = 0$ .

It is then possible [CdV99] to deduce the result of Levit-Smilansky [Lev-Smil77]:

**Theorem 12** *Let us consider the Dirichlet eigenvalues  $\lambda_k^i$  of  $L_{q_i}$ ,  $i = 1, 2$ , we have*

$$\prod_{k=1}^{\infty} \frac{\lambda_k^1}{\lambda_k^2} = \frac{\beta_1}{\beta_2},$$

where  $\beta_i$  are the corresponding entries of the matrices of  $\chi_i$ .

*Similar results holds for the 2 other boundary value problems.*

## References

- [AR86] Arnold, V., Varchenko, A. & Goussein-Zade, S., Singularités des applications différentiables. *Mir (Moscou)*, 1986.
- [BB70] Balian, R. & Bloch, C., Distribution of eigenfrequencies for the wave equation in a finite domain I. *Ann. of Physics*, 60:401, 1970.
- [BB71] Balian, R. & Bloch, C., Distribution of eigenfrequencies for the wave equation in a finite domain II. *Ann. of Physics*, 64:271, 1971.

- [BB72] Balian, R. & Bloch, C., Distribution of eigenfrequencies for the wave equation in a finite domain III. *Ann. of Physics*, 69:76, 1972.
- [BB74] Balian, R. & Bloch, C., Solution of the Schrödinger equation in terms of classical paths. *Ann. of Phys.*, 85:514, 1974.
- [BW97] Bates, S. & Weinstein, A., Lectures on the Geometry of Quantization. *Berkeley Math. Lecture Notes, AMS*, Vol. 8, 1997.
- [Be94] Bellissard, J. & al, Transition to Chaos in Classical and Quantum Mechanics. *Lecture Notes in Maths (Springer)*, no 1589, 1994.
- [Be00] Berger, M., Riemannian geometry during the second half of the twentieth century. *University Lecture Series, AMS*. 17, 2000.
- [BGM71] Berger, M., Gauduchon, P., & Mazet, E., Le spectre d'une variété riemannienne compacte. *Springer LNM*, 1971.
- [Be-Ta76] Berry, M. V. & Tabor, M., Closed orbits and the regular bound spectrum. *Proc. Roy. Soc. London Ser. A*, 349:101–123, 1976.
- [Bo-Pa-Sc00] Bogomolny, E., Pavloff, N. & Schmit, C., Diffractive corrections in the trace formula for polygonal billiards. *Phys. Rev. E (3)*, 61:3689–3711, 2000.
- [Bo-Gi-Sc84] Bohigas, O., Giannoni, M.-J. & Schmit, C., Characterization of chaotic quantum spectra and universality of level fluctuation laws. *Phys. Rev. Lett.*, 52:1–4, 1984
- [Bott56] Bott, R., On the iteration of closed geodesics and the Sturm intersection theory. *Comm. Pure Appl. Math.*, 9:171–206, 1956.
- [Bo-Gu81] Boutet de Monvel, L. & Guillemin, V., The spectral theory of Toeplitz operators. *Annals of Mathematics Studies, Princeton*, 99, 1981.
- [Br-Ur91] Brummelhuis, R. & Uribe, A., A trace formula for Schrödinger operators. *Comm. Math. Phys.*, 136: 567–584, 1991.
- [Cha-Po99] Charbonnel, A.-M. & Popov, G., A semi-classical trace formula for several commuting operators. *Comm. Partial Differential Equations*, 24:283–323, 1999.
- [Cha74] Chazarain, J., Formule de Poisson pour les variétés riemanniennes. *Invent. Math.*, 24: 65–82, 1974.
- [CdV73a] Colin de Verdière, Y., Spectre du Laplacien et longueurs des géodésiques périodiques I. *Compositio Mathematica*, 27: 80–106, 1973.

- [CdV73b] Colin de Verdière, Y., Spectre du Laplacien et longueurs des géodésiques périodiques II. *Compositio Mathematica*, 27: 159–184, 1973.
- [CdV79a] Colin de Verdière, Y., Sur le spectre des opérateurs elliptiques à bicaractéristiques toutes périodiques. *Comment. Math. Helv.*, 54: 508–522, 1979.
- [CdV99] Colin de Verdière, Y., Déterminants et intégrales de Fresnel. *Ann. Inst. Fourier*, 49:861–881, 1999.
- [CdV05] Colin de Verdière, Y., Bohr-Sommerfeld rules to all orders. *Henri Poincaré Acta*, 2005.
- [Dui76] Duistermaat, J., On the Morse index in variational calculus. *Advances in Math.*, 21:173–195, 1976.
- [Du-Gu75] Duistermaat, J. & Guillemin, V., The spectrum of positive elliptic operators and periodic geodesics. *Invent. Math.*, 29: 39–79, 1975.
- [Fe-Hi65] Feynman, R. & Hibbs, A., *Quantum mechanics and path integrals*. McGraw-Hill (New-York), 1965.
- [Fo89] Folland, G., *Harmonic Analysis in Phase Space*. Princeton UP, 1989.
- [Gu75] Guillemin, V., Clean intersection theory and Fourier integrals. *Fourier integral operators and partial differential equations (Colloq. Internat., Univ. Nice, Nice, 1974)*, Lecture Notes in Math., Vol. 459, 23–35, 1975.
- [Gu96] Guillemin, V., Wave-trace invariants. *Duke Math. Jour.*, 83: 287–352, 1996.
- [Gu-Me79] Guillemin, V. & Melrose, R., The Poisson summation formula for manifolds with boundary. *Adv. in Math.*, 32:204–232, 1979.
- [Gut71] Gutzwiller, M., Periodic orbits and classical quantization conditions. *J. of Math. Phys.*, 12:343–358, 1971.
- [Hej76] Hejhal, D., The Selberg trace formula and the Riemann  $\zeta$  function. *Duke Math. J.*, 43: 441–482, 1976.
- [Hill02] Hillairet, L., PhD thesis, Grenoble, 2002.
- [Ho-Ze94] Hofer, H. & Zehnder, E., *Symplectic Invariants and Hamiltonian Dynamics*. Birkhäuser, 1994.
- [Hör68] Hörmander, L., The spectral function of an elliptic operator. *Acta Math.*, 121:193–218, 1968.

- [Hör83] Hörmander, L., The Analysis of Linear Partial Differential Operators I. *Grundlehren (Springer)*, 1983.
- [Hör85] Hörmander, L., The Analysis of Linear Partial Differential Operators III. *Grundlehren (Springer)*, 1985.
- [Hub59] Huber, H., Zur analytischen Theorie hyperbolischen Raumformen und Bewegungsgruppen. *Math. Ann.*, 138:1–26, 1959.
- [Kac66] Kac, M., Can one hear the shape of a drum? *Am. Math. Mon.*, 73: 1–23, 1966.
- [K-T91] Kozlov, V. & Treshchëv, D., Billiards: a genetic introduction to the dynamics of systems with impacts. *Translations of mathematical monographs (American Mathematical Society)*, 089, 1991.
- [La57] Lax, P. D., Asymptotic solutions of oscillatory initial value problems. *Duke Math. J.*, 24:627–646, 1957.
- [Lev-Smil77] Levit, S. & Smilansky, U., A theorem on infinite products of eigenvalues of Sturm type operators. *Proc. AMS.*, 65: 299–303, 1977.
- [Mal75] Malgrange, B., Intégrales asymptotiques et monodromie, *Ann. Sci. École Norm. Sup.*, 7:405–430, 1974.
- [Mar03] Marklof, J., Selberg’s trace formula: an introduction. <http://www.maths.bris.ac.uk/~majm/bib/selberg.ps>
- [Mei92] Meinrenken, E., Semi-classical principal symbols and Gutzwiller’s trace formula. *Rep. Math. Phys.*, 31: 279–295, 1992.
- [Mei94] Meinrenken, E., Trace formulas and Conley-Zehnder index. *Jour. Geom. Phys.*, 13:1–15, 1994.
- [Mi01] Michel, L. (editor), Symmetry, invariants, topology. *Physics reports*, 341 (1-6), 2001.
- [Mil67] Milnor, J., *Morse theory*. Princeton, 1967.
- [Mo51] Morette, C., On the definition and approximation of Feynman’s path integrals. *Physical Rev. (2)*, 81:848–852, 1951.
- [Ra-Si71] Ray, D. B. & Singer, I. M.,  $R$ -torsion and the Laplacian on Riemannian manifolds. *Advances in Math.*, 7:145–210, 1971.
- [Sel56] Selberg, A., Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series. *J. Indian Math. Soc.*, 20: 47–87, 1956.

- [Se50] Serre, J.-P., Homologie singulière des espaces fibrés. Applications. *Ann. of Math. (2)*, 54:425–505, 1951.
- [Wei75] Weinstein, A., On Maslov’s quantization condition. *Fourier integral operators and partial differential equations (Colloq. Internat., Univ. Nice, Nice, 1974)*, Lecture Notes in Math., Vol. 459, 341–372, 1975.
- [Wei77] Weinstein, A., Asymptotics of eigenvalue clusters for the Laplacian plus a potential. *Duke Math. J.*, 44:883–892, 1977.
- [Yo69] Yorke, J., Periods of periodic solutions and the Lipschitz constant. *Proc. Amer. Math. Soc.*, 22:509–512, 1969.
- [Zel97] Zelditch, S., Wave trace invariants at elliptic closed geodesics. *GAF*, 7:145–213, 1997.
- [Zel98] Zelditch, S., Wave invariants for non-degenerate closed geodesics. *GAF*, 8:179–207, 1998.