

ANALYSIS & PDE

Volume 8

No. 4

2015

SEMYON DYATLOV, FRÉDÉRIC FAURE AND COLIN GUILLARMOU

**POWER SPECTRUM OF THE GEODESIC FLOW
ON HYPERBOLIC MANIFOLDS**



POWER SPECTRUM OF THE GEODESIC FLOW ON HYPERBOLIC MANIFOLDS

SEMYON DYATLOV, FRÉDÉRIC FAURE AND COLIN GUILLARMOU

We describe the complex poles of the power spectrum of correlations for the geodesic flow on compact hyperbolic manifolds in terms of eigenvalues of the Laplacian acting on certain natural tensor bundles. These poles are a special case of Pollicott–Ruelle resonances, which can be defined for general Anosov flows. In our case, resonances are stratified into bands by decay rates. The proof also gives an explicit relation between resonant states and eigenstates of the Laplacian.

1. Introduction	923
2. Outline and structure	929
3. Geometry of the hyperbolic space	933
4. Horocyclic operators	941
5. Pollicott–Ruelle resonances	952
6. Properties of the Laplacian	966
7. Expansions of eigenstates of the Laplacian	978
Appendix A. Some technical calculations	988
Appendix B. The special case of dimension 2	993
Appendix C. Eigenvalue asymptotics for symmetric tensors	994
Acknowledgements	997
References	998

1. Introduction

In this paper, we consider the characteristic frequencies of correlations,

$$\rho_{f,g}(t) = \int_{SM} (f \circ \varphi_{-t}) \cdot \bar{g} \, d\mu, \quad f, g \in C^\infty(SM), \tag{1-1}$$

for the geodesic flow φ_t on a compact hyperbolic manifold M of dimension $n + 1$ (that is, M has constant sectional curvature -1). Here φ_t acts on SM , the unit tangent bundle of M , and μ is the natural smooth probability measure. Such φ_t are classical examples of *Anosov flows*; for this family of examples, we are able to prove much more precise results than in the general Anosov case.

An important question, expanding on the notion of mixing, is the behavior of $\rho_{f,g}(t)$ as $t \rightarrow +\infty$. Following [Ruelle 1986], we take the *power spectrum*, which in our convention is the Laplace transform $\hat{\rho}_{f,g}(\lambda)$ of $\rho_{f,g}$ restricted to $t > 0$. The long-time behavior of $\rho_{f,g}(t)$ is related to the properties of

MSC2010: 37D40.

Keywords: Pollicott–Ruelle resonances, hyperbolic manifolds.




Figure 1. An illustration of [Theorem 1](#), with eigenvalues of the Laplacian on the left and the resonances of geodesic flow on the right. The red crosses mark exceptional points where the theorem does not apply.

the meromorphic extension of $\hat{\rho}_{f,g}(\lambda)$ to the entire complex plane. The poles of this extension, called *Pollicott–Ruelle resonances* (see [\[Pollicott 1986; Ruelle 1986; Faure and Sjöstrand 2011\]](#) and (1-7) below), are the complex characteristic frequencies of $\rho_{f,g}$, describing its decay and oscillation and not depending on f, g .

For the case of dimension $n + 1 = 2$, the following connection between resonances and the spectrum of the Laplacian was announced in [\[Faure and Tsujii 2013b, Section 4\]](#) (see [\[Flaminio and Forni 2003\]](#) for a related result and the remarks below regarding the zeta function techniques).

Theorem 1. *Assume that M is a compact hyperbolic surface ($n = 1$) and the spectrum of the positive Laplacian on M is (see [Figure 1](#))*

$$\text{Spec}(\Delta) = \{s_j(1 - s_j)\}, \quad s_j \in [0, 1] \cup \left(\frac{1}{2} + i\mathbb{R}\right).$$

Then Pollicott–Ruelle resonances for the geodesic flow on SM in $\mathbb{C} \setminus (-1 - \frac{1}{2}\mathbb{N}_0)$ are

$$\lambda_{j,m} = -m - 1 + s_j, \quad m \in \mathbb{N}_0. \tag{1-2}$$

Remark. We use the Laplace transform (which has poles in the left half-plane) rather than the Fourier transform as in [\[Ruelle 1986; Faure and Sjöstrand 2011\]](#) to simplify the relation to the parameter s used for Laplacians on hyperbolic manifolds.

Our main result concerns the case of higher dimensions $n + 1 > 2$. The situation is considerably more involved than in the case of [Theorem 1](#), featuring the spectrum of the Laplacian on certain tensor bundles. More precisely, for $\sigma \in \mathbb{R}$, denote

$$\text{Mult}_\Delta(\sigma, m) := \dim \text{Eig}^m(\sigma),$$




Figure 2. An illustration of [Theorem 2](#) for $n = 3$. The red crosses mark exceptional points where the theorem does not apply. Note that the points with $m = 2, \ell = 1$ are simply the points with $m = 0, \ell = 0$ shifted by -2 (modulo exceptional points), as illustrated by the arrow.

where $\text{Eig}^m(\sigma)$, defined in (5-19), is the space of *trace-free, divergence-free symmetric sections* of $\otimes^m T^*M$ satisfying $\Delta f = \sigma f$. Denote by $\text{Mult}_R(\lambda)$ the geometric multiplicity of λ as a Pollicott–Ruelle resonance of the geodesic flow on M (see [Theorem 3](#) and the remarks preceding it for a definition).

Theorem 2. *Let M be a compact hyperbolic manifold of dimension $n+1 \geq 2$. Assume $\lambda \in \mathbb{C} \setminus (-\frac{1}{2}n - \frac{1}{2}\mathbb{N}_0)$. Then, for $\lambda \notin -2\mathbb{N}$, we have (see [Figure 2](#))*

$$\text{Mult}_R(\lambda) = \sum_{m \geq 0} \sum_{\ell=0}^{\lfloor m/2 \rfloor} \text{Mult}_\Delta(-(\lambda + m + \frac{1}{2}n)^2 + \frac{1}{4}n^2 + m - 2\ell, m - 2\ell) \tag{1-3}$$

and, for $\lambda \in -2\mathbb{N}$, we have

$$\text{Mult}_R(\lambda) = \sum_{\substack{m \geq 0 \\ m \neq -\lambda}} \sum_{\ell=0}^{\lfloor m/2 \rfloor} \text{Mult}_\Delta(-(\lambda + m + \frac{1}{2}n)^2 + \frac{1}{4}n^2 + m - 2\ell, m - 2\ell). \tag{1-4}$$

Remark. (i) If $\text{Mult}_\Delta(-(\lambda + m + \frac{1}{2}n)^2 + \frac{1}{4}n^2 + m - 2\ell, m - 2\ell) > 0$, then [Lemma 6.1](#) and the fact that $\Delta \geq 0$ on functions imply that either $\lambda \in -m - \frac{1}{2}n + i\mathbb{R}$ or

$$\begin{aligned} \lambda &\in [-1 - m, -m] && \text{if } n = 1, m > 2\ell, \\ \lambda &\in [1 - n - m, -1 - m] && \text{if } n > 1, m > 2\ell, \\ \lambda &\in [-n - m, -m] && \text{if } m = 2\ell. \end{aligned} \tag{1-5}$$

In particular, we confirm that resonances lie in $\{\text{Re } \lambda \leq 0\}$ and the only resonance on the imaginary axis is $\lambda = 0$ with $\text{Mult}_R(0) = 1$, corresponding to $m = \ell = 0$. We call the set of resonances corresponding

to some m the m -th band. This is a special case of the band structure for general contact Anosov flows established in the work of Faure and Tsujii [2013a; 2013b; 2014].

(ii) The case $n = 1$ fits into Theorem 2 as follows: for $m \geq 2$, the spaces $\text{Eig}^m(\sigma)$ are trivial unless σ is an exceptional point (since the corresponding spaces $\text{Bd}^{m,0}(\lambda)$ of Lemma 5.6 would have to be trace-free sections of a one-dimensional vector bundle), and the spaces $\text{Eig}^1(\sigma + 1)$ and $\text{Eig}^0(\sigma)$ are isomorphic, as shown in Appendix C2.

(iii) The band with $m = 0$ corresponds to the spectrum of the scalar Laplacian; the band with $m = 1$ corresponds to the spectrum of the Hodge Laplacian on coclosed 1-forms; see Appendix C2.

(iv) As seen from (1-3) and (1-4), for $m \geq 2$ the m -th band of resonances contains shifted copies of bands $m - 2, m - 4, \dots$. The special case (1-4) means that the resonance 0 of the $m = 0$ band is not copied to other bands.

(v) A Weyl law holds for the spaces $\text{Eig}^m(\sigma)$; see Appendix C1. It implies the following Weyl law for resonances in the m -th band:

$$\sum_{\lambda \in -n/2 - m + i[-R, R]} \text{Mult}_R(\lambda) = \frac{2^{-n} \pi^{-(n+1)/2}}{\Gamma(\frac{1}{2}(n+3))} \cdot \frac{(m+n-1)!}{m!(n-1)!} \text{Vol}(M) R^{n+1} + \mathcal{O}(R^n). \tag{1-6}$$

The power R^{n+1} agrees with the Weyl law of [Faure and Tsujii 2013b, (5.3)] and with the earlier upper bound of [Datchev et al. 2014]. We also see that, if $n > 1$, then each m and $\ell \in [0, \frac{1}{2}m]$ produce a nontrivial contribution to the set of resonances. The factor $(m+n-1)!/m!(n-1)!$ is the dimension of the space of homogeneous polynomials of order m in n variables; it is natural in light of [Faure and Tsujii 2013a, Proposition 5.11], which locally reduces resonances to such polynomials.

The proof of Theorem 2 is outlined in Section 2. We use in particular the microlocal method of Faure and Sjöstrand [2011], defining Pollicott–Ruelle resonances as the points $\lambda \in \mathbb{C}$ for which the (unbounded nonselfadjoint) operator

$$X + \lambda : \mathcal{H}^r \rightarrow \mathcal{H}^r, \quad r > -C_0 \text{Re } \lambda, \tag{1-7}$$


is not invertible. Here X is the vector field on SM generating the geodesic flow, so that $\varphi_t = e^{tX}$, \mathcal{H}^r is a certain anisotropic Sobolev space, and C_0 is a fixed constant independent of r ; see Section 5A for details. Resonances do not depend on the choice of r . The relation to correlations (1-1) is given by the formula

$$\hat{\rho}_{f,g}(\lambda) = \int_0^\infty e^{-\lambda t} \rho_{f,g}(t) dt = \int_0^\infty e^{-\lambda t} \langle e^{-tX} f, g \rangle dt = \langle (X + \lambda)^{-1} f, g \rangle_{L^2(SM)},$$

valid for $\text{Re } \lambda > 0$ and $f, g \in C^\infty(SM)$. See also Theorem 4 below.

We stress that our method provides an explicit relation between classical and quantum states, that is, between Pollicott–Ruelle resonant states (elements of the kernel of (1-7)) and eigenstates of the Laplacian; namely, in addition to the poles of $\hat{\rho}_{f,g}(\lambda)$, we describe its residues. For instance, for the $m = 0$ band, if $u(x, \xi)$, $x \in M$, $\xi \in S_x M$, is a resonant state, then the corresponding eigenstate of the Laplacian, $f(x)$, is obtained by integration of u along the fibers $S_x M$; see (2-3). On the other hand, to obtain u from f one needs to take the boundary distribution w of f , which is a distribution on the conformal boundary \mathbb{S}^n of

the hyperbolic space \mathbb{H}^{n+1} appearing as the leading coefficient of a weak asymptotic expansion at \mathbb{S}^n of the lift of f to \mathbb{H}^{n+1} . Then u is described by w via an explicit formula, (2-4); this formula features the Poisson kernel P and the map $B_- : S\mathbb{H}^{n+1} \rightarrow \mathbb{S}^n$ mapping a tangent vector to the endpoint in negative infinite time of the corresponding geodesic of \mathbb{H}^{n+1} . The explicit relation can be schematically described as follows:



For $m > 0$, one needs to also use *horocyclic differential operators*; see Section 2.

Theorem 2 used the notion of *geometric multiplicity* of a resonance λ , that is, the dimension of the kernel of $X + \lambda$ on \mathcal{H}^r . For nonselfadjoint problems, it is often more natural to consider the *algebraic multiplicity*, the dimension of the space of elements of \mathcal{H}^r which are killed by some power of $X + \lambda$.

Theorem 3. *If $\lambda \notin -\frac{1}{2}n - \frac{1}{2}\mathbb{N}_0$, then the algebraic and geometric multiplicities of λ as a Pollicott–Ruelle resonance coincide.*

Theorem 3 relies on a *pairing formula* (Lemma 5.10), which states that

$$\langle u, u^* \rangle_{L^2(SM)} = F_{m,\ell}(\lambda) \langle f, f^* \rangle_{L^2(M; \otimes^{m-2\ell} T^*M)},$$

where u is a resonant state at some resonance λ corresponding to some m, ℓ in Theorem 2, u^* is a coresonant state (that is, an element of the kernel of the adjoint of $(X + \lambda)$), f, f^* are the corresponding eigenstates of the Laplacian, and $F_{m,\ell}(\lambda)$ is an explicit function. Here $\langle u, u^* \rangle_{L^2}$ refers to the integral $\int u \overline{u^*}$, which is well-defined despite the fact that neither u nor u^* lie in L^2 ; see (5-6). This pairing formula is of independent interest as a step towards understanding the high frequency behavior of resonant states and attempting to prove *quantum ergodicity of resonant states* in the present setting. Anantharaman and Zelditch [2007] obtained the pairing formula in dimension 2 and studied concentration of Patterson–Sullivan distributions, which are directly related to resonant states; see also [Hansen et al. 2012].

To motivate the study of Pollicott–Ruelle resonances, we also apply to our setting the following *resonance expansion*, proved by Tsujii [2010, Corollary 1.2] and Nonnenmacher and Zworski [2015, Corollary 5]:

Theorem 4. *Fix $\varepsilon > 0$. Then, for N large enough and f, g in the Sobolev space $H^N(SM)$,*

$$\rho_{f,g}(t) = \int f d\mu \int g d\mu + \sum_{\lambda \in (-n/2, 0)} \sum_{k=1}^{\text{Mult}_R(\lambda)} e^{\lambda t} \langle f, u_{\lambda,k}^* \rangle_{L^2} \langle u_{\lambda,k}, g \rangle_{L^2} + \mathcal{O}_{f,g}(e^{-(n/2-\varepsilon)t}), \quad (1-8)$$

where $u_{\lambda,k}$ is any basis of the space of resonant states associated to λ and $u_{\lambda,k}^*$ is the dual basis of the space of coresonant states (so that $\sum_k u_{\lambda,k} \otimes_{L^2} u_{\lambda,k}^*$ is the spectral projector of $-X$ at λ).

Here we use [Theorem 3](#) to see that there are no powers of t in the expansion and that there exists the dual basis of coresonant states to a basis of resonant states.

Combined with [Theorem 2](#), the expansion (1-8) in particular gives the optimal exponent in the decay of correlations in terms of the small eigenvalues of the Laplacian; more precisely, the difference between $\rho_{f,g}(t)$ and the product of the integrals of f and g is $\mathcal{O}(e^{-\nu_0 t})$, where

$$\nu_0 = \min_{0 \leq m < n/2} \min \{ \nu + m \mid \nu \in (0, \frac{1}{2}n - m), \nu(n - \nu) + m \in \text{Spec}^m(\Delta) \},$$

or $\mathcal{O}(e^{-(n/2-\varepsilon)t})$ for each $\varepsilon > 0$ if the set above is empty. Here $\text{Spec}^m(\Delta)$ denotes the spectrum of the Laplacian on trace-free, divergence-free symmetric tensors of order m . Using (1-5), we see that in fact one has $\nu \in [1, \frac{1}{2}n - m)$ for $m > 0$.

In order to go beyond the $\mathcal{O}(e^{-(n/2-\varepsilon)t})$ remainder in (1-8), one would need to handle the infinitely many resonances in the $m = 0$ band. This is thought to be impossible in the general context of scattering theory, as the scattering resolvent can grow exponentially near the bands; however, there exist cases, such as Kerr–de Sitter black holes, where a resonance expansion with infinitely many terms holds; see [\[Bony and Häfner 2008; Dyatlov 2012\]](#). The case of black holes is somewhat similar to the one considered here, because in both cases the trapped set is normally hyperbolic; see [\[Dyatlov 2015; Faure and Tsujii 2014\]](#). What is more, one can try to prove a resonance expansion with remainder $\mathcal{O}(e^{-(n/2+1-\varepsilon)t})$, where the sum over resonances in the first band is replaced by $\langle (\Pi_0 f) \circ \varphi^{-t}, g \rangle$ and Π_0 is the projector onto the space of resonant states with $m = 0$, having the microlocal structure of a Fourier integral operator — see [\[Dyatlov 2015\]](#) for a similar result in the context of black holes.

Previous results. In the constant curvature setting in dimension $n+1=2$, the spectrum of the geodesic flow on L^2 was studied by Fomin and Gelfand [\[1952\]](#) using representation theory. An exponential rate of mixing was proved by Ratner [\[1987\]](#) and it was extended to higher dimensions by Moore [\[1987\]](#). In variable negative curvature for surfaces and, more generally, for Anosov flows with stable/unstable jointly nonintegrable foliations, exponential decay of correlations was first shown by Dolgopyat [\[1998\]](#) and then by Liverani [\[2004\]](#) for contact flows. The work of Tsujii [\[2010; 2012\]](#) established the asymptotic size of the resonance-free strip and the work of Nonnenmacher and Zworski [\[2015\]](#) extended this result to general normally hyperbolic trapped sets. Faure and Tsujii [\[2013a; 2013b; 2014\]](#) established the band structure for general smooth contact Anosov flows and proved an asymptotic for the number of resonances in the first band.

In dimension 2, the study of resonant states in the first band ($m = 0$) — that is, distributions which lie in the spectrum of X and are annihilated by the horocyclic vector field U_- — appears already in the works of Guillemin [\[1977, Lecture 3\]](#) and Zelditch [\[1987\]](#), both using the representation theory of $\text{PSL}(2; \mathbb{R})$, albeit without explicitly interpreting them as Pollicott–Ruelle resonant states. A more general study of the elements in the kernel of U_- was performed by Flaminio and Forni [\[2003\]](#).

An alternative approach to resonances involves the *Selberg and Ruelle zeta functions*. The singularities (zeros and poles) of the Ruelle zeta function correspond to Pollicott–Ruelle resonances on differential forms (see [\[Fried 1986; 1995; Giulietti et al. 2013; Dyatlov and Zworski 2015\]](#)), while the singularities of the Selberg zeta function correspond to eigenvalues of the Laplacian. The Ruelle and Selberg zeta

functions are closely related; see [Leboeuf 2004, Section 5.1, Figure 1; Dyatlov and Zworski 2015, (1.2)] in dimension 2 and [Fried 1986; Bunke and Olbrich 1995, Proposition 3.4] in arbitrary dimensions. However, the Ruelle zeta function does not recover all resonances on functions, due to cancellations with singularities coming from differential forms of different orders. For example, [Juhl 2001, Theorem 3.7] describes the spectral singularities of the Ruelle zeta function for $n = 3$ in terms of the spectrum of the Laplacian on functions and 1-forms, which is much smaller than the set obtained in Theorem 2.

The book of Juhl [2001] and the works of Bunke and Olbrich [1995; 1997; 1999; 2001] study Ruelle and Selberg zeta functions corresponding to various representations of the orthogonal group. They also consider general locally symmetric spaces and address the question of what happens at the exceptional points (which in our case are contained in $-\frac{1}{2}n - \frac{1}{2}\mathbb{N}_0$), relating the behavior of the zeta functions at these points to topological invariants. It is possible that the results [Juhl 2001; Bunke and Olbrich 1995; 1997; 1999; 2001] together with an appropriate representation-theoretic calculation recover our description of resonances, even though no explicit description featuring the spectrum of the Laplacian on trace-free, divergence-free symmetric tensors as in (1-3), (1-4) seems to be available in the literature. The direct spectral approach used in this paper, unlike the zeta function techniques, gives an explicit relation between resonant states and eigenstates of the Laplacian (see the remarks following (1-7)) and is a step towards a more quantitative understanding of decay of correlations.

An essential component of our work is the analysis of the correspondence between eigenstates of the Laplacian on \mathbb{H}^{n+1} and distributions on the conformal infinity \mathbb{S}^n . In the scalar case, such a correspondence for hyperfunctions on \mathbb{S}^n is due to Helgason [1970; 1974] (see also [Minemura 1975]); the correspondence between tempered eigenfunctions of Δ and distributions (instead of hyperfunctions) was shown by Oshima and Sekiguchi [1980] and van den Ban and Schlichtkrull [1987] (see also [Grellier and Otal 2005]). Olbrich [1995] studied Poisson transforms on general homogeneous vector bundles, which include the bundles of tensors used in the present paper. The question of regularity of equivariant distributions on \mathbb{S}^n by certain Kleinian groups of isometries of \mathbb{H}^{n+1} (geometrically finite groups) is interesting since it determines the regularity of resonant states for the flow; precise regularity was studied by Otal [1998] in the 2-dimensional cocompact case, Grellier and Otal [2005] in higher dimensions, and Bunke and Olbrich [1999] for geometrically finite groups. In dimension 2, the correspondence between the eigenfunctions of the Laplacian on the hyperbolic plane and distributions on the conformal boundary \mathbb{S}^1 appeared in [Pollicott 1989; Bunke and Olbrich 1997]; it is also an important tool in the theory developed by [Bunke and Olbrich 2001] to study Selberg zeta functions on convex cocompact hyperbolic manifolds (see also [Juhl 2001] in the compact setting). These distributions on the conformal boundary \mathbb{S}^n , of Patterson–Sullivan type, are also the central object of the recent work of Anantharaman and Zelditch [2007; 2012] studying quantum ergodicity on hyperbolic compact surfaces; a generalization to higher-rank, locally symmetric spaces was provided by Hansen, Hilgert and Schröder [Hansen et al. 2012].

2. Outline and structure

In this section, we give the ideas of the proof of Theorem 2, first in dimension 2 and then in higher dimensions, and describe the structure of the paper.

2A. Dimension 2. We start by using the following criterion ([Lemma 5.1](#)): $\lambda \in \mathbb{C}$ is a Pollicott–Ruelle resonance if and only if the space

$$\text{Res}_X(\lambda) := \{u \in \mathcal{D}'(SM) \mid (X + \lambda)u = 0, \text{WF}(u) \subset E_u^*\}$$

is nontrivial. Here $\mathcal{D}'(SM)$ is the space of *distributions* on SM (see [\[Hörmander 1983\]](#)), $\text{WF}(u) \subset T^*(SM)$ is the *wavefront set* of u (see [\[Hörmander 1983, Chapter 8\]](#)), and $E_u^* \subset T^*(SM)$ is the dual unstable foliation described in [\(3-15\)](#). It is more convenient to use the condition $\text{WF}(u) \subset E_u^*$ rather than $u \in \mathcal{H}^r$, because this condition is invariant under differential operators of any order.

The key tools for the proof are the *horocyclic vector fields* U_\pm on SM , pictured in [Figure 3\(a\)](#) below. To define them, we represent $M = \Gamma \backslash \mathbf{H}^2$, where $\mathbf{H}^2 = \{z \in \mathbb{C} \mid \text{Im } z > 0\}$ is the hyperbolic plane and $\Gamma \subset \text{PSL}(2; \mathbb{R})$ is a cocompact Fuchsian group of isometries acting by Möbius transformations. (See [Appendix B](#) for the relation of the notation we use in dimension 2, based on the half-plane model of the hyperbolic space, to the notation used elsewhere in the paper that is based on the hyperboloid model.) Then SM is covered by $S\mathbf{H}^2$, which is isomorphic to the group $G := \text{PSL}(2; \mathbb{R})$ by the map $\gamma \in G \mapsto (\gamma(i), d\gamma(i) \cdot i)$. Consider the left-invariant vector fields on G corresponding to the following elements of its Lie algebra:

$$X = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{pmatrix}, \quad U_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad U_- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}; \tag{2-1}$$

then X, U_\pm descend to vector fields on SM , with X becoming the generator of the geodesic flow. We have the commutation relations

$$[X, U_\pm] = \pm U_\pm \quad \text{and} \quad [U_+, U_-] = 2X. \tag{2-2}$$

For each λ and $m \in \mathbb{N}_0$, define the spaces

$$V_m(\lambda) := \{u \in \mathcal{D}'(SM) \mid (X + \lambda)u = 0, U_-^m u = 0, \text{WF}(u) \subset E_u^*\},$$

and put

$$\text{Res}_X^0(\lambda) := V_1(\lambda).$$

By [\(2-2\)](#), $U_-^m(\text{Res}_X(\lambda)) \subset \text{Res}_X(\lambda + m)$. Since there are no Pollicott–Ruelle resonances in the right half-plane, we conclude that

$$\text{Res}_X(\lambda) = V_m(\lambda) \quad \text{for } m > -\text{Re } \lambda.$$

We now use the diagram (writing $\text{Id} = U_\pm^0, U_\pm = U_\pm^1$ for uniformity of notation)

$$\begin{array}{ccccccc} 0 = V_0(\lambda) & \xrightarrow{\iota} & V_1(\lambda) & \xrightarrow{\iota} & V_2(\lambda) & \xrightarrow{\iota} & V_3(\lambda) & \xrightarrow{\iota} & \dots, \\ & & \begin{array}{c} U_+^0 \uparrow \downarrow U_-^0 \\ \text{Res}_X^0(\lambda) \end{array} & & \begin{array}{c} U_+^1 \uparrow \downarrow U_-^1 \\ \text{Res}_X^0(\lambda + 1) \end{array} & & \begin{array}{c} U_+^2 \uparrow \downarrow U_-^2 \\ \text{Res}_X^0(\lambda + 2) \end{array} & & \\ & & \text{Res}_X^0(\lambda) & & \text{Res}_X^0(\lambda + 1) & & \text{Res}_X^0(\lambda + 2) & & \end{array}$$

where ι denotes the inclusion maps and, unless $\lambda \in -1 - \frac{1}{2}\mathbb{N}_0$, we have

$$V_{m+1}(\lambda) = V_m(\lambda) \oplus U_+^m(\text{Res}_X^0(\lambda + m)),$$

and U_+^m is one-to-one on $\text{Res}_X^0(\lambda + m)$; indeed, using (2-2) we calculate

$$U_-^m U_+^m = m! \left(\prod_{j=1}^m (2\lambda + m + j) \right) \text{Id} \quad \text{on } \text{Res}_X^0(\lambda + m)$$

and the coefficient above is nonzero when $\lambda \notin -1 - \frac{1}{2}\mathbb{N}_0$. We then see that

$$\text{Res}_X(\lambda) = \bigoplus_{m \geq 0} U_+^m(\text{Res}_X^0(\lambda + m)).$$

It remains to describe the space of resonant states in the first band,

$$\text{Res}_X^0(\lambda) = \{u \in \mathcal{D}'(SM) \mid (X + \lambda)u = 0, U_-u = 0, \text{WF}(u) \subset E_u^*\}.$$

We can remove the condition $\text{WF}(u) \subset E_u^*$ as it follows from the other two; see the remark following Lemma 5.6. We claim that the pushforward map

$$u \mapsto f(x) := \int_{S_x M} u(x, \xi) dS(\xi) \tag{2-3}$$

is an isomorphism from $\text{Res}_X^0(\lambda)$ onto $\text{Eig}(-\lambda(1 + \lambda))$, where $\text{Eig}(\sigma) = \{u \in C^\infty(M) \mid \Delta u = \sigma u\}$; this would finish the proof. In other words, the eigenstate of the Laplacian corresponding to u is obtained by integrating u over the fibers of SM .

To show that (2-3) is an isomorphism, we reduce the elements of $\text{Res}_X^0(\lambda)$ to the conformal boundary \mathbb{S}^1 of the ball model \mathbb{B}^2 of the hyperbolic space as follows:

$$\text{Res}_X^0(\lambda) = \{P(y, B_-(y, \xi))^\lambda w(B_-(y, \xi)) \mid w \in \text{Bd}(\lambda)\}, \tag{2-4}$$

where $P(y, v)$ is the Poisson kernel: $P(y, v) = (1 - |y|^2)/|y - v|^2$, $y \in \mathbb{B}^2$, $v \in \mathbb{S}^1$; $B_- : \mathbb{S}\mathbb{B}^2 \rightarrow \mathbb{S}^1$ maps (y, ξ) to the limiting point of the geodesic $\varphi_t(y, \xi)$ as $t \rightarrow -\infty$ — see Figure 3(a) — and $\text{Bd}(\lambda) \subset \mathcal{D}'(\mathbb{S}^1)$ is the space of distributions satisfying a certain equivariance property with respect to Γ . Here we lifted $\text{Res}_X^0(\lambda)$ to distributions on $S\mathbb{H}^2$ and used the fact that the map B_- is invariant under both X and U_- ; see Lemma 5.6 for details.

It remains to show that the map $w \mapsto f$ defined via (2-3) and (2-4) is an isomorphism from $\text{Bd}(\lambda)$ to $\text{Eig}(-\lambda(1 + \lambda))$. This map is given by (see Lemma 6.6)

$$f(y) = \mathcal{P}_\lambda^- w(y) := \int_{\mathbb{S}^1} P(y, v)^{1+\lambda} w(v) dS(v) \tag{2-5}$$

and is the Poisson operator for the (scalar) Laplacian corresponding to the eigenvalue $s(1 - s)$, $s = 1 + \lambda$. This Poisson operator is known to be an isomorphism for $\lambda \notin -1 - \mathbb{N}$ — see the remark following Theorem 6 in Section 5B — finishing the proof.

2B. Higher dimensions. In higher dimensions, the situation is made considerably more difficult by the fact we can no longer define the vector fields U_{\pm} on SM . To get around this problem, we remark that, in dimension 2, U_-u is the derivative of u along a certain canonical vector in the one-dimensional *unstable foliation* $E_u \subset T(SM)$ and, similarly, U_+u is the derivative along an element of the stable foliation E_s ; see Section 4B. In dimension $n + 1 > 2$, the foliations E_u, E_s are n -dimensional and one cannot trivialize them. However, each of these foliations is canonically parametrized by the following vector bundle \mathcal{E} over SM :

$$\mathcal{E}(x, \xi) = \{\eta \in T_x M \mid \eta \perp \xi\}, \quad (x, \xi) \in SM.$$

This makes it possible to define *horocyclic operators*

$$\mathcal{U}_{\pm}^m : \mathcal{D}'(SM) \rightarrow \mathcal{D}'(SM; \otimes_S^m \mathcal{E}^*),$$

where \otimes_S^m stands for the m -th symmetric tensor power, and we have the diagram

$$\begin{array}{ccccccc} 0 = V_0(\lambda) & \xrightarrow{\iota} & V_1(\lambda) & \xrightarrow{\iota} & V_2(\lambda) & \xrightarrow{\iota} & V_3(\lambda) \xrightarrow{\iota} \dots, \\ & & \mathcal{V}_+^0 \updownarrow \mathcal{U}^0 & & \mathcal{V}_+^1 \updownarrow \mathcal{U}^1 & & \mathcal{V}_+^2 \updownarrow \mathcal{U}^2 \\ & & \text{Res}_{\mathcal{X}}^0(\lambda) & & \text{Res}_{\mathcal{X}}^1(\lambda + 1) & & \text{Res}_{\mathcal{X}}^2(\lambda + 2) \end{array}$$

where $\mathcal{V}_+^m = (-1)^m (\mathcal{U}_+^m)^*$ and we put, for a certain extension \mathcal{X} of X to $\otimes_S^m \mathcal{E}^*$,

$$\begin{aligned} V_m(\lambda) &:= \{u \in \mathcal{D}'(SM) \mid (X + \lambda)u = 0, \mathcal{U}_-^m u = 0, \text{WF}(u) \subset E_u^*\}, \\ \text{Res}_{\mathcal{X}}^m(\lambda) &:= \{v \in \mathcal{D}'(SM; \otimes_S^m \mathcal{E}^*) \mid (\mathcal{X} + \lambda)v = 0, \mathcal{U}_- v = 0, \text{WF}(v) \subset E_u^*\}. \end{aligned}$$

Similarly to in dimension 2, we reduce the problem to understanding the spaces $\text{Res}_{\mathcal{X}}^m(\lambda)$, and an operator similar to (2-3) maps these spaces to eigenspaces of the Laplacian on divergence-free symmetric tensors. However, to make this statement precise, we have to further decompose $\text{Res}_{\mathcal{X}}^m(\lambda)$ into terms coming from traceless tensors of degrees $m, m - 2, m - 4, \dots$, explaining the appearance of the parameter ℓ in the theorem. (Here the trace of a symmetric tensor of order m is the result of contracting two of its indices with the metric, yielding a tensor of order $m - 2$.) The procedure of reducing elements of $\text{Res}_{\mathcal{X}}^m(\lambda)$ to the conformal boundary \mathbb{S}^n is also made more difficult because the boundary distributions w are now sections of $\otimes_S^m(T^*\mathbb{S}^n)$.

A significant part of the paper is dedicated to proving that the higher-dimensional analog of (2-5) on symmetric tensors is indeed an isomorphism between appropriate spaces. To show that the Poisson operator \mathcal{P}_{λ}^- is injective, we prove a weak expansion of $f(y) \in C^\infty(\mathbb{B}^{n+1})$ in powers of $1 - |y|$ as $y \in \mathbb{B}^{n+1}$ approaches the conformal boundary \mathbb{S}^n ; since w appears as the coefficient in one of the terms of the expansion, $\mathcal{P}_{\lambda}^- w = 0$ implies $w = 0$. To show the surjectivity of \mathcal{P}_{λ}^- , we prove that the lift to \mathbb{H}^{n+1} of every trace-free, divergence-free eigenstate f of the Laplacian admits a weak expansion at the conformal boundary (this requires a fine analysis of the Laplacian and divergence operators on symmetric tensors); putting w to be the coefficient next to one of the terms of this expansion, we can prove that $f = \mathcal{P}_{\lambda}^- w$.

2C. Structure of the paper. In Section 3, we study in detail the geometry of the hyperbolic space \mathbb{H}^{n+1} , which is the covering space of M . In Section 4, we introduce and study the horocyclic operators. In Section 5, we prove Theorems 2 and 3, modulo properties of the Poisson operator. In Sections 6 and 7, we show the injectivity and the surjectivity of the Poisson operator. Appendix A contains several technical lemmas. Appendix B shows how the discussion of Section 2A fits into the framework of the remainder of the paper. Appendix C shows a Weyl law for divergence-free symmetric tensors and relates the $m = 1$ case to the Hodge Laplacian.

3. Geometry of the hyperbolic space

In this section, we review the structure of the hyperbolic space and its geodesic flow and introduce various objects to be used later, including:

- the isometry group G of the hyperbolic space and its Lie algebra, including the horocyclic vector fields U_i^\pm (Section 3B);
- the stable/unstable foliations E_s, E_u (Section 3C);
- the conformal compactification of the hyperbolic space, the maps B_\pm , the coefficients Φ_\pm , and the Poisson kernel (Section 3D);
- parallel transport to conformal infinity and the maps \mathcal{A}_\pm (Section 3F).

3A. Models of the hyperbolic space. Consider the Minkowski space $\mathbb{R}^{1,n+1}$ with the Lorentzian metric

$$g_M = dx_0^2 - \sum_{j=1}^{n+1} dx_j^2.$$

The corresponding scalar product is denoted $\langle \cdot, \cdot \rangle_M$. We denote by e_0, \dots, e_{n+1} the canonical basis of $\mathbb{R}^{1,n+1}$.

The hyperbolic space of dimension $n + 1$ is defined to be one sheet of the two-sheeted hyperboloid

$$\mathbb{H}^{n+1} := \{x \in \mathbb{R}^{1,n+1} \mid \langle x, x \rangle_M = 1, x_0 > 0\}$$

equipped with the Riemannian metric

$$g_H := -g_M|_{T\mathbb{H}^{n+1}}.$$

We denote the unit tangent bundle of \mathbb{H}^{n+1} by

$$S\mathbb{H}^{n+1} := \{(x, \xi) \mid x \in \mathbb{H}^{n+1}, \xi \in \mathbb{R}^{1,n+1}, \langle \xi, \xi \rangle_M = -1, \langle x, \xi \rangle_M = 0\}. \tag{3-1}$$

Another model of the hyperbolic space is the unit ball $\mathbb{B}^{n+1} = \{y \in \mathbb{R}^{n+1} \mid |y| < 1\}$, which is identified with $\mathbb{H}^{n+1} \subset \mathbb{R}^{1,n+1}$ via the map (here $x = (x_0, x') \in \mathbb{R} \times \mathbb{R}^{n+1}$)

$$\psi : \mathbb{H}^{n+1} \rightarrow \mathbb{B}^{n+1}, \quad \psi(x) = \frac{x'}{x_0 + 1}, \quad \psi^{-1}(y) = \frac{1}{1 - |y|^2}(1 + |y|^2, 2y). \tag{3-2}$$

and the metric g_H pulls back to the following metric on \mathbb{B}^{n+1} :

$$(\psi^{-1})^* g_H = \frac{4 dy^2}{(1 - |y|^2)^2}. \tag{3-3}$$

We will also use the upper half-space model $\mathbb{U}^{n+1} = \mathbb{R}_{z_0}^+ \times \mathbb{R}_z^n$ with the metric

$$(\psi^{-1} \psi_1^{-1})^* g_H = \frac{dz_0^2 + dz^2}{z_0^2}, \tag{3-4}$$

where the diffeomorphism $\psi_1 : \mathbb{B}^{n+1} \rightarrow \mathbb{U}^{n+1}$ is given by (here $y = (y_1, y') \in \mathbb{R} \times \mathbb{R}^n$)

$$\psi_1(y_1, y') = \frac{(1 - |y|^2, 2y')}{1 + |y|^2 - 2y_1}, \quad \psi_1^{-1}(z_0, z) = \frac{(z_0^2 + |z|^2 - 1, 2z)}{(1 + z_0)^2 + |z|^2}. \tag{3-5}$$

3B. Isometry group. We consider the group

$$G = \text{PSO}(1, n + 1) \subset \text{SL}(n + 2; \mathbb{R})$$

of all linear transformations of $\mathbb{R}^{1, n+1}$ preserving the Minkowski metric, the orientation, and the sign of x_0 on timelike vectors. For $x \in \mathbb{R}^{1, n+1}$ and $\gamma \in G$, denote by $\gamma \cdot x$ the result of multiplying x by the matrix γ . The group G is exactly the group of orientation-preserving isometries of \mathbb{H}^{n+1} ; under the identification (3-2), it corresponds to the group of direct Möbius transformations of \mathbb{R}^{n+1} preserving the unit ball.

The Lie algebra of G is spanned by the matrices

$$X = E_{0,1} + E_{1,0}, \quad A_k = E_{0,k} + E_{k,0}, \quad R_{i,j} = E_{i,j} - E_{j,i} \tag{3-6}$$

for $i, j \geq 1$ and $k \geq 2$, where $E_{i,j}$ is the elementary matrix if $0 \leq i, j \leq n + 1$ (that is, $E_{i,j} e_k = \delta_{jk} e_i$). Denote for $i = 1, \dots, n$

$$U_i^+ := -A_{i+1} - R_{1,i+1}, \quad U_i^- := -A_{i+1} + R_{1,i+1} \tag{3-7}$$

and observe that $X, U_i^+, U_i^-, R_{i+1,j+1}$ (for $1 \leq i < j \leq n$) also form a basis. Henceforth we identify elements of the Lie algebra of G with left-invariant vector fields on G .

We have the commutator relations (for $1 \leq i, j, k \leq n$ and $i \neq j$)

$$\begin{aligned} [X, U_i^\pm] &= \pm U_i^\pm, & [U_i^\pm, U_j^\pm] &= 0, & [U_i^+, U_i^-] &= 2X, & [U_i^\pm, U_j^\mp] &= 2R_{i+1,j+1}, \\ [R_{i+1,j+1}, X] &= 0, & [R_{i+1,j+1}, U_k^\pm] &= \delta_{jk} U_i^\pm - \delta_{ik} U_j^\pm. \end{aligned} \tag{3-8}$$

The Lie algebra elements U_i^\pm are very important in our argument, since they generate horocyclic flows; see Section 4B. The flows of U_\pm^1 in the case $n = 1$ are shown in Figure 3(a); for $n > 1$, the flows of U_\pm^j do not descend to $S\mathbb{H}^{n+1}$.

The group G acts on \mathbb{H}^{n+1} transitively, with the isotropy group of $e_0 \in \mathbb{H}^{n+1}$ isomorphic to $\text{SO}(n + 1)$. It also acts transitively on the unit tangent bundle $S\mathbb{H}^{n+1}$, by the rule $\gamma \cdot (x, \xi) = (\gamma \cdot x, \gamma \cdot \xi)$, with the isotropy group of $(e_0, e_1) \in S\mathbb{H}^{n+1}$ being

$$H = \{\gamma \in G \mid \gamma \cdot e_0 = e_0, \gamma \cdot e_1 = e_1\} \simeq \text{SO}(n). \tag{3-9}$$




Figure 3. (a) The horocyclic flows $\exp(\pm U_1^\pm)$ in dimension $n + 1 = 2$, pulled back to the ball model by the map ψ from (3-2). The thick lines are geodesics and the dashed lines are horocycles. (b) The map \mathcal{A}_+ and the parallel transport of an element of \mathcal{E} along a geodesic.

Note that H is the connected Lie subgroup of G with Lie algebra spanned by $R_{i+1, j+1}$ for $1 \leq i, j \leq n$. We can then write $S\mathbb{H}^{n+1} \simeq G/H$, where the projection $\pi_S : G \rightarrow S\mathbb{H}^{n+1}$ is given by

$$\pi_S(\gamma) = (\gamma \cdot e_0, \gamma \cdot e_1) \in S\mathbb{H}^{n+1}, \quad \gamma \in G. \tag{3-10}$$

3C. Geodesic flow. The geodesic flow,

$$\varphi_t : S\mathbb{H}^{n+1} \rightarrow S\mathbb{H}^{n+1}, \quad t \in \mathbb{R},$$

is given in the parametrization (3-1) by

$$\varphi_t(x, \xi) = (x \cosh t + \xi \sinh t, x \sinh t + \xi \cosh t). \tag{3-11}$$

We note that, with the projection $\pi_S : G \rightarrow S\mathbb{H}^{n+1}$ defined in (3-10),

$$\varphi_t(\pi_S(\gamma)) = \pi_S(\gamma \exp(tX)),$$

where X is as defined in (3-6). This means that the generator of the geodesic flow can be obtained by pushing forward the left-invariant field on G generated by X by the map π_S (which is possible since X is invariant under right multiplications by elements of the subgroup H defined in (3-9)). By abuse of notation, we then denote by X also the generator of the geodesic flow on $S\mathbb{H}^{n+1}$:

$$X = \xi \cdot \partial_x + x \cdot \partial_\xi. \tag{3-12}$$

We now provide the stable/unstable decomposition for the geodesic flow, demonstrating that it is hyperbolic (and thus the flow on a compact quotient by a discrete group will be Anosov). For $\rho = (x, \xi) \in S\mathbb{H}^{n+1}$, the tangent space $T_\rho(S\mathbb{H}^{n+1})$ can be written as

$$T_\rho(S\mathbb{H}^{n+1}) = \{(v_x, v_\xi) \in (\mathbb{R}^{1, n+1})^2 \mid \langle x, v_x \rangle_M = \langle \xi, v_\xi \rangle_M = \langle x, v_\xi \rangle_M + \langle \xi, v_x \rangle_M = 0\}.$$

The differential of the geodesic flow acts by

$$d\varphi_t(\rho) \cdot (v_x, v_\xi) = (v_x \cosh t + v_\xi \sinh t, v_x \sinh t + v_\xi \cosh t).$$

We have $T_\rho(S\mathbb{H}^{n+1}) = E^0(\rho) \oplus \tilde{T}_\rho(S\mathbb{H}^{n+1})$, where $E^0(\rho) := \mathbb{R}X$ is the flow direction and

$$\tilde{T}_\rho(S\mathbb{H}^{n+1}) = \{(v_x, v_\xi) \in (\mathbb{R}^{1,n+1})^2 \mid \langle x, v_x \rangle_M = \langle x, v_\xi \rangle_M = \langle \xi, v_x \rangle_M = \langle \xi, v_\xi \rangle_M = 0\},$$

and this splitting is invariant under $d\varphi_t$. A natural norm on $\tilde{T}_\rho(S\mathbb{H}^{n+1})$ is given by the formula

$$|(v_x, v_\xi)|^2 := -\langle v_x, v_x \rangle_M - \langle v_\xi, v_\xi \rangle_M, \tag{3-13}$$

using the fact that v_x and v_ξ are Minkowski orthogonal to the timelike vector x and thus must be spacelike or zero. Note that this norm is invariant under the action of G .

We now define the *stable/unstable decomposition* $\tilde{T}_\rho(S\mathbb{H}^{n+1}) = E_s(\rho) \oplus E_u(\rho)$, where

$$\begin{aligned} E_s(\rho) &:= \{(v, -v) \mid \langle x, v \rangle_M = \langle \xi, v \rangle_M = 0\}, \\ E_u(\rho) &:= \{(v, v) \mid \langle x, v \rangle_M = \langle \xi, v \rangle_M = 0\}. \end{aligned} \tag{3-14}$$

Then $T_\rho(S\mathbb{H}^{n+1}) = E_0(\rho) \oplus E_s(\rho) \oplus E_u(\rho)$, this splitting is invariant under φ_t and under the action of G , and, using the norm from (3-13),

$$|d\varphi_t(\rho) \cdot w| = e^{-t}|w|, \quad w \in E_s(\rho), \quad \text{and} \quad |d\varphi_t(\rho) \cdot w| = e^t|w|, \quad w \in E_u(\rho).$$

Finally, we remark that the vector subbundles E_s and E_u are spanned by the left-invariant vector fields U_1^+, \dots, U_n^+ and U_1^-, \dots, U_n^- from (3-7) in the sense that

$$\pi_S^* E_s = \text{span}(U_1^+, \dots, U_n^+) \oplus \mathfrak{h}, \quad \pi_S^* E_u = \text{span}(U_1^-, \dots, U_n^-) \oplus \mathfrak{h}.$$

Here $\pi_S^* E_s := \{(\gamma, w) \in TG \mid (\pi_S(\gamma), d\pi_S(\gamma) \cdot w) \in E_s\}$ and $\pi_S^* E_u$ is defined similarly; \mathfrak{h} is the left translation of the Lie algebra of H , or equivalently the kernel of $d\pi_S$. Note that, while the individual vector fields U_1^\pm, \dots, U_n^\pm are not invariant under right multiplications by elements of H in dimensions $n + 1 > 2$ (and thus do not descend to vector fields on $S\mathbb{H}^{n+1}$ by the map π_S), their spans are invariant under H , by (3-8).

The dual decomposition, used in the construction of Pollicott–Ruelle resonances, is

$$T_\rho^*(S\mathbb{H}^{n+1}) = E_0^*(\rho) \oplus E_s^*(\rho) \oplus E_u^*(\rho), \tag{3-15}$$

where $E_0^*(\rho), E_s^*(\rho), E_u^*(\rho)$ are dual to $E_0(\rho), E_u(\rho), E_s(\rho)$ in the original decomposition (that is, for instance, $E_s^*(\rho)$ consists of all covectors annihilating $E_0(\rho) \oplus E_s(\rho)$). The switching of the roles of E_s and E_u is due to the fact that the flow on the cotangent bundle is $(d\varphi_t^{-1})^*$.

3D. Conformal infinity. The metric (3-3) in the ball model \mathbb{B}^{n+1} is conformally compact; namely, the metric $(1 - |y|^2)^2(\psi^{-1})^*g_H$ continues smoothly to the closure $\bar{\mathbb{B}}^{n+1}$, which we call the *conformal compactification* of \mathbb{H}^{n+1} ; note that \mathbb{H}^{n+1} embeds into the interior of $\bar{\mathbb{B}}^{n+1}$ by the map (3-2). The boundary $\mathbb{S}^n = \partial\bar{\mathbb{B}}^{n+1}$, endowed with the standard metric on the sphere, is called *conformal infinity*. On the hyperboloid model, it is natural to associate to a point at conformal infinity $v \in \mathbb{S}^n$ the lightlike ray

$\{(s, s\nu) \mid s > 0\} \subset \mathbb{R}^{1,n+1}$; this ray is asymptotic to the curve $\{(\sqrt{1+s^2}, s\nu) \mid s > 0\} \subset \mathbb{H}^{n+1}$, which converges to ν in $\overline{\mathbb{B}^{n+1}}$.

Take $(x, \xi) \in S\mathbb{H}^{n+1}$. Then $\langle x \pm \xi, x \pm \xi \rangle_M = 0$ and $x_0 \pm \xi_0 > 0$, so we can write

$$x \pm \xi = \Phi_{\pm}(x, \xi)(1, B_{\pm}(x, \xi))$$

for some maps

$$\Phi_{\pm} : S\mathbb{H}^{n+1} \rightarrow \mathbb{R}^+, \quad B_{\pm} : S\mathbb{H}^{n+1} \rightarrow \mathbb{S}^n. \tag{3-16}$$

Then $B_{\pm}(x, \xi)$ is the limit as $t \rightarrow \pm\infty$ of the x -projection of the geodesic $\varphi_t(x, \xi)$ in $\overline{\mathbb{B}^{n+1}}$:

$$B_{\pm}(x, \xi) = \lim_{t \rightarrow \pm\infty} \pi(\varphi_t(x, \xi)), \quad \pi : S\mathbb{H}^{n+1} \rightarrow \mathbb{H}^{n+1}.$$

This implies that, for X defined in (3-12), $dB_{\pm} \cdot X = 0$, since $B_{\pm}(\varphi_s(x, \xi)) = B_{\pm}(x, \xi)$ for all $s \in \mathbb{R}$. Moreover, since $\Phi_{\pm}(\varphi_t(x, \xi)) = e^{\pm t}(x_0 + \xi_0) = e^{\pm t}\Phi_{\pm}(x, \xi)$ from (3-11), we find

$$X\Phi_{\pm} = \pm\Phi_{\pm}. \tag{3-17}$$

For $(x, \nu) \in \mathbb{H}^{n+1} \times \mathbb{S}^n$ (in the hyperboloid model), define the function

$$P(x, \nu) = (x_0 - x' \cdot \nu)^{-1} = (\langle x, (1, \nu) \rangle_M)^{-1} \quad \text{if } x = (x_0, x') \in \mathbb{H}^{n+1}. \tag{3-18}$$

Note that $P(x, \nu) > 0$ everywhere, and in the Poincaré ball model \mathbb{B}^{n+1} we have

$$P(\psi^{-1}(y), \nu) = \frac{1 - |y|^2}{|y - \nu|^2}, \quad y \in \mathbb{B}^{n+1}, \tag{3-19}$$

which is the usual Poisson kernel. Here ψ is as defined in (3-2).

For $(x, \nu) \in \mathbb{H}^{n+1} \times \mathbb{S}^n$, there exist unique $\xi_{\pm} \in S_x\mathbb{H}^{n+1}$ such that $B_{\pm}(x, \xi_{\pm}) = \nu$: these are given by

$$\xi_{\pm}(x, \nu) = \mp x \pm P(x, \nu)(1, \nu), \tag{3-20}$$

and we have

$$\Phi_{\pm}(x, \xi_{\pm}(x, \nu)) = P(x, \nu). \tag{3-21}$$

Notice that the equations $B_{\pm}(x, \xi_{\pm}(x, \nu)) = \nu$ imply that B_{\pm} are submersions. The map $\nu \rightarrow \xi_{\pm}(x, \nu)$ is conformal with the standard choice of metrics on \mathbb{S}^n and $S_x\mathbb{H}^{n+1}$; in fact, for $\zeta_1, \zeta_2 \in T_{\nu}\mathbb{S}^n$,

$$\langle \partial_{\nu}\xi_{\pm}(x, \nu) \cdot \zeta_1, \partial_{\nu}\xi_{\pm}(x, \nu) \cdot \zeta_2 \rangle_M = -P(x, \nu)^2 \langle \zeta_1, \zeta_2 \rangle_{\mathbb{R}^{n+1}}. \tag{3-22}$$

Using that $\langle x + \xi, x - \xi \rangle_M = 2$, we see that

$$\Phi_+(x, \xi)\Phi_-(x, \xi)(1 - B_+(x, \xi) \cdot B_-(x, \xi)) = 2. \tag{3-23}$$

One can parametrize $S\mathbb{H}^{n+1}$ by

$$(v_-, v_+, s) = \left(B_-(x, \xi), B_+(x, \xi), \frac{1}{2} \log \frac{\Phi_+(x, \xi)}{\Phi_-(x, \xi)} \right) \in (\mathbb{S}^n \times \mathbb{S}^n)_{\Delta} \times \mathbb{R}, \tag{3-24}$$

where $(\mathbb{S}^n \times \mathbb{S}^n)_\Delta$ is $\mathbb{S}^n \times \mathbb{S}^n$ minus the diagonal. In fact, the geodesic $\gamma(t) = \varphi_t(x, \xi)$ goes from v_- to v_+ in $\bar{\mathbb{B}}^{n+1}$ and $\gamma(-s)$ is the point of γ closest to $e_0 \in \mathbb{H}^{n+1}$ (corresponding to $0 \in \mathbb{B}^{n+1}$). In the parametrization (3-24), the geodesic flow φ_t is simply

$$(v_-, v_+, s) \mapsto (v_-, v_+, s + t).$$

We finally remark that the stable/unstable subspaces of the cotangent bundle, $E_s^*, E_u^* \subset T^*(S\mathbb{H}^{n+1})$, defined in (3-15), are in fact the conormal bundles of the fibers of the maps B_\pm :

$$E_s^*(\rho) = N^*(B_+^{-1}(B_+(\rho))), \quad E_u^*(\rho) = N^*(B_-^{-1}(B_-(\rho))), \quad \rho \in S\mathbb{H}^{n+1}. \tag{3-25}$$

This is equivalent to saying that the fibers of B_+ integrate (that is, are tangent to) the subbundle $E_0 \oplus E_s \subset T(S\mathbb{H}^{n+1})$, while the fibers of B_- integrate the subbundle $E_0 \oplus E_u$. To see the latter statement, say for B_+ , it is enough to note that $dB_+ \cdot X = 0$ and differentiation along vectors in E_s annihilates the function $x + \xi$ and thus the map B_+ ; therefore, the kernel of dB_+ contains $E_0 \oplus E_s$, and this containment is an equality since the dimensions of both spaces are equal to $n + 1$.

3E. Action of G on the conformal infinity. For $\gamma \in G$ and $v \in \mathbb{S}^n$, $\gamma \cdot (1, v)$ is a lightlike vector with positive zeroth component. We can then define $N_\gamma(v) > 0$, $L_\gamma(v) \in \mathbb{S}^n$ by

$$\gamma \cdot (1, v) = N_\gamma(v)(1, L_\gamma(v)). \tag{3-26}$$

The map L_γ gives the action of G on the conformal infinity $\mathbb{S}^n = \partial\bar{\mathbb{B}}^{n+1}$. This action is transitive and the isotropy groups of $\pm e_1 \in \mathbb{S}^n$ are given by

$$H_\pm = \{\gamma \in G \mid \exists s > 0 \ \gamma \cdot (e_0 \pm e_1) = s(e_0 \pm e_1)\}. \tag{3-27}$$

The isotropy groups H_\pm are the connected subgroups of G with the Lie algebras generated by $R_{i+1, j+1}$ for $1 \leq i < j \leq n$, X , and U_i^\pm for $1 \leq i \leq n$. To see that H_\pm are connected, for $n = 1$ we can check directly that every $\gamma \in H_\pm$ can be written as a product $e^{tX} e^{sU_i^\pm}$ for some $t, s \in \mathbb{R}$, and for $n > 1$ we can use the fact that $\mathbb{S}^n \simeq G/H_\pm$ is simply connected and G is connected, and the homotopy long exact sequence of a fibration.

The differentials of N_γ and L_γ (in v) can be written as

$$dN_\gamma(v) \cdot \zeta = \langle dx_0, \gamma \cdot (0, \zeta) \rangle, \quad (0, dL_\gamma(v) \cdot \zeta) = \frac{\gamma \cdot (0, \zeta) - (dN_\gamma(v) \cdot \zeta)(1, L_\gamma(v))}{N_\gamma(v)};$$

here $\zeta \in T_v\mathbb{S}^n$. We see that the map $v \mapsto L_\gamma(v)$ is conformal with respect to the standard metric on \mathbb{S}^n ; in fact, for $\zeta_1, \zeta_2 \in T_v\mathbb{S}^n$,

$$\langle dL_\gamma(v) \cdot \zeta_1, dL_\gamma(v) \cdot \zeta_2 \rangle_{\mathbb{R}^{n+1}} = N_\gamma(v)^{-2} \langle \zeta_1, \zeta_2 \rangle_{\mathbb{R}^{n+1}}.$$

The maps $B_\pm : S\mathbb{H}^{n+1} \rightarrow \mathbb{S}^n$ are equivariant under the action of G :

$$B_\pm(\gamma \cdot (x, \xi)) = L_\gamma(B_\pm(x, \xi)).$$

Moreover, the functions $\Phi_{\pm}(x, \xi)$ and $P(x, \nu)$ enjoy the following properties:

$$\Phi_{\pm}(\gamma.(x, \xi)) = N_{\gamma}(B_{\pm}(x, \xi))\Phi_{\pm}(x, \xi), \quad P(\gamma \cdot x, L_{\gamma}(\nu)) = N_{\gamma}(\nu)P(x, \nu). \tag{3-28}$$

3F. The bundle \mathcal{E} and parallel transport to the conformal infinity. Consider the vector bundle \mathcal{E} over $S\mathbb{H}^{n+1}$ defined as follows:

$$\mathcal{E} = \{(x, \xi, \nu) \in S\mathbb{H}^{n+1} \times T_x\mathbb{H}^{n+1} \mid g_H(\xi, \nu) = 0\},$$

i.e., the fibers $\mathcal{E}(x, \xi)$ consist of all tangent vectors in $T_x\mathbb{H}^{n+1}$ orthogonal to ξ ; equivalently, $\mathcal{E}(x, \xi)$ consists of all vectors in $\mathbb{R}^{1,n+1}$ orthogonal to x and ξ with respect to the Minkowski inner product. Note that G naturally acts on \mathcal{E} , by putting $\gamma.(x, \xi, \nu) := (\gamma \cdot x, \gamma \cdot \xi, \gamma \cdot \nu)$.

The bundle \mathcal{E} is invariant under parallel transport along geodesics. Therefore, one can consider the first-order differential operator

$$\mathcal{X} : C^{\infty}(S\mathbb{H}^{n+1}; \mathcal{E}) \rightarrow C^{\infty}(S\mathbb{H}^{n+1}; \mathcal{E}), \tag{3-29}$$

which is the generator of parallel transport; namely, if ν is a section of \mathcal{E} and $(x, \xi) \in S\mathbb{H}^{n+1}$, then $\mathcal{X}\nu(x, \xi)$ is the covariant derivative at $t = 0$ of the vector field $\nu(t) := \nu(\varphi_t(x, \xi))$ on the geodesic $\varphi_t(x, \xi)$. Note that $\mathcal{E}(\varphi_t(x, \xi))$ is independent of t as a subspace of $\mathbb{R}^{1,n+1}$, and, under this embedding, \mathcal{X} just acts as X on each coordinate of ν in $\mathbb{R}^{1,n+1}$. The operator $\frac{1}{t}\mathcal{X}$ is a symmetric operator with respect to the standard volume form on $S\mathbb{H}^{n+1}$ and the inner product on \mathcal{E} inherited from $T\mathbb{H}^{n+1}$.

We now consider parallel transport of vectors along geodesics going off to infinity. Let $(x, \xi) \in S\mathbb{H}^{n+1}$ and $\nu \in T_x\mathbb{H}^{n+1}$. We let $(x(t), \xi(t)) = \varphi_t(x, \xi)$ be the corresponding geodesic and $\nu(t) \in T_{x(t)}\mathbb{H}^{n+1}$ be the parallel transport of ν along this geodesic. We embed $\nu(t)$ into the unit ball model \mathbb{B}^{n+1} by defining

$$w(t) = d\psi(x(t)) \cdot \nu(t) \in \mathbb{R}^{n+1},$$

where ψ is as defined in (3-2). Then $w(t)$ converges to 0 as $t \rightarrow \pm\infty$, but the limits $\lim_{t \rightarrow \pm\infty} x_0(t)w(t)$ are nonzero for nonzero ν ; we call the transformation mapping ν to these limits the *transport to conformal infinity* as $t \rightarrow \pm\infty$. More precisely, if

$$\nu = c\xi + u, \quad u \in \mathcal{E}(x, \xi),$$

then we calculate

$$\lim_{t \rightarrow \pm\infty} x_0(t)w(t) = \pm cB_{\pm}(x, \xi) + u' - u_0B_{\pm}(x, \xi), \tag{3-30}$$

where $B_{\pm}(x, \xi) \in \mathbb{S}^n$ is as defined in Section 3D. We will in particular use the inverse of the map $\mathcal{E}(x, \xi) \ni u \mapsto u' - u_0B_{\pm}(x, \xi) \in T_{B_{\pm}(x, \xi)}\mathbb{S}^n$: for $(x, \xi) \in S\mathbb{H}^{n+1}$ and $\zeta \in T_{B_{\pm}(x, \xi)}\mathbb{S}^n$, define (see Figure 3(b))

$$\mathcal{A}_{\pm}(x, \xi)\zeta = (0, \zeta) - \langle (0, \zeta), x \rangle_M(x \pm \xi) = \pm \frac{\partial_{\nu}\xi_{\pm}(x, B_{\pm}(x, \xi)) \cdot \zeta}{P(x, B_{\pm}(x, \xi))} \in \mathcal{E}(x, \xi). \tag{3-31}$$

Here ξ_{\pm} is as defined in (3-20). Note that, by (3-22), \mathcal{A}_{\pm} is an isometry:

$$|\mathcal{A}_{\pm}(x, \xi)\zeta|_{g_H} = |\zeta|_{\mathbb{R}^n}, \quad \zeta \in T_{B_{\pm}(x, \xi)}\mathbb{S}^n. \tag{3-32}$$

Also, \mathcal{A}_\pm is equivariant under the action of G :

$$\mathcal{A}_\pm(\gamma \cdot x, \gamma \cdot \xi) \cdot dL_\gamma(B_\pm(x, \xi)) \cdot \zeta = N_\gamma(B_\pm(x, \xi))^{-1} \gamma \cdot (\mathcal{A}_\pm(x, \xi)\zeta). \tag{3-33}$$

We now write the limits (3-30) in terms of the 0-tangent bundle of Mazzeo and Melrose [1987]. Consider the boundary defining function $\rho_0 := 2(1 - |y|)/(1 + |y|)$ on $\bar{\mathbb{B}}^{n+1}$; note that in the hyperboloid model, with the map ψ defined in (3-2),

$$\rho_0(\psi(x)) = 2 \frac{\sqrt{x_0 + 1} - \sqrt{x_0 - 1}}{\sqrt{x_0 + 1} + \sqrt{x_0 - 1}} = x_0^{-1} + \mathcal{O}(x_0^{-2}) \quad \text{as } x_0 \rightarrow \infty. \tag{3-34}$$

The hyperbolic metric can be written near the boundary as $g_H = (d\rho_0^2 + h_{\rho_0})/\rho_0^2$ with h_{ρ_0} a smooth family of metrics on \mathbb{S}^n , and $h_0 = d\theta^2$ is the canonical metric on the sphere (with curvature 1).

Define the 0-tangent bundle ${}^0T\bar{\mathbb{B}}^{n+1}$ to be the smooth bundle over $\bar{\mathbb{B}}^{n+1}$ whose smooth sections are the elements of the Lie algebra $\mathcal{V}_0(\bar{\mathbb{B}}^{n+1})$ of smooth vectors fields vanishing at $\mathbb{S}^n = \bar{\mathbb{B}}^{n+1} \cap \{\rho_0 = 0\}$; near the boundary, this algebra is locally spanned over $C^\infty(\bar{\mathbb{B}}^{n+1})$ by the vector fields $\rho_0\partial_{\rho_0}, \rho_0\partial_{\theta_1}, \dots, \rho_0\partial_{\theta_n}$ if θ_i are local coordinates on \mathbb{S}^n . There is a natural map ${}^0T\bar{\mathbb{B}}^{n+1} \rightarrow T\bar{\mathbb{B}}^{n+1}$, which is an isomorphism when restricted to the interior \mathbb{B}^{n+1} . We denote by ${}^0T^*\bar{\mathbb{B}}^{n+1}$ the dual bundle to ${}^0T\bar{\mathbb{B}}^{n+1}$, generated locally near $\rho_0 = 0$ by the covectors $d\rho_0/\rho_0, d\theta_1/\rho_0, \dots, d\theta_n/\rho_0$. Note that $T^*\bar{\mathbb{B}}^{n+1}$ naturally embeds into ${}^0T^*\bar{\mathbb{B}}^{n+1}$ and this embedding is an isomorphism in the interior. The metric g_H is a smooth, nondegenerate, positive definite quadratic form on ${}^0T\bar{\mathbb{B}}^{n+1}$, that is, $g_H \in C^\infty(\bar{\mathbb{B}}^{n+1}; \otimes_S^2({}^0T^*\bar{\mathbb{B}}^{n+1}))$, where \otimes_S^2 denotes the space of symmetric 2-tensors. We refer the reader to [Mazzeo and Melrose 1987] for further details (in particular, for an explanation of why 0-bundles are smooth vector bundles); see also [Melrose 1993, §2.2] for the similar b -setting.

We can then interpret (3-30) as follows: for each $(y, \eta) \in S\mathbb{B}^{n+1}$ and each $w \in T_y\mathbb{B}^{n+1}$, the parallel transport $w(t)$ of w along the geodesic $\varphi_t(y, \eta)$ (this geodesic extends smoothly to a curve on \mathbb{B}^{n+1} , as it is part of a line or a circle) has limits as $t \rightarrow \pm\infty$ in the 0-tangent bundle ${}^0T\bar{\mathbb{B}}^{n+1}$. In fact (see [Guillarmou et al. 2010, Appendix A]), the parallel transport

$$\tau(y', y) : {}^0T_y\mathbb{B}^{n+1} \rightarrow {}^0T_{y'}\mathbb{B}^{n+1}$$

from y to $y' \in \mathbb{B}^{n+1}$ along the geodesic starting at y and ending at y' extends smoothly to the boundary $(y, y') \in \bar{\mathbb{B}}^{n+1} \times \bar{\mathbb{B}}^{n+1} \setminus \text{diag}(\mathbb{S}^n \times \mathbb{S}^n)$ as an endomorphism ${}^0T_y\bar{\mathbb{B}}^{n+1} \rightarrow {}^0T_{y'}\bar{\mathbb{B}}^{n+1}$, where $\text{diag}(\mathbb{S}^n \times \mathbb{S}^n)$ denotes the diagonal in the boundary; this parallel transport is an isometry with respect to g_H . The same properties hold for parallel transport of covectors in ${}^0T^*\bar{\mathbb{B}}^{n+1}$, using the duality provided by the metric g_H . An explicit relation to the maps \mathcal{A}_\pm is given by the following formula:

$$\mathcal{A}_\pm(x, \xi) \cdot \zeta = d\psi(x)^{-1} \cdot \tau(\psi(x), B_\pm(x, \xi)) \cdot (\rho_0\zeta), \tag{3-35}$$

where $\rho_0\zeta \in {}^0T_{B_\pm(x, \xi)}\bar{\mathbb{B}}^{n+1}$ is tangent to the conformal boundary \mathbb{S}^n .

4. Horocyclic operators

In this section, we build on the results of Section 3 to construct horocyclic operators

$$U_{\pm} : \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes^j \mathcal{E}^*) \rightarrow \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes^{j+1} \mathcal{E}^*).$$

4A. Symmetric tensors. In this subsection, we assume that E is a vector space of finite dimension N , equipped with an inner product g_E , and let E^* denote the dual space, which has a scalar product induced by g_E (also denoted g_E). (In what follows, we shall take either $E = \mathcal{E}(x, \xi)$ or $E = T_x \mathbb{H}^{n+1}$ for some $(x, \xi) \in S\mathbb{H}^{n+1}$, and the scalar product g_E in both cases is given by the hyperbolic metric g_H on those vector spaces.) We will work with tensor powers of E^* , but the constructions also apply to tensor powers of E by swapping E with E^* .

We introduce some notation for finite sequences to simplify the calculations below. Denote by \mathcal{A}^m the space of all sequences $K = k_1 \dots k_m$ with $1 \leq k_\ell \leq N$. For $k_1 \dots k_m \in \mathcal{A}^m$, $j_1 \dots j_r \in \mathcal{A}^r$, and a sequence of distinct numbers $1 \leq \ell_1, \dots, \ell_r \leq m$, denote by

$$\{\ell_1 \rightarrow j_1, \dots, \ell_r \rightarrow j_r\} K \in \mathcal{A}^m$$

the result of replacing the ℓ_p -th element of K by j_p for all p . We can also replace some j_p by blank space, which means that the corresponding indices are removed from K .

For $m \geq 0$ denote by $\otimes^m E^*$ the m -th tensor power of E^* and by $\otimes_S^m E^*$ the subset of those tensors which are symmetric, i.e., $u \in \otimes_S^m E^*$ if $u(v_{\sigma(1)}, \dots, v_{\sigma(m)}) = u(v_1, \dots, v_m)$ for all $\sigma \in \Pi_m$ and all $v_1, \dots, v_m \in E$, where Π_m is the permutation group of $\{1, \dots, m\}$. There is a natural linear projection $\mathcal{S} : \otimes^m E^* \rightarrow \otimes_S^m E^*$ defined by

$$\mathcal{S}(\eta_1^* \otimes \dots \otimes \eta_m^*) = \frac{1}{m!} \sum_{\sigma \in \Pi_m} \eta_{\sigma(1)}^* \otimes \dots \otimes \eta_{\sigma(m)}^*, \quad \eta_k^* \in E^*. \tag{4-1}$$

The metric g_E induces a scalar product on $\otimes^m E^*$,

$$\langle v_1^* \otimes \dots \otimes v_m^*, w_1^* \otimes \dots \otimes w_m^* \rangle_{g_E} = \prod_{j=1}^m \langle v_j^*, w_j^* \rangle_{g_E}, \quad v_i^*, w_i^* \in E^*.$$

The operator \mathcal{S} is selfadjoint and thus an orthogonal projection with respect to this scalar product.

Using the metric g_E , one can decompose the vector space $\otimes_S^m E^*$ as follows. Let $(e_i)_{i=1}^N$ be an orthonormal basis of E for the metric g_E and (e_i^*) be the dual basis. First of all, introduce the trace map $\mathcal{T} : \otimes^m E^* \rightarrow \otimes^{m-2} E^*$ contracting the first two indices by the metric: for $v_i \in E$, define

$$\mathcal{T}(u)(v_1, \dots, v_{m-2}) := \sum_{i=1}^N u(e_i, e_i, v_1, \dots, v_{m-2}) \tag{4-2}$$

(the result is independent of the choice of the basis). For $m < 2$, we define \mathcal{T} to be zero on $\otimes^m E^*$. Note that \mathcal{T} maps $\otimes_S^{m+2} E^*$ onto $\otimes_S^m E^*$. Set

$$e_K^* := e_{k_1}^* \otimes \dots \otimes e_{k_m}^* \in \otimes^m E^*, \quad K = k_1 \dots k_m \in \mathcal{A}^m.$$

Then

$$\mathcal{T}\left(\sum_{K \in \mathcal{A}^{m+2}} f_K e_K^*\right) = \sum_{K \in \mathcal{A}^m} \sum_{q \in \mathcal{A}} f_{qqK} e_K^*.$$

The adjoint of $\mathcal{T} : \otimes_S^{m+2} E^* \rightarrow \otimes_S^m E^*$ with respect to the scalar product g_E is given by the map $u \mapsto \mathcal{S}(g_E \otimes u)$. To simplify computations, we define a scaled version of it: let $\mathcal{I} : \otimes_S^m E^* \rightarrow \otimes_S^{m+2} E^*$ be defined by

$$\mathcal{I}(u) = \frac{(m+2)(m+1)}{2} \mathcal{S}(g_E \otimes u) = \frac{(m+2)(m+1)}{2} \mathcal{T}^*(u). \tag{4-3}$$

Then

$$\mathcal{I}\left(\sum_{K \in \mathcal{A}^m} f_K e_K^*\right) = \sum_{K \in \mathcal{A}^{m+2}} \sum_{\substack{\ell, r=1 \\ \ell < r}}^{m+2} \delta_{k_\ell k_r} f_{\{\ell \rightarrow, r \rightarrow\}K} e_K^*.$$

Note that, for $u \in \otimes_S^m E^*$,

$$\mathcal{T}(\mathcal{I}u) = (2m + N)u + \mathcal{I}(\mathcal{T}u). \tag{4-4}$$

By (4-3) and (4-4), the homomorphism $\mathcal{T}\mathcal{I} : \otimes_S^m E^* \rightarrow \otimes_S^m E^*$ is positive definite and thus an isomorphism. Therefore, for $u \in \otimes_S^m E^*$, we can decompose $u = u_1 + \mathcal{I}(u_2)$, where $u_1 \in \otimes_S^m E^*$ satisfies $\mathcal{T}(u_1) = 0$ and $u_2 = (\mathcal{T}\mathcal{I})^{-1}\mathcal{T}u \in \otimes_S^{m-2} E^*$. Iterating this process, we can decompose any $u \in \otimes_S^m E^*$ into

$$u = \sum_{r=0}^{\lfloor m/2 \rfloor} \mathcal{I}^r(u_r), \quad u_r \in \otimes_S^{m-2r} E^*, \quad \mathcal{T}(u_r) = 0, \tag{4-5}$$

with u_r determined uniquely by u .

Another operation on tensors which will be used is the interior product: if $v \in E$ and $u \in \otimes_S^m E^*$, we denote by $\iota_v(u) \in \otimes_S^{m-1} E^*$ the interior product of u by v given by

$$\iota_v u(v_1, \dots, v_{m-1}) := u(v, v_1, \dots, v_{m-1}).$$

If $v^* \in E^*$, we write $\iota_{v^*} u$ for the tensor $\iota_v u$ with $g_E(v, \cdot) = v^*$.

We conclude this subsection with a correspondence which will be useful in certain calculations later. There is a linear isomorphism between $\otimes_S^m E^*$ and the space $\text{Pol}^m(E)$ of homogeneous polynomials of degree m on E : to a tensor $u \in \otimes_S^m E^*$ we associate the function on E given by $x \rightarrow P_u(x) := u(x, \dots, x)$. If we write $x = \sum_{i=1}^N x_i e_i$ in a given orthonormal basis, then

$$P_{\mathcal{S}(e_K^*)}(x) = \prod_{j=1}^m x_{k_j}, \quad K = k_1 \cdots k_m \in \mathcal{A}^m.$$

The flat Laplacian associated to g_E is given by $\Delta_E = -\sum_{i=1}^N \partial_{x_i}^2$ in the coordinates induced by the basis (e_i) . Then it is direct to see that

$$\Delta_E P_u(x) = -m(m-1)P_{\mathcal{T}(u)}(x), \quad u \in \otimes_S^m E^*, \tag{4-6}$$

which means that the trace corresponds to applying the Laplacian (see [Dairbekov and Sharafutdinov 2010, Lemma 2.4]). In particular, trace-free symmetric tensors of order m correspond to homogeneous harmonic polynomials, and thus restrict to spherical harmonics on the sphere $|x|_{g_E} = 1$ of E . We also have

$$P_{\mathcal{I}(u)}(x) = \frac{1}{2}(m+2)(m+1)|x|^2 P_u(x), \quad u \in \otimes_S^m E^*. \tag{4-7}$$

4B. Horocyclic operators. We now consider the left-invariant vector fields $X, U_{\pm}^i, R_{i+1,j+1}$ on the isometry group G , identified with the elements of the Lie algebra of G introduced in (3-6), (3-7). Recall that G acts on $S\mathbb{H}^{n+1}$ transitively with the isotropy group $H \simeq \text{SO}(n)$ and this action gives rise to the projection $\pi_S : G \rightarrow S\mathbb{H}^{n+1}$; see (3-10). Note that, with the maps $\Phi_{\pm} : S\mathbb{H}^{n+1} \rightarrow \mathbb{R}^+$ and $B_{\pm} : S\mathbb{H}^{n+1} \rightarrow \mathbb{S}^n$ defined in (3-16), we have

$$B_{\pm}(\pi_S(\gamma)) = L_{\gamma}(\pm e_1) \quad \text{and} \quad \Phi_{\pm}(\pi_S(\gamma)) = N_{\gamma}(\pm e_1), \quad \gamma \in G,$$

where $N_{\gamma} : \mathbb{S}^n \rightarrow \mathbb{R}^+$ and $L_{\gamma} : \mathbb{S}^n \rightarrow \mathbb{S}^n$ are defined in (3-26). Since H_{\pm} , the isotropy group of $\pm e_1$ under the action L_{γ} , contains X and U_i^{\pm} in its Lie algebra (see (3-27) and Figure 3(a)), we find

$$d(B_{\pm} \circ \pi_S) \cdot U_i^{\pm} = 0 \quad \text{and} \quad d(B_{\pm} \circ \pi_S) \cdot X = 0. \tag{4-8}$$

We also calculate

$$d(\Phi_{\pm} \circ \pi_S) \cdot U_i^{\pm} = 0. \tag{4-9}$$

Define the differential operator on G

$$U_K^{\pm} := U_{k_1}^{\pm} \cdots U_{k_m}^{\pm}, \quad K = k_1 \cdots k_m \in \mathcal{A}^m.$$

Note that the order in which k_1, \dots, k_m are listed does not matter, by (3-8). Moreover, by (3-8),

$$[R_{i+1,j+1}, U_K^{\pm}] = \sum_{\ell=1}^m (\delta_{jk_{\ell}} U_{\{\ell \rightarrow i\}K}^{\pm} - \delta_{ik_{\ell}} U_{\{\ell \rightarrow j\}K}^{\pm}). \tag{4-10}$$

Since H is generated by the vector fields $R_{i+1,j+1}$, we see that in dimensions $n+1 > 2$ the horocyclic vector fields U_i^{\pm} , and more generally the operators U_K^{\pm} , are not invariant under right multiplication by elements of H and therefore do not descend to differential operators on $S\mathbb{H}^{n+1}$ —in other words, if $u \in \mathcal{D}'(S\mathbb{H}^{n+1})$, then $U_K^{\pm}(\pi_S^* u) \in \mathcal{D}'(G)$ is not in the image of π_S^* .

However, in this section we will show how to differentiate distributions on $S\mathbb{H}^{n+1}$ along the horocyclic vector fields, resulting in sections of the vector bundle \mathcal{E} introduced in Section 3F and its tensor powers. First of all, we note that by (3-14), the stable and unstable bundles $E_s(x, \xi)$ and $E_u(x, \xi)$ are canonically isomorphic to $\mathcal{E}(x, \xi)$, by the maps

$$\theta_+ : \mathcal{E}(x, \xi) \rightarrow E_s(x, \xi), \quad \theta_- : \mathcal{E}(x, \xi) \rightarrow E_u(x, \xi), \quad \theta_{\pm}(v) = (-v, \pm v).$$

For $u \in \mathcal{D}'(S\mathbb{H}^{n+1})$, we then define the horocyclic derivatives $\mathcal{U}_{\pm} u \in \mathcal{D}'(S\mathbb{H}^{n+1}; \mathcal{E}^*)$ by restricting the differential $du \in \mathcal{D}'(S\mathbb{H}^{n+1}; T^*(S\mathbb{H}^{n+1}))$ to the stable/unstable foliations and pulling it back by θ_{\pm} :

$$\mathcal{U}_{\pm} u(x, \xi) := du(x, \xi) \circ \theta_{\pm} \in \mathcal{E}^*(x, \xi). \tag{4-11}$$

To relate \mathcal{U}_\pm to the vector fields U_i^\pm on the group G , consider the orthonormal frame e_1^*, \dots, e_n^* of the bundle $\pi_S^* \mathcal{E}^*$ over G defined by

$$e_j^*(\gamma) := \gamma^{-*}(e_{j+1}^*) \in \mathcal{E}^*(\pi_S(\gamma)),$$

where the $e_j^* = dx_j$ form the dual basis to the canonical basis $(e_j)_{j=0, \dots, n+1}$ of $\mathbb{R}^{1, n+1}$, and $\gamma^{-*} = (\gamma^{-1})^* : (\mathbb{R}^{1, n+1})^* \rightarrow (\mathbb{R}^{1, n+1})^*$. More generally, we can define the orthonormal frame e_K^* of $\pi_S^*(\otimes^m \mathcal{E}^*)$ by

$$e_K^* := e_{k_1}^* \otimes \dots \otimes e_{k_m}^*, \quad K = k_1 \dots k_m \in \mathcal{A}^m.$$

We compute, for $u \in \mathcal{D}'(S\mathbb{H}^{n+1})$, $du(\pi_S(\gamma)) \cdot \theta_\pm(\gamma(e_{j+1})) = U_j^\pm(\pi_S^* u)(\gamma)$, and thus

$$\pi_S^*(\mathcal{U}_\pm u) = \sum_{j=1}^n U_j^\pm(\pi_S^* u) e_j^*. \tag{4-12}$$

We next use the formula (4-12) to define \mathcal{U}_\pm as an operator

$$\mathcal{U}_\pm : \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes^m \mathcal{E}^*) \rightarrow \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes^{m+1} \mathcal{E}^*) \tag{4-13}$$

as follows: for $u \in \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes^m \mathcal{E}^*)$, define $\mathcal{U}_\pm u$ by

$$\pi_S^*(\mathcal{U}_\pm u) = \sum_{r=1}^n \sum_{K \in \mathcal{A}^m} (U_r^\pm u_K) e_{rK}^*, \quad \pi_S^* u = \sum_{K \in \mathcal{A}^m} u_K e_K^*. \tag{4-14}$$

This definition makes sense (that is, the right-hand side of the first formula in (4-14) lies in the image of π_S^*) since a section

$$f = \sum_{K \in \mathcal{A}^m} f_K e_K^* \in \mathcal{D}'(G; \pi_S^*(\otimes^m \mathcal{E}^*)), \quad f_K \in \mathcal{D}'(G),$$

lies in the image of π_S^* if and only if $R_{i+1, j+1} f = 0$ for $1 \leq i < j \leq n$ (the differentiation is well defined since the fibers of $\pi_S^*(\otimes^m \mathcal{E}^*)$ are the same along each integral curve of $R_{i+1, j+1}$), and this translates to

$$R_{i+1, j+1} f_K = \sum_{\ell=1}^m (\delta_{jk_\ell} f_{\{\ell \rightarrow i\}K} - \delta_{ik_\ell} f_{\{\ell \rightarrow j\}K}), \quad 1 \leq i < j \leq n, \quad K \in \mathcal{A}^m; \tag{4-15}$$

to show (4-15) for $f_{rK} = U_r^\pm u_K$, we use (3-8):

$$\begin{aligned} R_{i+1, j+1} f_{rK} &= [R_{i+1, j+1}, U_r^\pm] u_K + U_r^\pm R_{i+1, j+1} u_K \\ &= \delta_{jr} U_i^\pm u_K - \delta_{ir} U_j^\pm u_K + \sum_{\ell=1}^m \delta_{jk_\ell} U_r^\pm u_{\{\ell \rightarrow i\}K} - \delta_{ik_\ell} U_r^\pm u_{\{\ell \rightarrow j\}K}. \end{aligned}$$

To interpret the operator (4-13) in terms of the stable/unstable foliations in a manner similar to (4-11), consider the connection ∇^S on the bundle \mathcal{E} over $S\mathbb{H}^{n+1}$ defined as follows: for $(x, \xi) \in S\mathbb{H}^{n+1}$, $(v, w) \in T_{(x, \xi)}(S\mathbb{H}^{n+1})$, and $u \in \mathcal{D}'(S\mathbb{H}^{n+1}; \mathcal{E})$, let $\nabla_{(v, w)}^S u(x, \xi)$ be the orthogonal projection of

$$\nabla_{(v, w)}^{\mathbb{R}^{1, n+1}} u(x, \xi) \quad \text{onto} \quad \mathcal{E}(x, \xi) \subset \mathbb{R}^{1, n+1},$$

where $\nabla^{\mathbb{R}^{1,n+1}}$ is the canonical connection on the trivial bundle $S\mathbb{H}^{n+1} \times \mathbb{R}^{1,n+1}$ over $S\mathbb{H}^{n+1}$ (corresponding to differentiating the coordinates of u in $\mathbb{R}^{1,n+1}$). Then ∇^S naturally induces a connection on $\otimes^m \mathcal{E}^*$, also denoted ∇^S , and we have, for $v, v_1, \dots, v_m \in \mathcal{E}(x, \xi)$ and $u \in \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes^m \mathcal{E}^*)$,

$$\mathcal{U}_\pm u(x, \xi)(v, v_1, \dots, v_m) = (\nabla_{\theta_\pm(v)}^S u)(v_1, \dots, v_m). \tag{4-16}$$

Indeed, if $\gamma(t) = \gamma(0)e^{tU_j^\pm}$ is an integral curve of U_j^\pm on G , then $\gamma(t)e_2, \dots, \gamma(t)e_{n+1}$ form a parallel frame of \mathcal{E} over the curve $(x(t), \xi(t)) = \pi_S(\gamma(t))$ with respect to ∇^S , since the covariant derivative of $\gamma(t)e_k$ in t with respect to $\nabla^{\mathbb{R}^{1,n+1}}$ is simply $\gamma(t)U_j^\pm e_k$; by (3-7) this is a linear combination of $x(t) = \gamma(t)e_0$ and $\xi(t) = \gamma(t)e_1$ and thus $\nabla_t^S(\gamma(t)e_k) = 0$.

Note also that the operator \mathcal{X} defined in (3-29) can be interpreted as the covariant derivative on \mathcal{E} along the generator X of the geodesic flow by the connection ∇^S . One can naturally generalize \mathcal{X} to a first-order differential operator

$$\mathcal{X} : \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes^m \mathcal{E}^*) \rightarrow \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes^m \mathcal{E}^*), \tag{4-17}$$

and $\frac{1}{i}\mathcal{X}$ is still symmetric with respect to the natural measure on $S\mathbb{H}^{n+1}$ and the inner product on $\otimes^m \mathcal{E}^*$ induced by the Minkowski metric. A characterization of X in terms of the frame e_K^* is given by

$$\pi_S^*(\mathcal{X}u) = \sum_{K \in \mathcal{A}^m} (Xu_K) e_K^*, \quad \pi_S^*u = \sum_{K \in \mathcal{A}^m} u_K e_K^*. \tag{4-18}$$

It follows from (3-8) that, for $u \in \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes^m \mathcal{E}^*)$,

$$\mathcal{X}\mathcal{U}_\pm u - \mathcal{U}_\pm \mathcal{X}u = \pm \mathcal{U}_\pm u. \tag{4-19}$$

We also observe that, since $[U_i^\pm, U_j^\pm] = 0$, for each scalar distribution $u \in \mathcal{D}'(S\mathbb{H}^{n+1})$ and $m \in \mathbb{N}$ we have $\mathcal{U}_\pm^m u \in \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes_S^m \mathcal{E}^*)$, where $\otimes_S^m \mathcal{E}^* \subset \otimes^m \mathcal{E}^*$ denotes the space of all symmetric cotensors of order m . Inversion of the operator \mathcal{U}_\pm^m is the topic of the next subsection. We conclude with the following lemma, describing how the operator \mathcal{U}_\pm^m acts on distributions invariant under the left action of an element of G :

Lemma 4.1. *Let $\gamma \in G$ and $u \in \mathcal{D}'(S\mathbb{H}^{n+1})$. Assume also that u is invariant under left multiplications by γ , namely $u(\gamma.(x, \xi)) = u(x, \xi)$ for all¹ $(x, \xi) \in S\mathbb{H}^{n+1}$. Then $v = \mathcal{U}_\pm^m u$ is equivariant under left multiplication by γ in the following sense:*

$$v(\gamma.(x, \xi)) = \gamma.v(x, \xi), \tag{4-20}$$

where the action of γ on $\otimes_S^m \mathcal{E}^*$ is naturally induced by its action on \mathcal{E} (by taking inverse transposes), which in turn comes from the action of γ on $\mathbb{R}^{1,n+1}$.

Proof. We have, for $\gamma' \in G$,

$$\mathcal{U}_\pm^m u(\pi_S(\gamma')) = \sum_{K \in \mathcal{A}^m} (U_K^\pm(u \circ \pi_S)(\gamma')) e_K^*(\gamma').$$

¹Strictly speaking, this statement should be formulated in terms of the pullback of the distribution u by the map $(x, \xi) \mapsto \gamma.(x, \xi)$.

Therefore, since U_j^\pm are left-invariant vector fields on G ,

$$\mathcal{U}_\pm^m u(\gamma.\pi_S(\gamma')) = \mathcal{U}_\pm^m u(\pi_S(\gamma\gamma')) = \sum_{K \in \mathcal{A}^m} (U_K^\pm(u \circ \pi_S)(\gamma')) \mathbf{e}_K^*(\gamma\gamma').$$

It remains to note that $\mathbf{e}_K^*(\gamma\gamma') = \gamma.\mathbf{e}_K^*(\gamma')$. □

4C. Inverting horocyclic operators. In this subsection, we show that distributions $v \in \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes_S^m \mathcal{E}^*)$ satisfying certain conditions are in fact in the image of \mathcal{U}_\pm^m acting on $\mathcal{D}'(S\mathbb{H}^{n+1})$. This is an important step in our construction of Pollicott–Ruelle resonances, as it will make it possible to recover a scalar resonant state corresponding to a resonance in the m -th band. More precisely, we prove:

Lemma 4.2. *Assume that $v \in \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes_S^m \mathcal{E}^*)$ satisfies $\mathcal{U}_\pm v = 0$, and $\mathcal{X}v = \pm\lambda v$ for $\lambda \notin \frac{1}{2}\mathbb{Z}$. Then there exists $u \in \mathcal{D}'(S\mathbb{H}^{n+1})$ such that $\mathcal{U}_\pm^m u = v$ and $Xu = \pm(\lambda - m)u$. Moreover, if v is equivariant under left multiplication by some $\gamma \in G$ in the sense of (4-20), then u is invariant under left multiplication by γ .*

The proof of Lemma 4.2 is modeled on the following well-known formula recovering a homogeneous polynomial of degree m from its coefficients: given constants a_α for each multiindex α of length m , we have

$$\partial_x^\beta \sum_{|\alpha|=m} \frac{1}{\alpha!} x^\alpha a_\alpha = a_\beta, \quad |\beta| = m. \tag{4-21}$$

The formula recovering u from v in Lemma 4.2 is morally similar to (4-21), with U_j^\pm taking the role of ∂_{x_j} , the condition $\mathcal{U}_\pm v = 0$ corresponding to a_α being constants, and U_j^\mp taking the role of the multiplication operators x_j . However, the commutation structure of U_j^\pm , given by (3-8), is more involved than that of ∂_{x_j} and x_j , and in particular it involves the vector field X , explaining the need for the condition $\mathcal{X}v = \pm\lambda v$ (which is satisfied by resonant states).

To prove Lemma 4.2, we define the operator

$$\mathcal{V}_\pm : \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes^{m+1} \mathcal{E}^*) \rightarrow \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes^m \mathcal{E}^*), \quad \mathcal{V}_\pm := \mathcal{T}\mathcal{U}_\pm,$$

where \mathcal{T} is as defined in Section 4A. Then, by (4-14),

$$\pi_S^*(\mathcal{V}_\pm u) = \sum_{K \in \mathcal{A}^m} \sum_{q \in \mathcal{A}} (U_q^\pm u_{qK}) \mathbf{e}_K^*, \quad u = \sum_{K \in \mathcal{A}^{m+1}} u_K \mathbf{e}_K^*.$$

For later use, we record the following fact:

Lemma 4.3. $\mathcal{U}_\pm^* = -\mathcal{V}_\pm$, where the adjoint is understood in the formal sense.

Proof. If $u \in C_0^\infty(S\mathbb{H}^{n+1}; \otimes^m \mathcal{E}^*)$, $v \in C^\infty(S\mathbb{H}^{n+1}; \otimes^{m+1} \mathcal{E}^*)$, and u_K, v_J are the coordinates of $\pi_S^* u$ and $\pi_S^* v$ in the bases $(\mathbf{e}_K^*)_{K \in \mathcal{A}^m}$ and $(\mathbf{e}_J^*)_{J \in \mathcal{A}^{m+1}}$, then, by (4-14), we compute the following pointwise identity on $S\mathbb{H}^{n+1}$:

$$\langle \mathcal{U}_\pm u, \bar{v} \rangle + \langle u, \overline{\mathcal{V}_\pm v} \rangle = \mathcal{V}_\pm w, \quad w \in C_0^\infty(S\mathbb{H}^{n+1}; \mathcal{E}^*), \quad \pi_S^* w = \sum_{\substack{K \in \mathcal{A}^m \\ q \in \mathcal{A}}} u_K \overline{v_{qK}} \mathbf{e}_q^*.$$

It remains to show that, for each w , the integral of $\mathcal{V}_\pm w$ is equal to zero. Since \mathcal{V}_\pm is a differential operator of order 1, we must have

$$\int_{S\mathbb{H}^{n+1}} \mathcal{V}_\pm w = \int_{S\mathbb{H}^{n+1}} \langle w, \eta_\pm \rangle$$

for all w and some $\eta_\pm \in C^\infty(S\mathbb{H}^{n+1}; \mathcal{E}^*)$ independent of w . Then η_\pm is equivariant under the action of the isometry group G and, in particular, $|\eta_\pm|$ is a constant function on $S\mathbb{H}^{n+1}$. Moreover, using that $\int Xf = 0$ for all $f \in C_0^\infty(S\mathbb{H}^{n+1})$ and $\mathcal{V}_\pm(\mathcal{X}w) = (X \mp 1)\mathcal{V}_\pm w$, we get, for all $w \in C_0^\infty$,

$$\mp \int_{S\mathbb{H}^{n+1}} \langle w, \eta_\pm \rangle = \int_{S\mathbb{H}^{n+1}} \mathcal{V}_\pm(\mathcal{X}w) = - \int_{S\mathbb{H}^{n+1}} \langle w, \mathcal{X}\eta_\pm \rangle.$$

This implies that $\mathcal{X}\eta_\pm = \pm\eta_\pm$ and, in particular,

$$X|\eta_\pm|^2 = 2\langle \mathcal{X}\eta_\pm, \eta_\pm \rangle = \pm 2|\eta_\pm|^2.$$

Since $|\eta_\pm|^2$ is a constant function, this implies $\eta_\pm = 0$, finishing the proof. \square

To construct u from v in Lemma 4.2, we first handle the case when $\mathcal{T}(v) = 0$; this condition is automatically satisfied when $m \leq 1$.

Lemma 4.4. *Assume that $v \in \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes_S^m \mathcal{E}^*)$ and $\mathcal{U}_\pm v = 0, \mathcal{T}(v) = 0$. Define $u = \mathcal{V}_\mp^m v \in \mathcal{D}'(S\mathbb{H}^{n+1})$. Then*

$$\mathcal{U}_\pm^m u = 2^m m! \left(\prod_{\ell=n-1}^{n+m-2} (\ell \pm \mathcal{X}) \right) v. \tag{4-22}$$

Proof. Assume that

$$\pi_S^* v = \sum_{K \in \mathcal{A}^m} f_K \mathbf{e}_K^*, \quad f_K \in \mathcal{D}'(G).$$

Then

$$\pi_S^* u = \sum_{K \in \mathcal{A}^m} U_K^\mp f_K, \quad \pi_S^*(\mathcal{U}_\pm^m u) = \sum_{K, J \in \mathcal{A}^m} U_J^\pm U_K^\mp f_K \mathbf{e}_J^*.$$

For $0 \leq r < m, J \in \mathcal{A}^{m-1-r}$, and $p \in \mathcal{A}$, we have, by (3-8),

$$\sum_{\substack{K \in \mathcal{A}^r \\ q \in \mathcal{A}}} [U_p^\pm, U_q^\mp] U_K^\mp f_{qKJ} = \pm 2X \sum_{K \in \mathcal{A}^r} U_K^\mp f_{pKJ} + 2 \sum_{\substack{K \in \mathcal{A}^r \\ q \in \mathcal{A}}} R_{p+1, q+1} U_K^\mp f_{qKJ}.$$

To compute the second term on the right-hand side, we commute $R_{p+1, q+1}$ with U_K^\mp by (4-10) and use (4-15) to get

$$\begin{aligned} \sum_{\substack{K \in \mathcal{A}^r \\ q \in \mathcal{A}}} R_{p+1, q+1} U_K^\mp f_{qKJ} &= \sum_{\substack{K \in \mathcal{A}^r \\ q \in \mathcal{A}}} \left(\sum_{\ell=1}^r (\delta_{qk_\ell} U_{\{\ell \rightarrow p\}K}^\mp f_{qKJ} - \delta_{pk_\ell} U_{\{\ell \rightarrow q\}K}^\mp f_{qKJ}) + U_K^\mp f_{pKJ} - \delta_{pq} U_K^\mp f_{qKJ} \right. \\ &\quad + \sum_{\ell=1}^r (\delta_{qk_\ell} U_K^\mp f_{q(\{\ell \rightarrow p\}K)J} - \delta_{pk_\ell} U_K^\mp f_{q(\{\ell \rightarrow q\}K)J}) \\ &\quad \left. + \sum_{\ell=1}^{m-1-r} (\delta_{qj_\ell} U_K^\mp f_{qK(\{\ell \rightarrow p\}J)} - \delta_{pj_\ell} U_K^\mp f_{qK(\{\ell \rightarrow q\}J)}) \right). \end{aligned}$$

Since v is symmetric and $\mathcal{T}(v) = 0$, the expressions $\sum_{K \in \mathcal{A}^r, q \in \mathcal{A}} \delta_{qk_\ell} U_{\{\ell \rightarrow p\}K}^\mp f_{qKJ}$, $\sum_{q \in \mathcal{A}} f_{q(\{\ell \rightarrow q\}K)J}$, and $\sum_{q \in \mathcal{A}} f_{qK(\{\ell \rightarrow q\}J)}$ are zero. Further using the symmetry of v , we find

$$\sum_{\substack{K \in \mathcal{A}^r \\ q \in \mathcal{A}}} R_{p+1, q+1} U_K^\mp f_{qKJ} = (n + m - r - 2) \sum_{K \in \mathcal{A}^r} U_K^\mp f_{pKJ},$$

and thus

$$\sum_{\substack{K \in \mathcal{A}^r \\ q \in \mathcal{A}}} [U_p^\pm, U_q^\mp] U_K^\mp f_{qKJ} = 2 \sum_{K \in \mathcal{A}^r} U_K^\mp (\pm X + n + m - 2r - 2) f_{pKJ}. \tag{4-23}$$

Then, using that $\mathcal{U}_\pm v = 0$, we find

$$\begin{aligned} \sum_{\substack{K \in \mathcal{A}^{r+1} \\ q \in \mathcal{A}}} U_p^\pm U_K^\mp f_{qKJ} &= \sum_{\substack{K \in \mathcal{A}^r \\ q \in \mathcal{A}}} \sum_{\ell=1}^{r+1} U_{k_\ell \dots k_r}^\mp [U_p^\pm, U_q^\mp] U_{k_1 \dots k_{\ell-1}}^\mp f_{qKJ} \\ &= 2 \sum_{K \in \mathcal{A}^r} \sum_{\ell=1}^{r+1} U_K^\mp (\pm X + n + m - 2\ell) f_{pKJ} \\ &= 2(r + 1) \sum_{K \in \mathcal{A}^r} U_K^\mp (\pm X + n + m - r - 2) f_{pKJ}. \end{aligned} \tag{4-24}$$

By iterating (4-24) we obtain (using also that v is symmetric), for $J \in \mathcal{A}^m$,

$$\begin{aligned} U_J^\pm \sum_{K \in \mathcal{A}^m} U_K^\mp f_K &= 2m U_{j_1 \dots j_{m-1}}^\pm \sum_{K \in \mathcal{A}^{m-1}} U_K^\mp (\pm X + n - 1) f_{Kj_m} \\ &= 4m(m - 1) U_{j_1 \dots j_{m-2}}^\pm \sum_{K \in \mathcal{A}^{m-2}} U_K^\mp (\pm X + n) (\pm X + n - 1) f_{Kj_{m-1}j_m} \\ &\quad \vdots \\ &= 2^m m! \prod_{\ell=n-1}^{n+m-2} (\pm X + \ell) f_J, \end{aligned}$$

which achieves the proof. □

To handle the case $\mathcal{T}(v) \neq 0$, define also the horocyclic Laplacians

$$\Delta_\pm := -\mathcal{T} \mathcal{U}_\pm^2 = -\mathcal{V}_\pm \mathcal{U}_\pm : \mathcal{D}'(\mathbb{S}\mathbb{H}^{n+1}) \rightarrow \mathcal{D}'(\mathbb{S}\mathbb{H}^{n+1}),$$

so that, for $u \in \mathcal{D}'(\mathbb{S}\mathbb{H}^{n+1})$,

$$\pi_S^* \Delta_\pm u = - \sum_{q=1}^n U_q^\pm U_q^\pm (\pi_S^* u).$$

Note that, by the commutation relation (3-8),

$$[X, \Delta_\pm] = \pm 2 \Delta_\pm. \tag{4-25}$$

Also, by Lemma 4.3, Δ_\pm are symmetric operators.

Lemma 4.5. *Assume that $u \in \mathcal{D}'(S\mathbb{H}^{n+1})$ and $U_{\pm}^{m+1}u = 0$. Then*

$$U_{\pm}^{m+2}\Delta_{\mp}u = -4(\mathcal{X} \mp m)(2\mathcal{X} \pm (n-2))\mathcal{I}(U_{\pm}^m u) - 4\mathcal{I}^2(\mathcal{T}(U_{\pm}^m u)).$$

Proof. We have

$$\pi_S^*(U_{\pm}^{m+2}\Delta_{\mp}u) = - \sum_{\substack{K \in \mathcal{A}^{m+2} \\ q \in \mathcal{A}}} U_K^{\pm} U_q^{\mp} U_q^{\mp} u \mathbf{e}_K^*.$$

Using (3-8), we compute, for $K \in \mathcal{A}^{m+2}$ and $q \in \mathcal{A}$,

$$\begin{aligned} & [U_K^{\pm}, U_q^{\mp}] \\ &= \sum_{\ell=1}^{m+2} U_{k_1 \dots k_{\ell-1}}^{\pm} [U_{k_{\ell}}^{\pm}, U_q^{\mp}] U_{k_{\ell+1} \dots k_{m+2}}^{\pm} \\ &= 2 \sum_{\ell=1}^{m+2} (\delta_{qk_{\ell}} U_{\{\ell \rightarrow\}K}^{\pm} (\pm X + m - \ell + 2) + U_{k_1 \dots k_{\ell-1}}^{\pm} R_{k_{\ell+1}, q+1} U_{k_{\ell+1} \dots k_{m+2}}^{\pm}) \\ &= 2 \sum_{\ell=1}^{m+2} \left(U_{\{\ell \rightarrow\}K}^{\pm} (\delta_{qk_{\ell}} (\pm X + m - \ell + 2) + R_{k_{\ell+1}, q+1}) + \sum_{r=\ell+1}^{m+2} (\delta_{qk_r} U_{\{r \rightarrow\}K}^{\pm} - \delta_{k_{\ell}k_r} U_{\{\ell \rightarrow, r \rightarrow q\}K}^{\pm}) \right) \\ &= 2 \sum_{\ell=1}^{m+2} \left(U_{\{\ell \rightarrow\}K}^{\pm} (\delta_{qk_{\ell}} (\pm X + m + 1) + R_{k_{\ell+1}, q+1}) - \sum_{r=\ell+1}^{m+2} \delta_{k_{\ell}k_r} U_{\{\ell \rightarrow, r \rightarrow q\}K}^{\pm} \right). \end{aligned}$$

Since $U_{\pm}^{m+1}u = 0$, for $K \in \mathcal{A}^{m+2}$ and $q \in \mathcal{A}$ we have $U_K^{\pm}u = [U_K^{\pm}, U_q^{\mp}]u = 0$, and thus

$$U_K^{\pm} U_q^{\mp} U_q^{\mp} u = [[U_K^{\pm}, U_q^{\mp}], U_q^{\mp}]u.$$

We calculate

$$\sum_{q \in \mathcal{A}} [\delta_{qk_{\ell}} (\pm X + m + 1) + R_{k_{\ell+1}, q+1}, U_q^{\mp}] = (n-2)U_{k_{\ell}}^{\mp},$$

and thus, for $K \in \mathcal{A}^{m+2}$,

$$\sum_{q \in \mathcal{A}} U_K^{\pm} U_q^{\mp} U_q^{\mp} u = 2 \sum_{\ell=1}^{m+2} \left([U_{\{\ell \rightarrow\}K}^{\pm}, U_{k_{\ell}}^{\mp}] (\pm X + m + n - 1) - \sum_{r=\ell+1}^{m+2} \delta_{k_{\ell}k_r} \sum_{q \in \mathcal{A}} [U_{\{\ell \rightarrow, r \rightarrow q\}K}^{\pm}, U_q^{\mp}] \right) u.$$

Now, for $K \in \mathcal{A}^{m+2}$,

$$\begin{aligned} & \sum_{\ell=1}^{m+2} [U_{\{\ell \rightarrow\}K}^{\pm}, U_{k_{\ell}}^{\mp}] (\pm X + m + n - 1) u \\ &= 2 \sum_{\substack{\ell, s=1 \\ \ell \neq s}}^{m+2} \left(\delta_{k_{\ell}k_s} U_{\{\ell \rightarrow, s \rightarrow\}K}^{\pm} (\pm X + m) - \sum_{\substack{r=s+1 \\ r \neq \ell}}^{m+2} \delta_{k_s k_r} U_{\{s \rightarrow, r \rightarrow\}K}^{\pm} \right) (\pm X + m + n - 1) u \\ &= 2 \sum_{\substack{\ell, r=1 \\ \ell < r}}^{m+2} \delta_{k_{\ell}k_r} U_{\{\ell \rightarrow, r \rightarrow\}K}^{\pm} (\pm 2X + m) (\pm X + m + n - 1) u. \end{aligned}$$

Furthermore, we have, for $K \in \mathcal{A}^m$,

$$\sum_{q \in \mathcal{A}} [U_{qK}^\pm, U_q^\mp]u = 2U_K^\pm((m+n)(\pm X+m) - m)u - 2 \sum_{q \in \mathcal{A}} \sum_{\substack{s,p=1 \\ s < p}}^m \delta_{k_s k_p} U_{qq\{s \rightarrow, p \rightarrow\}K}^\pm u.$$

We finally compute

$$\begin{aligned} & \sum_{q \in \mathcal{A}} U_K^\pm U_q^\mp U_q^\mp u \\ &= 4 \sum_{\substack{\ell, r=1 \\ \ell < r}}^{m+2} \delta_{k_\ell k_r} U_{\{\ell \rightarrow, r \rightarrow\}K}^\pm X(2X \pm (n+2m-2))u + 4 \sum_{q \in \mathcal{A}} \sum_{\substack{\ell, r=1 \\ \ell < r}}^{m+2} \sum_{\substack{s,p=1 \\ s < p \\ \{s,p\} \cap \{\ell,r\} = \emptyset}}^{m+2} \delta_{k_\ell k_r} \delta_{k_s k_p} U_{qq\{\ell \rightarrow, r \rightarrow, s \rightarrow, p \rightarrow\}K}^\pm u, \end{aligned}$$

which finishes the proof. □

Arguing by induction using (4-4) and applying Lemma 4.5 to $\Delta_{\mp}^r u$, we get:

Lemma 4.6. *Assume that $u \in \mathcal{D}'(S\mathbb{H}^{n+1})$ and $\mathcal{U}_\pm^{m+1}u = 0$, $\mathcal{T}(\mathcal{U}_\pm^m u) = 0$. Then, for each $r \geq 0$,*

$$\mathcal{U}_\pm^{m+2r} \Delta_{\mp}^r u = (-1)^r 2^{2r} \left(\prod_{j=0}^{r-1} (\mathcal{X} \mp (m+j)) \right) \left(\prod_{j=1}^r (2\mathcal{X} \pm (n-2j)) \right) \mathcal{I}^r (\mathcal{U}_\pm^m u).$$

Moreover, for $r \geq 1$,

$$\mathcal{T}(\mathcal{U}_\pm^{m+2r} \Delta_{\mp}^r u) = (-1)^r 2^{2r} r(n+2m+2r-2) \left(\prod_{j=0}^{r-1} (\mathcal{X} \mp (m+j)) \right) \left(\prod_{j=1}^r (2\mathcal{X} \pm (n-2j)) \right) \mathcal{I}^{r-1} (\mathcal{U}_\pm^m u).$$

We are now ready to complete the proof of Lemma 4.2. Following (4-5), we decompose v as $v = \sum_{r=0}^{\lfloor m/2 \rfloor} \mathcal{I}^r(v_r)$ with $v_r \in \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes_S^{m-2r} \mathcal{E}^*)$ and $\mathcal{T}(v_r) = 0$. Since X commutes with \mathcal{T} and \mathcal{I} , we find $Xv_r = \pm \lambda v_r$. Moreover, since $\mathcal{U}_\pm v = 0$, we have $\mathcal{U}_\pm v_r = 0$. Put

$$u_r := (-\Delta_{\mp})^r \mathcal{V}_{\mp}^{m-2r} v_r \in \mathcal{D}'(S\mathbb{H}^{n+1}).$$

By Lemma 4.4 (applied to v_r) and Lemma 4.6 (applied to $\mathcal{V}_{\mp}^{m-2r} v_r$ and with m replaced by $m-2r$),

$$\begin{aligned} \mathcal{U}_\pm^m u_r &= 2^{2r} \left(\prod_{j=0}^{r-1} (\lambda - (m-2r+j)) \right) \left(\prod_{j=1}^r (2\lambda + n - 2j) \right) \mathcal{I}^r (\mathcal{U}_\pm^{m-2r} \mathcal{V}_{\mp}^{m-2r} v_r) \\ &= 2^m (m-2r)! \left(\prod_{j=n-1}^{n+m-2r-2} (\lambda + j) \right) \left(\prod_{j=m-2r}^{m-r-1} (\lambda - j) \right) \left(\prod_{j=1}^r (2\lambda + n - 2j) \right) \mathcal{I}^r(v_r). \end{aligned}$$

Since $\lambda \notin \frac{1}{2}\mathbb{Z}$, we see that $v = \mathcal{U}_\pm^m u$, where u is a linear combination of $u_0, \dots, u_{\lfloor m/2 \rfloor}$. The relation $Xu = \pm(\lambda - m)u$ follows immediately from (4-19) and (4-25). Finally, the equivariance property under G follows similarly to Lemma 4.1.

4D. Reduction to the conformal boundary. We now describe the tensors $v \in \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes_S^m \mathcal{E}^*)$ that satisfy $\mathcal{U}_\pm v = 0$ and $Xv = 0$ via symmetric tensors on the conformal boundary \mathbb{S}^n . For that we define the operators

$$\mathcal{Q}_\pm : \mathcal{D}'(\mathbb{S}^n; \otimes^m(T^*\mathbb{S}^n)) \rightarrow \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes^m \mathcal{E}^*)$$

by the following formula: if $w \in \mathcal{C}^\infty(\mathbb{S}^n; \otimes^m(T^*\mathbb{S}^n))$, we set, for $\eta_i \in \mathcal{E}(x, \xi)$,

$$\mathcal{Q}_\pm w(x, \xi)(\eta_1, \dots, \eta_m) := (w \circ B_\pm(x, \xi))(\mathcal{A}_\pm^{-1}(x, \xi)\eta_1, \dots, \mathcal{A}_\pm^{-1}(x, \xi)\eta_m), \tag{4-26}$$

where $\mathcal{A}_\pm(x, \xi) : T_{B_\pm(x, \xi)}\mathbb{S}^n \rightarrow \mathcal{E}(x, \xi)$ is the parallel transport defined in (3-31), and we see that the operator (4-26) extends continuously to $\mathcal{D}'(\mathbb{S}^n; \otimes_S^m(T^*\mathbb{S}^n))$, since the map $B_\pm : S\mathbb{H}^{n+1} \rightarrow \mathbb{S}^n$ defined in (3-16) is a submersion; see [Hörmander 1983, Theorem 6.1.2]; the result can be written as $\mathcal{Q}_\pm w = (\otimes^m(\mathcal{A}_\pm^{-1})^T) \cdot w \circ B_\pm$, where T denotes the transpose.

Lemma 4.7. *The operator \mathcal{Q}_\pm is a linear isomorphism from $\mathcal{D}'(\mathbb{S}^n; \otimes_S^m(T^*\mathbb{S}^n))$ onto the space*

$$\{v \in \mathcal{D}'(S\mathbb{H}^{n+1}; \otimes_S^m \mathcal{E}^*) \mid \mathcal{U}_\pm v = 0, \mathcal{X}v = 0\}. \tag{4-27}$$

Proof. It is clear that \mathcal{Q}_\pm is injective. Next, we show that the image of \mathcal{Q}_\pm is contained in (4-27). For that it suffices to show that, for $w \in \mathcal{C}^\infty(\mathbb{S}^n; \otimes_S^m(T^*\mathbb{S}^n))$, we have $\mathcal{U}_\pm(\mathcal{Q}_\pm w) = 0$ and $\mathcal{X}(\mathcal{Q}_\pm w) = 0$. We prove the first statement; the second one is established similarly. Let $\gamma \in G$, $w_1, \dots, w_m \in \mathcal{C}^\infty(\mathbb{S}^n; T\mathbb{S}^n)$, and $w_i^* = \langle w_i, \cdot \rangle_{g_{\mathbb{S}^n}}$ be the duals through the metric. Then

$$\begin{aligned} \mathcal{Q}_\pm(w_1^* \otimes \dots \otimes w_m^*)(\pi_S(\gamma)) &= \sum_{k_1, \dots, k_m=1}^n \left(\prod_{j=1}^m (w_j^* \circ B_\pm \circ \pi_S(\gamma))(\mathcal{A}_\pm^{-1}(\pi_S(\gamma))\gamma \cdot e_{k_j+1}) \right) \mathbf{e}_K^*(\gamma) \\ &= (-1)^m \sum_{k_1, \dots, k_m=1}^n \left(\prod_{j=1}^m \langle (\mathcal{A}_\pm \cdot w_j \circ B_\pm) \circ \pi_S(\gamma), \gamma \cdot e_{k_j+1} \rangle_M \right) \mathbf{e}_K^*(\gamma), \end{aligned}$$

where we have used (3-32) in the second identity. Now we have, from (3-31),

$$\mathcal{A}_\pm(\pi_S(\gamma))\zeta = (0, \zeta) - \langle (0, \zeta), \gamma \cdot e_0 \rangle_M \gamma (e_0 + e_1);$$

thus

$$\mathcal{Q}_\pm(w_1^* \otimes \dots \otimes w_m^*)(\pi_S(\gamma)) = \sum_{k_1, \dots, k_m=1}^n \left(\prod_{j=1}^m \langle (0, -w_j(B_\pm(\pi_S(\gamma)))) \rangle_M, \gamma \cdot e_{k_j+1} \right) \mathbf{e}_K^*(\gamma).$$

Since $d(B_\pm \circ \pi_S) \cdot U_\ell^\pm = 0$ by (4-8) and $U_\ell^\pm(\gamma \cdot e_{k_j+1}) = \gamma \cdot U_\ell^\pm \cdot e_{k_j+1}$ is a multiple of $\gamma \cdot (e_0 \pm e_1) = \Phi_\pm(\pi_S(\gamma))(1, B_\pm(\pi_S(\gamma)))$, we see that $\mathcal{U}_\pm(\mathcal{Q}_\pm w) = 0$ for all w .

It remains to show that, for v in (4-27), we have $v = \mathcal{Q}_\pm(w)$ for some w . For that, define

$$\tilde{v} = (\otimes^m \mathcal{A}_\pm^T) \cdot v \in \mathcal{D}'(S\mathbb{H}^{n+1}; B_\pm^*(\otimes_S^m T^*\mathbb{S}^n)),$$

where \mathcal{A}_\pm^T denotes the transpose of \mathcal{A}_\pm . Then $\mathcal{U}_\pm v = 0$ and $\mathcal{X}v = 0$ imply that $U_\ell^\pm(\pi_S^* \tilde{v}) = 0$ and $X\tilde{v} = 0$ (where, to define differentiation, we embed $T^*\mathbb{S}^n$ into \mathbb{R}^{n+1}). Additionally, $R_{i+1, j+1}(\pi_S^* \tilde{v}) = 0$; therefore $\pi_S^* \tilde{v}$ is constant on the right cosets of the subgroup $H_\pm \subset G$ defined in (3-27). Since

$(B_{\pm} \circ \pi_S)^{-1}(B_{\pm} \circ \pi_S(\gamma)) = \gamma H_{\pm}$, we see that \tilde{v} is the pullback under B_{\pm} of some $w \in \mathcal{D}'(\mathbb{S}^n; \otimes_S^m T^*\mathbb{S}^n)$, and it follows that $v = \mathcal{Q}_{\pm}(w)$. □

In fact, using (3-31) and the expression of $\xi_{\pm}(x, \nu)$ in (3-20) in terms of the Poisson kernel, it is not difficult to show that $\mathcal{Q}_{\pm}(w)$ belongs to a smaller space of *tempered* distributions: in the ball model, this can be described as the dual space to the Fréchet space of smooth sections of $\otimes^m({}^0S\bar{\mathbb{B}}^{n+1})$ over $\bar{\mathbb{B}}^{n+1}$ which vanish to infinite order at the conformal boundary $\mathbb{S}^n = \partial\bar{\mathbb{B}}^{n+1}$.

We finally give a useful criterion for invariance of $\mathcal{Q}_{\pm}(w)$ under the left action of an element of G :

Lemma 4.8. *Take $\gamma \in G$ and let $w \in \mathcal{D}'(\mathbb{S}^n; \otimes_S^m(T^*\mathbb{S}^n))$. Take $s \in \mathbb{C}$ and define $v = \Phi_{\pm}^s \mathcal{Q}_{\pm}(w)$. Then v is equivariant under left multiplication by γ , in the sense of (4-20), if and only if w satisfies the condition*

$$L_{\gamma}^* w(v) = N_{\gamma}(v)^{-s-m} w(v), \quad v \in \mathbb{S}^n. \tag{4-28}$$

Here $L_{\gamma}(v) \in \mathbb{S}^n$ and $N_{\gamma}(v) > 0$ are defined in (3-26).

Proof. The lemma follows by a direct calculation from (3-28) and (3-33). □

5. Pollicott–Ruelle resonances

In this section, we first recall the results of Butterley and Liverani [2007] and Faure and Sjöstrand [2011] on the Pollicott–Ruelle resonances for Anosov flows. We next state several useful microlocal properties of these resonances and prove Theorem 2, modulo properties of Poisson kernels (Lemma 5.8 and Theorem 6), which will be proved in Sections 6 and 7. Finally, we prove a pairing formula for resonances and Theorem 3.

5A. Definition and properties. We follow the presentation of [Faure and Sjöstrand 2011]; a more recent treatment using different technical tools is given in [Dyatlov and Zworski 2015]. We refer the reader to these two papers for the necessary notions of microlocal analysis.

Let \mathcal{M} be a smooth compact manifold of dimension $2n + 1$ and $\varphi_t = e^{tX}$ be an Anosov flow on \mathcal{M} , generated by a smooth vector field X . (In our case, $\mathcal{M} = SM$, $M = \Gamma \backslash \mathbb{H}^{n+1}$, and φ_t is the geodesic flow — see Section 5B.) The Anosov property is defined as follows: there exists a continuous splitting

$$T_y \mathcal{M} = E_0(y) \oplus E_u(y) \oplus E_s(y), \quad y \in \mathcal{M}, \quad E_0(y) := \mathbb{R}X(y), \tag{5-1}$$

invariant under $d\varphi_t$ and such that the stable/unstable subbundles $E_s, E_u \subset T\mathcal{M}$ satisfy, for some fixed smooth norm $|\cdot|$ on the fibers of $T\mathcal{M}$ and some constants C and $\theta > 0$,

$$\begin{aligned} |d\varphi_t(y)v| &\leq C e^{-\theta t} |v|, \quad v \in E_s(y), \\ |d\varphi_{-t}(y)v| &\leq C e^{-\theta t} |v|, \quad v \in E_u(y). \end{aligned} \tag{5-2}$$

We make an additional assumption that \mathcal{M} is equipped with a smooth measure μ which is invariant under φ_t , that is, $\mathcal{L}_X \mu = 0$.

We will use the dual decomposition to (5-1), given by

$$T_y^* \mathcal{M} = E_0^*(y) \oplus E_u^*(y) \oplus E_s^*(y), \quad y \in \mathcal{M}, \tag{5-3}$$

where E_0^*, E_u^*, E_s^* are dual to E_0, E_s, E_u respectively (note that E_u and E_s switch places), so for example $E_u^*(y)$ consists of covectors annihilating $E_0(y) \oplus E_u(y)$.

Following [Faure and Sjöstrand 2011, (1.24)], we now consider, for each $r \geq 0$, an *anisotropic Sobolev space*

$$\mathcal{H}^r(\mathcal{M}), \quad \text{where } \mathcal{C}^\infty(\mathcal{M}) \subset \mathcal{H}^r(\mathcal{M}) \subset \mathcal{D}'(\mathcal{M}).$$

Here we put $u := -r, s := r$ in [Faure and Sjöstrand 2011, Lemma 1.2]. Microlocally near E_u^* , the space \mathcal{H}^r is equivalent to the Sobolev space H^{-r} , in the sense that, for each pseudodifferential operator A of order 0 whose wavefront set is contained in a small enough conic neighborhood of E_u^* , the operator A is bounded, $\mathcal{H}^r \rightarrow H^{-r}$ and $H^{-r} \rightarrow \mathcal{H}^r$. Similarly, microlocally near E_s^* , the space \mathcal{H}^r is equivalent to the Sobolev space H^r . We also have $\mathcal{H}^0 = L^2$. The first-order differential operator X admits a unique closed unbounded extension from \mathcal{C}^∞ to \mathcal{H}^r ; see [Faure and Sjöstrand 2011, Lemma A.1].

The following theorem, defining Pollicott–Ruelle resonances associated to φ_t , is due to Faure and Sjöstrand [2011, Theorems 1.4 and 1.5]; see also [Dyatlov and Zworski 2015, Section 3.2].

Theorem 5. *Fix $r \geq 0$. Then the closed unbounded operator*

$$-X : \mathcal{H}^r(\mathcal{M}) \rightarrow \mathcal{H}^r(\mathcal{M})$$

has discrete spectrum in the region $\{\operatorname{Re} \lambda > -r/C_0\}$ for some constant C_0 independent of r . The eigenvalues of $-X$ on \mathcal{H}^r , called Pollicott–Ruelle resonances, and taken with multiplicities, do not depend on the choice of r as long as they lie in the appropriate region.

We have the following criterion for Pollicott–Ruelle resonances which does not use the \mathcal{H}^r spaces explicitly:

Lemma 5.1. *A number $\lambda \in \mathbb{C}$ is a Pollicott–Ruelle resonance of X if and only the space*

$$\operatorname{Res}_X(\lambda) := \{u \in \mathcal{D}'(\mathcal{M}) \mid (X + \lambda)u = 0, \operatorname{WF}(u) \subset E_u^*\} \tag{5-4}$$

is nontrivial. Here WF denotes the wavefront set; see, for instance, [Faure and Sjöstrand 2011, Definition 1.6]. The elements of $\operatorname{Res}_X(\lambda)$ are called resonant states associated to λ and the dimension of this space is called the geometric multiplicity of λ .

Proof. Assume first that λ is a Pollicott–Ruelle resonance. Take $r > 0$ such that $\operatorname{Re} \lambda > -r/C_0$. Then λ is an eigenvalue of $-X$ on \mathcal{H}^r , which implies that there exists nonzero $u \in \mathcal{H}^r$ such that $(X + \lambda)u = 0$. By [Faure and Sjöstrand 2011, Theorem 1.7], we have $\operatorname{WF}(u) \subset E_u^*$; thus u lies in (5-4).

Assume now that $u \in \mathcal{D}'(\mathcal{M})$ is a nonzero element of (5-4). For large enough r , we have $\operatorname{Re} \lambda > -r/C_0$ and $u \in H^{-r}(\mathcal{M})$. Since $\operatorname{WF}(u) \subset E_u^*$ and \mathcal{H}^r is equivalent to H^{-r} microlocally near E_u^* , we have $u \in \mathcal{H}^r$. Together with the identity $(X + \lambda)u$, this shows that λ is an eigenvalue of $-X$ on \mathcal{H}^r and thus a Pollicott–Ruelle resonance. □

For each λ with $\operatorname{Re} \lambda > -r/C_0$, the operator $X + \lambda : \mathcal{H}^r \rightarrow \mathcal{H}^r$ is Fredholm of index zero on its domain; this follows from the proof of Theorem 5. Therefore, $\dim \operatorname{Res}_X(\lambda)$ is equal to the dimension of the kernel

of the adjoint operator $X^* + \bar{\lambda}$ on the L^2 dual of \mathcal{H}^r , which we denote by \mathcal{H}^{-r} . Since $\frac{1}{i}X$ is symmetric on L^2 , we see that $\text{Res}_X(\lambda)$ has the same dimension as the following space of *coresonant states* at λ :

$$\text{Res}_{X^*}(\lambda) := \{u \in \mathcal{D}'(\mathcal{M}) \mid (X - \bar{\lambda})u = 0, \text{WF}(u) \subset E_s^*\}. \tag{5-5}$$

The main difference of (5-5) from (5-4) is that the subbundle E_s^* is used instead of E_u^* ; this can be justified by applying Lemma 5.1 to the vector field $-X$ instead of X , since the roles of the stable/unstable spaces for the corresponding flow φ_{-t} are reversed.

Note also that, for any $\lambda, \lambda^* \in \mathbb{C}$, one can define a pairing

$$\langle u, u^* \rangle \in \mathbb{C}, \quad u \in \text{Res}_X(\lambda), \quad u^* \in \text{Res}_{X^*}(\lambda^*). \tag{5-6}$$

One way to do that is to use the fact that wavefront sets of u and u^* intersect only at the zero section and apply [Hörmander 1983, Theorem 8.2.10]. An equivalent definition is obtained by noting that u is in \mathcal{H}^r and u^* is in \mathcal{H}^{-r} for $r > 0$ large enough and using the duality of \mathcal{H}^r and \mathcal{H}^{-r} . Note that, for $\lambda \neq \lambda^*$, we have $\langle u, u^* \rangle = 0$; indeed, $X(u\bar{u}^*) = (\lambda^* - \lambda)u\bar{u}^*$ integrates to 0. The question of computing the product $\langle u, u^* \rangle$ for $\lambda = \lambda^*$ is much more subtle and related to algebraic multiplicities; see Section 5C.

Since $\frac{1}{i}X$ is selfadjoint on $L^2 = \mathcal{H}^0$ (see [Faure and Sjöstrand 2011, Appendix A.1]), it has no eigenvalues on this space away from the real line; this implies that there are no Pollicott–Ruelle resonances in the right half-plane. In other words, we have:

Lemma 5.2. *The spaces $\text{Res}_X(\lambda)$ and $\text{Res}_{X^*}(\lambda)$ are trivial for $\text{Re } \lambda > 0$.*

Finally, we note that the results above apply to certain operators on vector bundles. More precisely, let \mathcal{E} be a smooth vector bundle over \mathcal{M} and assume that \mathcal{X} is a first-order differential operator on $\mathcal{D}'(\mathcal{M}; \mathcal{E})$ whose principal part is given by X , namely

$$\mathcal{X}(f\mathbf{u}) = f \mathcal{X}(\mathbf{u}) + (Xf)\mathcal{X}(\mathbf{u}), \quad f \in \mathcal{D}'(\mathcal{M}), \quad \mathbf{u} \in \mathcal{C}^\infty(\mathcal{M}; \mathcal{E}). \tag{5-7}$$

Assume moreover that \mathcal{E} is endowed with an inner product $\langle \cdot, \cdot \rangle_{\mathcal{E}}$ and $\frac{1}{i}\mathcal{X}$ is symmetric on L^2 with respect to this inner product and the measure μ . By an easy adaptation of the results of [Faure and Sjöstrand 2011] (see [Faure and Tsujii 2014; Dyatlov and Zworski 2015]), one can construct anisotropic Sobolev spaces $\mathcal{H}^r(\mathcal{M}; \mathcal{E})$ and Theorem 5 and Lemmas 5.1 and 5.2 apply to \mathcal{X} on these spaces.

5B. Proof of the main theorem. We now concentrate on the case

$$\mathcal{M} = SM = \Gamma \backslash (S\mathbb{H}^{n+1}), \quad M = \Gamma \backslash \mathbb{H}^{n+1},$$

with φ_t the geodesic flow. Here $\Gamma \subset G = \text{PSO}(1, n + 1)$ is a cocompact discrete subgroup with no fixed points, so that M is a compact smooth manifold. Henceforth we identify functions on the sphere bundle SM with functions on $S\mathbb{H}^{n+1}$ invariant under Γ , and similar identifications will be used for other geometric objects. It is important to note that *the constructions of the previous sections, except those involving the conformal infinity, are invariant under left multiplication by elements of G and thus descend naturally to SM .*

The lift of the geodesic flow on SM is the generator of the geodesic flow on $S\mathbb{H}^{n+1}$ (see Section 3C); both are denoted X . The lifts of the stable/unstable spaces E_s, E_u to $S\mathbb{H}^{n+1}$ are given in (3-14), and we see that (5-1) holds with $\theta = 1$. The invariant measure μ on SM is just the product of the volume measure on M and the standard measure on the fibers of SM induced by the metric.

Consider the bundle \mathcal{E} on SM defined in Section 3F. Then, for each m , the operator

$$\mathcal{X} : \mathcal{D}'(SM; \otimes_S^m \mathcal{E}^*) \rightarrow \mathcal{D}'(SM; \otimes_S^m \mathcal{E}^*)$$

defined in (4-17) satisfies (5-7) and $\frac{1}{i}\mathcal{X}$ is symmetric. The results of Section 5A apply both to X and \mathcal{X} .

Recall the operator \mathcal{U}_- introduced in Section 4B and its powers, for $m \geq 0$,

$$\mathcal{U}_-^m : \mathcal{D}'(SM) \rightarrow \mathcal{D}'(SM; \otimes_S^m \mathcal{E}^*).$$

The significance of \mathcal{U}_-^m for Pollicott–Ruelle resonances is explained by the following:

Lemma 5.3. *Assume that $\lambda \in \mathbb{C}$ is a Pollicott–Ruelle resonance of X and $u \in \text{Res}_X(\lambda)$ is a corresponding resonant state as defined in (5-4). Then*

$$\mathcal{U}_-^m u = 0 \quad \text{for } m > -\text{Re } \lambda.$$

Proof. By (4-19),

$$(\mathcal{X} + \lambda + m)\mathcal{U}_-^m u = 0.$$

Note also that $\text{WF}(\mathcal{U}_-^m u) \subset E_u^*$, since $\text{WF}(u) \subset E_u^*$ and \mathcal{U}_-^m is a differential operator. Since $\lambda + m$ lies in the right half-plane, it remains to apply Lemma 5.2 to $\mathcal{U}_-^m u$. \square

We can then use the operators \mathcal{U}_-^m to split the resonance spectrum into bands:

Lemma 5.4. *Assume that $\lambda \in \mathbb{C} \setminus \frac{1}{2}\mathbb{Z}$. Then*

$$\dim \text{Res}_X(\lambda) = \sum_{m \geq 0} \dim \text{Res}_{\mathcal{X}}^m(\lambda + m), \tag{5-8}$$

where

$$\text{Res}_{\mathcal{X}}^m(\lambda) := \{v \in \mathcal{D}'(SM; \otimes_S^m \mathcal{E}^*) \mid (\mathcal{X} + \lambda)v = 0, \mathcal{U}_- v = 0, \text{WF}(v) \subset E_u^*\}. \tag{5-9}$$

The space $\text{Res}_{\mathcal{X}}^m(\lambda)$ is trivial for $\text{Re } \lambda > 0$ (by Lemma 5.2). If $\lambda \in \frac{1}{2}\mathbb{Z}$, then we have

$$\dim \text{Res}_X(\lambda) \leq \sum_{m \geq 0} \dim \text{Res}_{\mathcal{X}}^m(\lambda + m). \tag{5-10}$$

Proof. Denote, for $m \geq 1$,

$$V_m(\lambda) := \{u \in \mathcal{D}'(SM) \mid (X + \lambda)u = 0, \mathcal{U}_-^m u = 0, \text{WF}(u) \subset E_u^*\}.$$

Clearly, $V_m(\lambda) \subset V_{m+1}(\lambda)$. Moreover, by Lemma 5.3 we have $\text{Res}_X(\lambda) = V_m(\lambda)$ for m large enough depending on λ . By (4-19), the operator \mathcal{U}_-^m acts as

$$\mathcal{U}_-^m : V_{m+1}(\lambda) \rightarrow \text{Res}_{\mathcal{X}}^m(\lambda + m), \tag{5-11}$$

and the kernel of (5-11) is exactly $V_m(\lambda)$, with the convention that $V_0(\lambda) = 0$. Therefore,

$$\dim V_{m+1}(\lambda) \leq \dim V_m(\lambda) + \dim \text{Res}_{\mathcal{X}}^m(\lambda + m)$$

and (5-10) follows.

To show (5-8), it remains to prove that the operator (5-11) is onto; this follows from Lemma 4.2 (which does not enlarge the wavefront set of the resulting distribution, since it only employs differential operators in the proof). \square

The space $\text{Res}_{\mathcal{X}}^m(\lambda + m)$ is called the space of *resonant states at λ associated to the m -th band*; later we see that most of the corresponding Pollicott–Ruelle resonances satisfy $\text{Re } \lambda = -\frac{1}{2}n - m$. Similarly, we can describe $\text{Res}_{\mathcal{X}^*}(\lambda)$ via the spaces $\text{Res}_{\mathcal{X}^*}^m(\lambda + m)$, where

$$\text{Res}_{\mathcal{X}^*}^m(\lambda) := \{v \in \mathcal{D}'(SM; \otimes_S^m \mathcal{E}^*) \mid (\mathcal{X} - \bar{\lambda})v = 0, \mathcal{U}_+ v = 0, \text{WF}(v) \subset E_S^*\}; \tag{5-12}$$

note that here \mathcal{U}_+ is used in place of \mathcal{U}_- .

We further decompose $\text{Res}_{\mathcal{X}}^m(\lambda)$ using *trace-free* tensors:

Lemma 5.5. *Recall the homomorphisms $\mathcal{T} : \otimes_S^m \mathcal{E}^* \rightarrow \otimes_S^{m-2} \mathcal{E}^*$, $\mathcal{I} : \otimes_S^m \mathcal{E}^* \rightarrow \otimes_S^{m-2} \mathcal{E}^*$ defined in Section 4A (we put $\mathcal{T} = 0$ for $m = 0, 1$). Define the space*

$$\text{Res}_{\mathcal{X}}^{m,0}(\lambda) := \{v \in \text{Res}_{\mathcal{X}}^m(\lambda) \mid \mathcal{T}(v) = 0\}. \tag{5-13}$$

Then for all $m \geq 0$ and λ ,

$$\dim \text{Res}_{\mathcal{X}}^m(\lambda) = \sum_{\ell=0}^{\lfloor m/2 \rfloor} \dim \text{Res}_{\mathcal{X}}^{m-2\ell,0}(\lambda). \tag{5-14}$$

In fact,

$$\text{Res}_{\mathcal{X}}^{m,0}(\lambda) = \bigoplus_{\ell=0}^{\lfloor m/2 \rfloor} \mathcal{I}^\ell(\text{Res}_{\mathcal{X}}^{m-2\ell,0}(\lambda)). \tag{5-15}$$

Proof. The identity (5-15) follows immediately from (4-5); it is straightforward to see that the defining properties of $\text{Res}_{\mathcal{X}}^m(\lambda)$ are preserved by the canonical tensorial operations involved. The identity (5-14) then follows since \mathcal{I} is one-to-one by the paragraph following (4-4). \square

The elements of $\text{Res}_{\mathcal{X}}^{m,0}(\lambda)$ can be expressed via distributions on the conformal boundary \mathbb{S}^n :

Lemma 5.6. *Let Q_- be the operator defined in (4-26); recall that it is injective. If $\pi_\Gamma : S\mathbb{H}^{n+1} \rightarrow SM$ is the natural projection map, then*

$$\pi_\Gamma^* \text{Res}_{\mathcal{X}}^{m,0}(\lambda) = \Phi_-^\lambda Q_-(\text{Bd}^{m,0}(\lambda)),$$

where $\text{Bd}^{m,0}(\lambda) \subset \mathcal{D}'(\mathbb{S}^n; \otimes_S^m(T^*\mathbb{S}^n))$ consists of all distributions w such that $\mathcal{T}(w) = 0$ and

$$L_\gamma^* w(v) = N_\gamma(v)^{-\lambda-m} w(v), \quad v \in \mathbb{S}^n, \gamma \in \Gamma; \tag{5-16}$$

L_γ and N_γ are as defined in (3-26). Similarly

$$\pi_\Gamma^* \text{Res}_{\mathcal{X}^*}^{m,0}(\lambda) = \Phi_+^{\bar{\lambda}} Q_+(\text{Bd}^{m,0}(\bar{\lambda})), \quad \text{Bd}^{m,0}(\bar{\lambda}) = \overline{\text{Bd}^{m,0}(\lambda)}.$$

Proof. Assume first that $w \in \text{Bd}^{m,0}(\lambda)$ and put $\tilde{v} = \Phi_-^\lambda \mathcal{Q}_-(w)$. Then, by Lemma 4.8 and (5-16), \tilde{v} is invariant under Γ and thus descends to a distribution $v \in \mathcal{D}'(SM; \otimes_S^m \mathcal{E}^*)$. Since $X\Phi_-^\lambda = -\lambda\Phi_-^\lambda$ and $U_j^-(\Phi_-^\lambda \circ \pi_S) = 0$ by (3-17) and (4-8), and \mathcal{X} and \mathcal{U}_- annihilate the image of \mathcal{Q}_- by Lemma 4.7, we have $(\mathcal{X} + \lambda)v = 0$ and $\mathcal{U}_-v = 0$. Moreover, by [Hörmander 1983, Theorem 8.2.4] the wavefront set of \tilde{v} is contained in the conormal bundle to the fibers of the map B_- ; by (3-25), we see that $\text{WF}(v) \subset E_u^*$. Finally, $\mathcal{T}(v) = 0$ since the map $\mathcal{A}_-(x, \xi)$ used in the definition of \mathcal{Q}_- is an isometry. Therefore, $v \in \text{Res}_\mathcal{X}^{m,0}(\lambda)$ and we have proved the containment $\pi_\Gamma^* \text{Res}_\mathcal{X}^{m,0}(\lambda) \supset \Phi_-^\lambda \mathcal{Q}_-(\text{Bd}^{m,0}(\lambda))$. The opposite containment is proved by reversing this argument. \square

Remark. It follows from the proof of Lemma 5.6 that the condition $\text{WF}(v) \subset E_u^*$ in (5-9) is unnecessary. This could also be seen by applying [Hörmander 1994, Theorem 18.1.27] to the equations $(\mathcal{X} + \lambda)v = 0$ and $\mathcal{U}_-v = 0$, since \mathcal{X} differentiates along the direction E_0 , \mathcal{U}_- differentiates along the direction E_u (see (4-11) and (4-16)), and the annihilator of $E_0 \oplus E_u$ (that is, the joint critical set of $\mathcal{X} + \lambda, \mathcal{U}_-$) is exactly E_u^* .

It now remains to relate the space $\text{Bd}^{m,0}(\lambda)$ to an eigenspace of the Laplacian on symmetric tensors. For that, we introduce the following operator, obtained by integrating the corresponding elements of $\text{Res}_\mathcal{X}^{m,0}(\lambda)$ along the fibers of \mathbb{S}^n :

Definition 5.7. Take $\lambda \in \mathbb{C}$. The *Poisson operators*

$$\mathcal{P}_\lambda^\pm : \mathcal{D}'(\mathbb{S}^n; \otimes^m T^*\mathbb{S}^n) \rightarrow C^\infty(\mathbb{H}^{n+1}; \otimes^m T^*\mathbb{H}^{n+1})$$

are defined by the formulas

$$\begin{aligned} \mathcal{P}_\lambda^- w(x) &= \int_{S_x \mathbb{H}^{n+1}} \Phi_-(x, \xi)^\lambda \mathcal{Q}_-(w)(x, \xi) dS(\xi), \\ \mathcal{P}_\lambda^+ w(x) &= \int_{S_x \mathbb{H}^{n+1}} \Phi_+(x, \xi)^{\bar{\lambda}} \mathcal{Q}_+(w)(x, \xi) dS(\xi). \end{aligned} \tag{5-17}$$

Here, integration of elements of $\otimes^m \mathcal{E}^*(x, \xi)$ is performed by embedding them in $\otimes^m T_x^* \mathbb{H}^{n+1}$ using composition with the orthogonal projection $T_x \mathbb{H}^{n+1} \rightarrow \mathcal{E}(x, \xi)$.

The operators \mathcal{P}_λ^\pm are related by the identity

$$\overline{\mathcal{P}_\lambda^\pm w} = \mathcal{P}_\lambda^\mp \bar{w}. \tag{5-18}$$

By Lemma 5.6, \mathcal{P}_λ^- maps $\text{Bd}^{m,0}(\lambda)$ onto symmetric Γ -equivariant tensors, which can thus be considered as elements of $C^\infty(M; \otimes_S^m T^*M)$. The relation with the Laplacian is given by the following fact, proved in Section 6C:

Lemma 5.8. *The image of $\text{Bd}^{m,0}(\lambda)$ under \mathcal{P}_λ^- is contained in the eigenspace $\text{Eig}^m(-\lambda(n + \lambda) + m)$ for each λ , where*

$$\text{Eig}^m(\sigma) := \{f \in C^\infty(M; \otimes_S^m T^*M) \mid \Delta f = \sigma f, \nabla^* f = 0, \mathcal{T}(f) = 0\}. \tag{5-19}$$

Here the trace \mathcal{T} was defined in Section 4A and the Laplacian Δ and the divergence ∇^* are introduced in Section 6A. (A similar result for \mathcal{P}_λ^+ follows from (5-18).)

Furthermore, in Sections 6C and 7 we show the crucial:

Theorem 6. Assume that $\lambda \notin \mathcal{R}_m$, where

$$\mathcal{R}_m = \begin{cases} -\frac{1}{2}n - \frac{1}{2}\mathbb{N}_0 & \text{if } n > 1 \text{ or } m = 0, \\ -\frac{1}{2}\mathbb{N}_0 & \text{if } n = 1 \text{ and } m > 0. \end{cases} \tag{5-20}$$

Then the map $\mathcal{P}_\lambda^- : \text{Bd}^{m,0}(\lambda) \rightarrow \text{Eig}^m(-\lambda(n + \lambda) + m)$ is an isomorphism.

Remark. In Theorem 6, the set of exceptional points where we do not show isomorphism is not optimal but is sufficient for our application (we only need $\mathcal{R}_m \subset m - \frac{1}{2}n - \frac{1}{2}\mathbb{N}_0$); we expect the exceptional set to be contained in $-n + 1 - \mathbb{N}_0$. This result is known for functions, that is for $m = 0$, with the exceptional set being $-n - \mathbb{N}$. This was proved by Helgason [1974] and Minemura [1975] in the case of hyperfunctions on \mathbb{S}^n and by Oshima and Sekiguchi [1980] and Schlichtkrull and van den Ban [1987] for distributions; Grellier and Otal [2005] studied the sharp functional spaces on \mathbb{S}^n of the boundary values of bounded eigenfunctions on \mathbb{H}^{n+1} . The extension to $m > 0$ does not seem to be known in the literature and is not trivial: it takes most of Sections 6 and 7.

We finally provide the following refinement of Lemma 5.4, needed to handle the case $\lambda \in (-\frac{1}{2}n, \infty) \cap \frac{1}{2}\mathbb{Z}$:

Lemma 5.9. Assume that $\lambda \in -\frac{1}{2}n + \frac{1}{2}\mathbb{N}$. If $\lambda \in -2\mathbb{N}$, then

$$\dim \text{Res}_X(\lambda) = \sum_{\substack{m \geq 0 \\ m \neq -\lambda}} \dim \text{Res}_X^m(\lambda + m).$$

If $\lambda \notin -2\mathbb{N}$, then (5-8) holds.

Proof. We use the proof of Lemma 5.4. We first show that, for m odd or $\lambda \neq -m$,

$$\mathcal{U}_-^m(V_{m+1}(\lambda)) = \text{Res}_X^m(\lambda + m). \tag{5-21}$$

Using (5-15), it suffices to prove that, for $0 \leq \ell \leq \frac{1}{2}m$, the space $\mathcal{I}^\ell(\text{Res}_X^{m-2\ell,0}(\lambda + m))$ is contained in $\mathcal{U}_-^m(V_{m+1}(\lambda))$. This follows from the proof of Lemma 4.2 as long as

$$\begin{aligned} \lambda + m &\notin \mathbb{Z} \cap ([2\ell + 2 - n - m, 1 - n] \cup [m - 2\ell, m - \ell - 1]), \\ \lambda + m + \frac{1}{2}n &\notin \mathbb{Z} \cap [1, \ell]; \end{aligned}$$

using that $\lambda > -\frac{1}{2}n$, it suffices to prove that

$$\lambda \notin \mathbb{Z} \cap [-2\ell, -\ell - 1]. \tag{5-22}$$

On the other hand, by Lemma 5.6, Theorem 6, and Lemma 6.1, if $\ell < \frac{1}{2}m$ and the space $\text{Res}_X^{m-2\ell,0}(\lambda + m)$ is nontrivial, then

$$-(\lambda + m + \frac{1}{2}n)^2 + \frac{1}{4}n^2 + m - 2\ell \geq m - 2\ell + n - 1,$$

implying

$$|\lambda + m + \frac{1}{2}n| \leq |\frac{1}{2}n - 1|, \tag{5-23}$$

and (5-22) follows. For the case $\ell = \frac{1}{2}m$, since $\Delta \geq 0$ on functions we have

$$-(\lambda + m + \frac{1}{2}n)^2 + \frac{1}{4}n^2 \geq 0,$$

which implies that $\lambda \leq -m$ and thus (5-22) holds unless $\lambda = -m$.

It remains to consider the case when $m = 2\ell$ is even and $\lambda = -m$. We have

$$\text{Res}_{\mathcal{X}}^m(0) = \mathcal{I}^\ell(\text{Res}_{\mathcal{X}}^{0,0}(0));$$

that is, $\text{Res}_{\mathcal{X}}^{m-2\ell',0}(0)$ is trivial for $\ell' < \frac{1}{2}m$. For $n > 1$, this follows immediately from (5-23), and, for $n = 1$, since the bundle \mathcal{E}^* is one-dimensional, we get $\text{Res}_{\mathcal{X}}^{m',0}(\lambda) = 0$ for $m' \geq 2$. Now, $\text{Res}_{\mathcal{X}}^{0,0}(0) = \text{Res}_{\mathcal{X}}^0(0)$ corresponds via Lemma 5.6 and Theorem 6 to the kernel of the scalar Laplacian, that is, to the space of constant functions. Therefore, $\text{Res}_{\mathcal{X}}^{0,0}$ is one-dimensional and it is spanned by the constant function 1 on SM ; it follows that $\text{Res}_{\mathcal{X}}^m(0)$ is spanned by $\mathcal{I}^\ell(1)$. However, by Lemma 4.3, for each $u \in \mathcal{D}'(SM)$,

$$\langle \mathcal{I}^\ell(1), \mathcal{U}_-^m u \rangle_{L^2} = (-1)^m \langle \mathcal{V}_-^m \mathcal{I}^\ell(1), u \rangle_{L^2} = 0.$$

Since $\mathcal{U}_-^m(V_{m+1}(\lambda)) \subset \text{Res}_{\mathcal{X}}^m(0)$, we have $\mathcal{U}_-^m = 0$ on $V_{m+1}(\lambda)$, which implies that $V_{m+1}(\lambda) = V_m(\lambda)$, finishing the proof. □

To prove Theorem 2, it now suffices to combine Lemmas 5.4–5.9 with Theorem 6.

5C. Resonance pairing and algebraic multiplicity. In this section, we prove Theorem 3. The key component is a pairing formula which states that the inner product between a resonant and a coresonant state, defined in (5-6), is determined by the inner product between the corresponding eigenstates of the Laplacian. The nondegeneracy of the resulting inner product as a bilinear operator on $\text{Res}_{\mathcal{X}}(\lambda) \times \text{Res}_{\mathcal{X}^*}(\lambda)$ for $\lambda \notin \frac{1}{2}\mathbb{Z}$ immediately implies the fact that the algebraic and geometric multiplicities of λ coincide (that is, $X + \lambda$ does not have any nontrivial Jordan cells).

To state the pairing formula, we first need a decomposition of the space $\text{Res}_{\mathcal{X}}(\lambda)$, which is an effective version of the formulas (5-8) and (5-14). Take $m \geq 0$, $\ell \leq \lfloor m/2 \rfloor$ and $w \in \text{Bd}^{m-2\ell,0}(\lambda)$. Let \mathcal{I} be the operator defined in Section 4A. Then (5-15) and Lemma 5.6 show that

$$\text{Res}_{\mathcal{X}}^m(\lambda) = \bigoplus_{\ell=0}^{\lfloor m/2 \rfloor} \mathcal{I}^\ell(\text{Res}_{\mathcal{X}}^{m-2\ell,0}(\lambda)) = \bigoplus_{\ell=0}^{\lfloor m/2 \rfloor} \mathcal{I}^\ell(\Phi_-^\lambda \mathcal{Q}_-(\text{Bd}^{m-2\ell,0}(\lambda))).$$

Next, let

$$\mathcal{V}_\pm^m : \mathcal{D}'(SM; \otimes_S^m \mathcal{E}^*) \rightarrow \mathcal{D}'(SM) \quad \text{and} \quad \Delta_\pm : \mathcal{D}'(SM) \rightarrow \mathcal{D}'(SM)$$

be the operators introduced in Section 4C. Then the proofs of Lemma 5.4 and Lemma 4.2 show that, for $\lambda \notin \frac{1}{2}\mathbb{Z}$,

$$\begin{aligned} \text{Res}_{\mathcal{X}}(\lambda) &= \bigoplus_{m \geq 0} \bigoplus_{\ell=0}^{\lfloor m/2 \rfloor} V_{m\ell}(\lambda), & V_{m\ell}(\lambda) &:= \Delta_+^\ell \mathcal{V}_+^{m-2\ell}(\Phi_-^{\lambda+m} \mathcal{Q}_-(\text{Bd}^{m-2\ell,0}(\lambda+m))), \\ \text{Res}_{\mathcal{X}^*}(\lambda) &= \bigoplus_{m \geq 0} \bigoplus_{\ell=0}^{\lfloor m/2 \rfloor} V_{m\ell}^*(\lambda), & V_{m\ell}^*(\lambda) &:= \Delta_-^\ell \mathcal{V}_-^{m-2\ell}(\Phi_+^{\bar{\lambda}+m} \overline{\mathcal{Q}_+(\text{Bd}^{m-2\ell,0}(\lambda+m))}), \end{aligned} \tag{5-24}$$

and the operators in the definitions of $V_{m\ell}(\lambda)$, $V_{m\ell}^*(\lambda)$ are one-to-one on the corresponding spaces. By the proof of [Lemma 5.9](#), the decomposition (5-24) is also valid for $\lambda \in (-\frac{1}{2}n, \infty) \setminus (-2\mathbb{N})$; for $\lambda \in (-\frac{1}{2}n, \infty) \cap (-2\mathbb{N})$, we have

$$\text{Res}_X(\lambda) = \bigoplus_{\substack{m \geq 0 \\ m \neq -\lambda}} \bigoplus_{\ell=0}^{\lfloor m/2 \rfloor} V_{m\ell}(\lambda), \quad \text{Res}_{X^*}(\lambda) = \bigoplus_{\substack{m \geq 0 \\ m \neq -\lambda}} \bigoplus_{\ell=0}^{\lfloor m/2 \rfloor} V_{m\ell}^*(\lambda). \tag{5-25}$$

We can now state the pairing formula:

Lemma 5.10. *Let $\lambda \notin -\frac{1}{2}n - \frac{1}{2}\mathbb{N}_0$ and $u \in \text{Res}_X(\lambda)$, $u^* \in \text{Res}_{X^*}(\lambda)$. Let $\langle u, u^* \rangle_{L^2(SM)}$ be defined by (5-6). Then:*

- (1) *If $u \in V_{m\ell}(\lambda)$, $u^* \in V_{m'\ell'}^*(\lambda)$, and $(m, \ell) \neq (m', \ell')$, then $\langle u, u^* \rangle_{L^2(SM)} = 0$.*
- (2) *If $u \in V_{m\ell}(\lambda)$, $u^* \in V_{m\ell}^*(\lambda)$, and $w \in \text{Bd}^{m-2\ell, 0}(\lambda + m)$ and $w^* \in \overline{\text{Bd}^{m-2\ell, 0}(\lambda + m)}$ are the elements generating u and u^* according to (5-24), then*

$$\langle u, u^* \rangle_{L^2(SM)} = c_{m\ell}(\lambda) \langle \mathcal{P}_{\lambda+m}^-(w), \mathcal{P}_{\lambda+m}^+(w^*) \rangle_{L^2(M)}, \tag{5-26}$$

where

$$c_{m\ell}(\lambda) = 2^{m+2\ell-n} \pi^{-1-n/2} \ell! (m-2\ell)! \sin(\pi(\frac{1}{2}n + \lambda)) \times \frac{\Gamma(m + \frac{1}{2}n - \ell) \Gamma(\lambda + n + 2m - 2\ell) \Gamma(-\lambda - \ell) \Gamma(-\lambda - m - \frac{1}{2}n + \ell + 1)}{\Gamma(m + \frac{1}{2}n - 2\ell) \Gamma(-\lambda - 2\ell)},$$

and, under the conditions (i) either $\lambda \notin -2\mathbb{N}$ or $m \neq -\lambda$ and (ii) $V_{m\ell}(\lambda)$ is nontrivial, we have $c_{m\ell}(\lambda) \neq 0$.

Remark. (i) The proofs below are rather technical, and it is suggested that the reader start with the case of resonances in the first band, $m = \ell = 0$, which preserves the essential analytic difficulties of the proof but considerably reduces the amount of calculation needed (in particular, one can go immediately to [Lemma 5.11](#), and the proof of this lemma for the case $m = \ell = 0$ does not involve the operator \mathcal{C}_η). We have

$$c_{00}(\lambda) = (4\pi)^{-n/2} \frac{\Gamma(n + \lambda)}{\Gamma(\frac{1}{2}n + \lambda)}.$$

(ii) In the special case of $n = 1$, $m = \ell = 0$, [Lemma 5.10](#) is a corollary of [[Anantharaman and Zelditch 2007](#), Theorem 1.2], where the product $uu^* \in \mathcal{D}'(SM)$ lifts to a Patterson–Sullivan distribution on $S\mathbb{H}^2$. In general, if $|\text{Re } \lambda| \leq C$ and $\text{Im } \lambda \rightarrow \infty$, then $c_{m\ell}(\lambda)$ grows like $|\lambda|^{n/2+m}$.

[Lemma 5.10](#) immediately gives:

Proof of Theorem 3. By [Theorem 6](#), we know that

$$\mathcal{P}_\lambda^- : \text{Bd}^{m-2\ell, 0}(\lambda + m) \rightarrow \text{Eig}^{m-2\ell} \left(-(\lambda + m + \frac{1}{2}n)^2 + \frac{1}{4}n^2 + m - 2\ell \right)$$

is an isomorphism. Given (5-18), we also get the isomorphism

$$\mathcal{P}_\lambda^+ : \overline{\text{Bd}^{m-2\ell, 0}(\lambda + m)} \rightarrow \text{Eig}^{m-2\ell} \left(-(\lambda + m + \frac{1}{2}n)^2 + \frac{1}{4}n^2 + m - 2\ell \right).$$

Here we used that the target space is invariant under complex conjugation. By [Lemma 5.10](#), the bilinear product

$$\text{Res}_X(\lambda) \times \text{Res}_{X^*}(\lambda) \rightarrow \mathbb{C}, \quad (u, u^*) \mapsto \langle u, u^* \rangle_{L^2(SM)} \tag{5-27}$$

is nondegenerate, since the $L^2(M)$ inner product restricted to $\text{Eig}^{m-2\ell}(-(\lambda + m + \frac{1}{2}n)^2 + \frac{1}{4}n^2 + m - 2\ell)$ is nondegenerate for all m, ℓ .

Assume now that $\tilde{u} \in \mathcal{D}'(SM)$ satisfies $(X + \lambda)^2\tilde{u} = 0$ and $\tilde{u} \in \mathcal{H}^r$ for some $r, \text{Re } \lambda > -r/C_0$; we need to show that $(X + \lambda)\tilde{u} = 0$. Put $u := (X + \lambda)\tilde{u}$. Then $u \in \text{Res}_X(\lambda)$. However, u also lies in the image of $X + \lambda$ on \mathcal{H}^r ; therefore we have $\langle u, u^* \rangle = 0$ for each $u^* \in \text{Res}_{X^*}(\lambda)$. Since the product (5-27) is nondegenerate, we see that $u = 0$, finishing the proof. \square

In the remaining part of this section, we prove [Lemma 5.10](#). Take some $m, m', \ell, \ell' \geq 0$ such that $2\ell \leq m, 2\ell' \leq m'$, and consider $u \in V_{m\ell}(\lambda), u^* \in V_{m'\ell'}(\lambda)$ given by

$$u = \Delta_+^\ell \mathcal{V}_+^{m-2\ell} v, \quad u^* = \Delta_-^{\ell'} \mathcal{V}_-^{m'-2\ell'} v^*,$$

where, for some $w \in \text{Bd}^{m-2\ell,0}(\lambda + m)$ and $w^* \in \overline{\text{Bd}^{m'-2\ell',0}(\lambda + m')}$,

$$v = \Phi_-^{\lambda+m} \mathcal{Q}_-(w) \in \text{Res}_X^{m-2\ell,0}(\lambda + m), \quad v^* = \Phi_+^{\bar{\lambda}+m'} \mathcal{Q}_+(w^*) \in \text{Res}_{X^*}^{m'-2\ell',0}(\lambda + m').$$

Using [Lemma 4.3](#) and the fact that Δ_\pm are symmetric, we get

$$\langle u, u^* \rangle_{L^2(SM)} = (-1)^{m'} \langle \mathcal{U}_-^{m'-2\ell'} \Delta_-^{\ell'} \Delta_+^\ell \mathcal{V}_+^{m-2\ell} v, v^* \rangle_{L^2(SM; \otimes^{m'-2\ell'} \mathcal{E}^*)}.$$

By [Lemmas 4.4](#) and [4.6](#), $\mathcal{U}_-^{m+1} \Delta_+^\ell \mathcal{V}_+^{m-2\ell} v = 0$. Therefore, if $m' > m$, we derive that $\langle u, u^* \rangle_{L^2(SM)} = 0$; by swapping u and u^* , one can similarly handle the case $m' < m$. We therefore assume that $m = m'$. Then, by [Lemmas 4.4](#) and [4.6](#) (see the proof of [Lemma 4.2](#)),

$$\begin{aligned} & (-1)^{\ell+\ell'} \mathcal{U}_-^{m-2\ell'} \Delta_-^{\ell'} \Delta_+^\ell \mathcal{V}_+^{m-2\ell} v \\ &= \mathcal{T}^{\ell'} \mathcal{U}_-^m (-\Delta_+)^{\ell'} \mathcal{V}_+^{m-2\ell} v \\ &= 2^{m+\ell} (m-2\ell)! \frac{\Gamma(\lambda + n + 2m - 2\ell - 1) \Gamma(-\lambda - \ell) \Gamma(-\lambda - m - \frac{1}{2}n + \ell + 1)}{\Gamma(\lambda + m + n - 1) \Gamma(-\lambda - 2\ell) \Gamma(-\lambda - m - \frac{1}{2}n + 1)} \mathcal{T}^{\ell'} \mathcal{I}^\ell v. \end{aligned}$$

If $\ell' > \ell$, this implies that $\langle u, u^* \rangle_{L^2(SM)} = 0$, and the case $\ell' < \ell$ is handled similarly. (Recall that $\mathcal{T}(v) = 0$.) We therefore assume that $m = m', \ell = \ell'$. In this case, by (4-4),

$$\mathcal{T}^\ell \mathcal{I}^\ell v = 2^\ell \ell! \frac{\Gamma(m + \frac{1}{2}n - \ell)}{\Gamma(m + \frac{1}{2}n - 2\ell)} v,$$

which implies that

$$\begin{aligned} \langle u, u^* \rangle_{L^2(SM)} &= (-2)^{m+2\ell} \ell! (m-2\ell)! \frac{\Gamma(m + \frac{1}{2}n - \ell) \Gamma(\lambda + n + 2m - 2\ell - 1)}{\Gamma(m + \frac{1}{2}n - 2\ell) \Gamma(\lambda + n + m - 1)} \\ &\quad \times \frac{\Gamma(-\lambda - \ell) \Gamma(-\lambda - m - \frac{1}{2}n + \ell + 1)}{\Gamma(-\lambda - 2\ell) \Gamma(-\lambda - m - \frac{1}{2}n + 1)} \langle v, v^* \rangle_{L^2(SM; \otimes^{m-2\ell} \mathcal{E}^*)}. \end{aligned}$$

Note that, under assumptions (i) and (ii) of [Lemma 5.10](#), the coefficient in the formula above is nonzero; see the proof of [Lemma 5.9](#).

It then remains to prove the following identity (note that the coefficient there is nonzero for $\lambda \notin \mathbb{Z}$ or $\operatorname{Re} \lambda > m - \frac{1}{2}n$):

Lemma 5.11. *Assume that $v \in \operatorname{Res}_{\mathcal{X}}^{m,0}(\lambda)$ and $v^* \in \operatorname{Res}_{\mathcal{X}^*}^{m,0}(\lambda)$. Define*

$$f(x) := \int_{S_x M} v(x, \xi) dS(\xi), \quad f^*(x) := \int_{S_x M} v^*(x, \xi) dS(\xi),$$

where integration of tensors is understood as in [Definition 5.7](#). If $\lambda \notin -(\frac{1}{2}n + \mathbb{N}_0)$, then

$$\langle f, f^* \rangle_{L^2(M; \otimes^m T^* M)} = 2^n \pi^{n/2} \frac{\Gamma(\frac{1}{2}n + \lambda)}{(n + \lambda + m - 1)\Gamma(n - 1 + \lambda)} \langle v, v^* \rangle_{L^2(SM; \otimes^m \mathcal{E}^*)}.$$

Proof. We write

$$\langle f, f^* \rangle_{L^2(M; \otimes^m T^* M)} = \int_{S^2 M} \langle v(y, \eta_-), \overline{v^*(y, \eta_+)} \rangle_{\otimes^m T^*_y M} dy d\eta_- d\eta_+, \tag{5-28}$$

where the bundle $S^2 M$ is given by

$$S^2 M = \{(y, \eta_-, \eta_+) \mid y \in M, \eta_{\pm} \in S_y M\}.$$

Define also

$$S^2_{\Delta} M = \{(y, \eta_-, \eta_+) \in S^2 M \mid \eta_- + \eta_+ \neq 0\}.$$

On the other hand,

$$\langle v, v^* \rangle_{L^2(SM; \otimes^m \mathcal{E}^*)} = \int_{SM} \langle v(x, \xi), \overline{v^*(x, \xi)} \rangle_{\otimes^m \mathcal{E}^*(x, \xi)} dx d\xi. \tag{5-29}$$

The main idea of the proof is to reduce [\(5-28\)](#) to [\(5-29\)](#) by applying the coarea formula to a correctly chosen map $S^2_{\Delta} M \rightarrow SM$. More precisely, consider the following map $\Psi : \mathcal{E} \rightarrow S^2_{\Delta} \mathbb{H}^{n+1}$: for $(x, \xi) \in S\mathbb{H}^{n+1}$ and $\eta \in \mathcal{E}(x, \xi)$, define $\Psi(x, \xi, \eta) := (y, \eta_-, \eta_+)$, with

$$\begin{pmatrix} y \\ \eta_- \\ \eta_+ \end{pmatrix} = A(|\eta|^2) \begin{pmatrix} x \\ \xi \\ \eta \end{pmatrix}, \quad A(s) = \begin{pmatrix} \sqrt{s+1} & 0 & 1 \\ \frac{s}{\sqrt{s+1}} & \frac{1}{\sqrt{s+1}} & 1 \\ -\frac{s}{\sqrt{s+1}} & \frac{1}{\sqrt{s+1}} & -1 \end{pmatrix}.$$

Note that, with $|\eta|$ denoting the Riemannian length of η (that is, $|\eta|^2 = -\langle \eta, \eta \rangle_M$),

$$\Phi_{\pm}(y, \eta_{\pm}) = \frac{\Phi_{\pm}(x, \xi)}{\sqrt{1 + |\eta|^2}}, \quad B_{\pm}(y, \eta_{\pm}) = B_{\pm}(x, \xi), \quad |\eta_+ + \eta_-| = \frac{2}{\sqrt{1 + |\eta|^2}}.$$

Also,

$$\det A(s) = -\frac{2}{s+1}, \quad A(s)^{-1} = \begin{pmatrix} \sqrt{s+1} & -\frac{1}{2}\sqrt{s+1} & \frac{1}{2}\sqrt{s+1} \\ 0 & \frac{1}{2}\sqrt{s+1} & \frac{1}{2}\sqrt{s+1} \\ -s & \frac{1}{2}(s+1) & -\frac{1}{2}(s+1) \end{pmatrix}.$$




Figure 4. (a) The map $\Psi : (x, \xi, \eta) \mapsto (y, \eta_-, \eta_+)$. (b) The vectors $\mathcal{A}_\pm(x, \xi)\zeta_\pm$ (which are equal in the case drawn) and $\mathcal{A}_\pm(y, \eta_\pm)\zeta_\pm$.

The map Ψ is a diffeomorphism; the inverse is given by the formulas

$$x = \frac{2y + \eta_+ - \eta_-}{|\eta_+ + \eta_-|}, \quad \xi = \frac{\eta_+ + \eta_-}{|\eta_+ + \eta_-|}, \quad \eta = \frac{2(\eta_- - \eta_+) - |\eta_+ - \eta_-|^2 y}{|\eta_+ + \eta_-|^2}.$$

The map Ψ^{-1} can be visualized as follows (see Figure 4(a)): given (y, η_-, η_+) , the corresponding tangent vector (x, ξ) is the closest to y on the geodesic going from $v_- = B_-(y, \eta_-)$ to $v_+ = B_+(y, \eta_+)$ and the vector η measures both the distance between x and y and the direction of the geodesic from x to y . The exceptional set $\{\eta_+ + \eta_- = 0\}$ corresponds to $|\eta| = \infty$.

A calculation using (3-31) shows that, for $\zeta_\pm \in T_{B_\pm(x, \xi)}\mathbb{S}^n$,

$$\mathcal{A}_\pm(y, \eta_\pm)\zeta_\pm = \mathcal{A}_\pm(x, \xi)\zeta_\pm + \frac{(\mathcal{A}_\pm(x, \xi)\zeta_\pm) \cdot \eta}{\sqrt{1 + |\eta|^2}}(x \pm \xi).$$

Here, \cdot stands for the Riemannian inner product on \mathcal{E} , which is equal to $-\langle \cdot, \cdot \rangle_M$ restricted to \mathcal{E} . Then (see Figure 4(b))

$$\begin{aligned} & (\mathcal{A}_+(y, \eta_+)\zeta_+) \cdot (\mathcal{A}_-(y, \eta_-)\zeta_-) \\ &= (\mathcal{A}_+(x, \xi)\zeta_+) \cdot (\mathcal{A}_-(x, \xi)\zeta_-) - \frac{2}{1 + |\eta|^2} ((\mathcal{A}_+(x, \xi)\zeta_+) \cdot \eta) ((\mathcal{A}_-(x, \xi)\zeta_-) \cdot \eta) \\ &= (\mathcal{C}_\eta(\mathcal{A}_+(x, \xi)\zeta_+)) \cdot (\mathcal{A}_-(x, \xi)\zeta_-), \end{aligned}$$

where $\mathcal{C}_\eta : \mathcal{E}(x, \xi) \rightarrow \mathcal{E}(x, \xi)$ is given by

$$\mathcal{C}_\eta(\tilde{\eta}) = \tilde{\eta} - \frac{2}{1 + |\eta|^2} (\tilde{\eta} \cdot \eta)\eta.$$

We can similarly define $\mathcal{C}_\eta^* : \mathcal{E}(x, \xi)^* \rightarrow \mathcal{E}(x, \xi)^*$. Then, for $\zeta_\pm \in \otimes^m T_{B_\pm(x, \xi)}^*\mathbb{S}^n$,

$$\begin{aligned} & \langle \otimes^m (\mathcal{A}_+^{-1}(y, \eta_+)^T)\zeta_+, \otimes^m (\mathcal{A}_-^{-1}(y, \eta_-)^T)\zeta_- \rangle_{\otimes^m T_{\mathbb{H}^{n+1}}^*} \\ &= \langle \otimes^m \mathcal{C}_\eta^* \otimes^m (\mathcal{A}_+^{-1}(x, \xi)^T)\zeta_+, \otimes^m (\mathcal{A}_-^{-1}(x, \xi)^T)\zeta_- \rangle_{\otimes^m \mathcal{E}^*(x, \xi)}. \end{aligned} \quad (5-30)$$

The Jacobian of Ψ with respect to naturally arising volume forms on \mathcal{E} and $S^2_\Delta \mathbb{H}^{n+1}$ is given by (see [Appendix A2](#) for the proof)

$$J_\Psi(x, \xi, \eta) = 2^n (1 + |\eta|^2)^{-n}. \tag{5-31}$$

Now, Ψ is equivariant under G , therefore it descends to a diffeomorphism

$$\Psi : \mathcal{E}_M \rightarrow S^2_\Delta M, \quad \mathcal{E}_M := \{(x, \xi, \eta) \mid (x, \xi) \in SM, \eta \in \mathcal{E}(x, \xi)\}.$$

Using [Lemma 5.6](#) and (5-30), we calculate, for $(x, \xi, \eta) \in \mathcal{E}_M$ and $(y, \eta_-, \eta_+) = \Psi(x, \xi, \eta)$,

$$\langle v(y, \eta_-), \overline{v^*(y, \eta_+)} \rangle_{\otimes^m T^*_y M} = (1 + |\eta|^2)^{-\lambda} \langle \otimes^m \mathcal{C}_\eta^* v(x, \xi), \overline{v^*(x, \xi)} \rangle_{\otimes^m \mathcal{E}^*(x, \xi)}. \tag{5-32}$$

We would now like to plug this expression into (5-28), make the change of variables from (y, η_-, η_+) to (x, ξ, η) , and integrate η out, obtaining a multiple of (5-29). However, this is not directly possible because (i) the integral in η typically diverges and (ii) since the expression integrated in (5-28) is a distribution, one cannot simply replace $S^2 M$ by $S^2_\Delta M$ in the integral.

We will instead use the asymptotic behavior of both integrals as one approaches the set $\{\eta_+ + \eta_- = 0\}$, and Hadamard regularization in η in the (x, ξ, η) variables. For that, fix $\chi \in C^\infty_0(\mathbb{R})$ such that $\chi = 1$ near 0, and define, for $\varepsilon > 0$,

$$\chi_\varepsilon(y, \eta_-, \eta_+) = \chi(\varepsilon |\eta(y, \eta_-, \eta_+)|),$$

where $\eta(y, \eta_-, \eta_+)$ is the corresponding component of Ψ^{-1} ; we can write

$$\chi_\varepsilon(y, \eta_-, \eta_+) = \chi\left(\varepsilon \frac{|\eta_+ - \eta_-|}{|\eta_+ + \eta_-|}\right).$$

Then $\chi_\varepsilon \in \mathcal{D}'(S^2 M)$. In fact, χ_ε is supported inside $S^2_\Delta M$; by making the change of variables $(y, \eta_-, \eta_+) = \Psi(x, \xi, \eta)$ and, using (5-31) and (5-32), we get

$$\begin{aligned} & \int_{S^2 M} \chi_\varepsilon(y, \eta_-, \eta_+) \langle v(y, \eta_-), \overline{v^*(y, \eta_+)} \rangle_{\otimes^m T^*_y M} dy d\eta_- d\eta_+ \\ &= 2^n \int_{\mathcal{E}_M} \chi(\varepsilon |\eta|) (1 + |\eta|^2)^{-\lambda-n} \langle \otimes^m \mathcal{C}_\eta^* v(x, \xi), \overline{v^*(x, \xi)} \rangle_{\otimes^m \mathcal{E}^*(x, \xi)} dx d\xi d\eta. \end{aligned} \tag{5-33}$$

By [Lemma A.4](#), (5-33) has the asymptotic expansion

$$2^n \pi^{n/2} \frac{\Gamma(\frac{1}{2}n + \lambda)}{(n + \lambda + m - 1)\Gamma(n - 1 + \lambda)} \langle v, v^* \rangle_{L^2(SM; \otimes^m \mathcal{E}^*)} + \sum_{0 \leq j \leq -\operatorname{Re} \lambda - n/2} c_j \varepsilon^{n+2\lambda+2j} + o(1) \tag{5-34}$$

for some constants c_j .

It remains to prove the following asymptotic expansion as $\varepsilon \rightarrow 0$:

$$\int_{S^2 M} (1 - \chi_\varepsilon(y, \eta_-, \eta_+)) \langle v(y, \eta_-), \overline{v^*(y, \eta_+)} \rangle_{\otimes^m T^*_y M} dy d\eta_- d\eta_+ \sim \sum_{j=0}^\infty c'_j \varepsilon^{n+2\lambda+2j}, \tag{5-35}$$

where the c'_j are some constants. Indeed, $\langle f, f^* \rangle_{L^2(M; \otimes^m T^*_M)}$ is equal to the sum of (5-33) and (5-35); since (5-35) does not have a constant term, $\langle f, f^* \rangle$ is equal to the constant term in the expansion (5-34).

To show (5-35), we use the dilation vector field $\eta \cdot \partial_\eta$ on \mathcal{E} , which under Ψ becomes the following vector field on $S^2_\Delta M$ extending smoothly to $S^2 M$:

$$L_{(y, \eta_-, \eta_+)} = \left(\frac{1}{2}(\eta_- - \eta_+), \frac{1}{4}|\eta_+ - \eta_-|^2 y - \frac{1}{2}\eta_+ + \frac{1}{2}(\eta_- \cdot \eta_+)\eta_-, -\frac{1}{4}|\eta_+ - \eta_-|^2 y - \frac{1}{2}\eta_- + \frac{1}{2}(\eta_- \cdot \eta_+)\eta_+ \right).$$

The vector field L is tangent to the submanifold $\{\eta_+ + \eta_- = 0\}$; in fact,

$$L(|\eta_+ - \eta_-|^2) = -L(|\eta_+ + \eta_-|^2) = \frac{1}{2}|\eta_+ - \eta_-|^2 \cdot |\eta_+ + \eta_-|^2.$$

We can then compute (following the identity $L|\eta| = |\eta|$)

$$L\left(\frac{|\eta_+ - \eta_-|}{|\eta_+ + \eta_-|}\right) = \frac{|\eta_+ - \eta_-|}{|\eta_+ + \eta_-|} \quad \text{on } S^2_\Delta M.$$

Using the (x, ξ, η) coordinates and (5-31), we can compute the divergence of L with respect to the standard volume form on $S^2 M$:

$$\text{Div } L = n(\eta_+ \cdot \eta_-).$$

Moreover, $B_\pm(y, \eta_\pm)$ are constant along the trajectories of L , and

$$L(\Phi_\pm(y, \eta_\pm)) = -\frac{1}{4}|\eta_+ - \eta_-|^2 \Phi_\pm(y, \eta_\pm).$$

We also use (3-31) to calculate, for $\zeta_\pm \in T_{B_\pm(y, \eta_\pm)} \mathbb{S}^n$,

$$\begin{aligned} L((\mathcal{A}_+(y, \eta_+)\zeta_+) \cdot (\mathcal{A}_-(y, \eta_-)\zeta_-)) &= ((\mathcal{A}_+(y, \eta_+)\zeta_+) \cdot \eta_-)((\mathcal{A}_-(y, \eta_-)\zeta_-) \cdot \eta_+), \\ L((\mathcal{A}_\pm(y, \eta_\pm)\zeta_\pm) \cdot \eta_\mp) &= (\eta_+ \cdot \eta_-)((\mathcal{A}_\pm(y, \eta_\pm)\zeta_\pm) \cdot \eta_\mp). \end{aligned}$$

Combining these identities and using Lemma 5.6, we get

$$(L + \frac{1}{2}\lambda|\eta_+ - \eta_-|^2)\langle v(y, \eta_-), \overline{v^*(y, \eta_+)} \rangle_{\otimes^m T_y^* M} = m\langle \iota_{\eta_+} v(y, \eta_-), \iota_{\eta_-} \overline{v^*(y, \eta_+)} \rangle_{\otimes^{m-1} T_y^* M}. \quad (5-36)$$

Integrating by parts, we find

$$\begin{aligned} \varepsilon \partial_\varepsilon \int_{S^2 M} (1 - \chi_\varepsilon(y, \eta_-, \eta_+)) \langle v(y, \eta_-), \overline{v^*(y, \eta_+)} \rangle_{\otimes^m T_y^* M} dy d\eta_- d\eta_+ \\ = \int_{S^2 M} L(1 - \chi_\varepsilon(y, \eta_-, \eta_+)) \langle v(y, \eta_-), \overline{v^*(y, \eta_+)} \rangle_{\otimes^m T_y^* M} dy d\eta_- d\eta_+ \\ = \int_{S^2 M} \left(\frac{1}{2}\lambda|\eta_+ - \eta_-|^2 - n(\eta_+ \cdot \eta_-) \right) (1 - \chi_\varepsilon(y, \eta_-, \eta_+)) \langle v(y, \eta_-), \overline{v^*(y, \eta_+)} \rangle_{\otimes^m T_y^* M} dy d\eta_- d\eta_+ \\ \quad - m \int_{S^2 M} (1 - \chi_\varepsilon(y, \eta_-, \eta_+)) \langle \iota_{\eta_+} v(y, \eta_-), \iota_{\eta_-} \overline{v^*(y, \eta_+)} \rangle_{\otimes^{m-1} T_y^* M} dy d\eta_- d\eta_+. \end{aligned}$$

Arguing similarly, we see that if, for integers $0 \leq r \leq m$, $p \geq 0$, we put

$$I_{r,p}(\varepsilon) := \int_{S^2 M} |\eta_- + \eta_+|^{2p} (1 - \chi_\varepsilon(y, \eta_-, \eta_+)) \langle \iota_{\eta_+}^r v(y, \eta_-), \iota_{\eta_-}^r \overline{v^*(y, \eta_+)} \rangle_{\otimes^{m-r} T_y^* M} dy d\eta_- d\eta_+,$$

then $(\varepsilon\partial_\varepsilon - 2\lambda - n - 2(r + p))I_{r,p}(\varepsilon)$ is a finite linear combination of $I_{r',p'}(\varepsilon)$, where $r' \geq r$, $p' \geq p$, and $(r', p') \neq (r, p)$. For example, the calculation above shows that

$$(\varepsilon\partial_\varepsilon - 2\lambda - n)I_{0,0}(\varepsilon) = -\frac{1}{2}(\lambda + n)I_{0,1}(\varepsilon) - mI_{1,0}(\varepsilon).$$

Moreover, if N is fixed and p is large enough depending on N , then $I_{r,p}(\varepsilon) = \mathcal{O}(\varepsilon^N)$; to see this, note that $I_{r,p}(\varepsilon)$ is bounded by some fixed C^∞ -seminorm of $|\eta_- + \eta_+|^{2p}(1 - \chi_\varepsilon(y, \eta_-, \eta_+))$. It follows that, if N is fixed and \tilde{N} is large depending on N , then

$$\left(\prod_{j=0}^{\tilde{N}} (\varepsilon\partial_\varepsilon - 2\lambda - n - 2j) \right) I_{0,0}(\varepsilon) = \mathcal{O}(\varepsilon^N),$$

which implies the existence of the decomposition (5-35) and finishes the proof. □

6. Properties of the Laplacian

In this section, we introduce the Laplacian and study its basic properties (Section 6A). We then give formulas for the Laplacian on symmetric tensors in the half-plane model (Section 6B), which will be the basis for the analysis of the following sections. Using these formulas, we study the Poisson kernel and in particular prove Lemma 5.8 and the injectivity of the Poisson kernel (Section 6C).

6A. Definition and Bochner identity. The Levi-Civita connection associated to the hyperbolic metric g_H is the operator

$$\nabla : \mathcal{C}^\infty(\mathbb{H}^{n+1}, T\mathbb{H}^{n+1}) \rightarrow \mathcal{C}^\infty(\mathbb{H}^{n+1}, T^*\mathbb{H}^{n+1} \otimes T\mathbb{H}^{n+1}),$$

which induces a natural covariant derivative, still denoted ∇ , on sections of $\otimes^m T^*\mathbb{H}^{n+1}$. We can work in the ball model \mathbb{B}^{n+1} and use the 0-tangent structure (see Section 3F), and nabla can be viewed as a differential operator of order 1:

$$\nabla : \mathcal{C}^\infty(\bar{\mathbb{B}}^{n+1}; \otimes^m({}^0T^*\bar{\mathbb{B}}^{n+1})) \rightarrow \mathcal{C}^\infty(\bar{\mathbb{B}}^{n+1}, \otimes^{m+1}({}^0T^*\bar{\mathbb{B}}^{n+1})).$$

We denote by ∇^* its adjoint with respect to the L^2 scalar product, called the *divergence*; it is given by $\nabla^*u = -\mathcal{T}(\nabla u)$, where \mathcal{T} denotes the trace; see Section 4A. Define the rough Laplacian acting on $\mathcal{C}^\infty(\bar{\mathbb{B}}^{n+1}; \otimes^m({}^0T^*\bar{\mathbb{B}}^{n+1}))$ by

$$\Delta := \nabla^*\nabla; \tag{6-1}$$

this operator maps symmetric tensors to symmetric tensors. It also extends to $\mathcal{D}'(\mathbb{B}^{n+1}; \otimes_S^m({}^0T^*\bar{\mathbb{B}}^{n+1}))$ by duality. The operator Δ commutes with \mathcal{T} and \mathcal{I} :

$$\Delta\mathcal{T}(u) = \mathcal{T}(\Delta u) \quad \text{and} \quad \Delta\mathcal{I}(u) = \mathcal{I}(\Delta u) \tag{6-2}$$

for all $u \in \mathcal{D}'(\mathbb{B}^{n+1}; \otimes_S^m({}^0T^*\bar{\mathbb{B}}^{n+1}))$.

There is another natural operator given by

$$\Delta_D = D^*D$$

if

$$D : \mathcal{C}^\infty(\bar{\mathbb{B}}^{n+1}; \otimes_S^m({}^0T^*\bar{\mathbb{B}}^{n+1})) \rightarrow \mathcal{C}^\infty(\bar{\mathbb{B}}^{n+1}; \otimes_S^{m+1}({}^0T^*\bar{\mathbb{B}}^{n+1}))$$

is defined by $D := \mathcal{S} \circ \nabla$, where \mathcal{S} is the symmetrization defined by (4-1), and $D^* = \nabla^*$ is the formal adjoint. There is a Bochner–Weitzenböck formula relating Δ and Δ_D , and, using that the curvature is constant, we have on trace-free symmetric tensors of order m , by [Dairbekov and Sharafutdinov 2010, Lemma 8.2],

$$\Delta_D = \frac{1}{m+1}(mDD^* + \Delta + m(m+n-1)). \tag{6-3}$$

In particular, since $|\mathcal{S}\nabla u|^2 \leq |\nabla u|^2$ pointwise by the fact that \mathcal{S} is an orthogonal projection, we see that, for u smooth and compactly supported, $\|Du\|_{L^2}^2 \leq \|\nabla u\|_{L^2}^2$ and thus, for $m \geq 1$, $u \in \mathcal{C}_0^\infty(\mathbb{H}^{n+1}; \otimes_S^m(T^*\mathbb{H}^{n+1}))$, and $\mathcal{T}u = 0$,

$$\langle \Delta u, u \rangle_{L^2} \geq (m+n-1)\|u\|^2. \tag{6-4}$$

Since the Bochner identity is local, the same inequality clearly descends to cocompact quotients $\Gamma \backslash \mathbb{H}^{n+1}$ (where Δ is selfadjoint and has compact resolvent by standard theory of elliptic operators, as its principal part is given by the scalar Laplacian), and this implies:

Lemma 6.1. *The spectrum of Δ acting on trace-free symmetric tensors of order $m \geq 1$ on hyperbolic compact manifolds of dimension $n + 1$ is bounded below by $m + n - 1$.*

We finally define

$$E^{(m)} := \otimes_S^m({}^0T^*\bar{\mathbb{B}}^{n+1}) \cap \ker \mathcal{T} \tag{6-5}$$

to be the bundle of trace-free symmetric m -cotensors over the ball model of hyperbolic space.

6B. Laplacian in the half-plane model. We now give concrete formulas concerning the Laplacian on symmetric tensors in the half-space model \mathbb{U}^{n+1} (see (3-4)). We fix $\nu \in \mathbb{S}^n$ and map \mathbb{B}^{n+1} to \mathbb{U}^{n+1} by a composition of a rotation of \mathbb{B}^{n+1} and the map (3-5); the rotation is chosen so that ν is mapped to $0 \in \bar{\mathbb{U}}^{n+1}$ and $-\nu$ is mapped to infinity.

The 0-cotangent and tangent bundles ${}^0T^*\bar{\mathbb{B}}^{n+1}$ and ${}^0T\bar{\mathbb{B}}^{n+1}$ pull back to the half-space; we denote them ${}^0T^*\mathbb{U}^{n+1}$ and ${}^0T\mathbb{U}^{n+1}$. The coordinates on \mathbb{U}^{n+1} are $(z_0, z) \in \mathbb{R}^+ \times \mathbb{R}^n$ and $z = (z_1, \dots, z_n)$. We use the following orthonormal bases of ${}^0T\mathbb{U}^{n+1}$ and ${}^0T^*\mathbb{U}^{n+1}$:

$$Z_i = z_0 \partial_{z_i} \quad \text{and} \quad Z_i^* = \frac{dz_i}{z_0}, \quad 0 \leq i \leq n.$$

Note that in the compactification $\bar{\mathbb{B}}^{n+1}$ this basis is smooth only on $\bar{\mathbb{B}}^{n+1} \setminus \{-\nu\}$.

Let $\mathcal{A} := \{1, \dots, n\}$. We can decompose the vector bundle $\otimes_S^m({}^0T^*\mathbb{U}^{n+1})$ into an orthogonal direct sum

$$\otimes_S^m({}^0T^*\mathbb{U}^{n+1}) = \bigoplus_{k=0}^m E_k^{(m)}, \quad E_k^{(m)} = \text{span}(\mathcal{S}((Z_0^*)^{\otimes k} \otimes Z_I^*)_{I \in \mathcal{A}^{m-k}}),$$

and we let π_i be the orthogonal projection onto $E_i^{(m)}$. Now, each tensor $u \in \otimes_S^m({}^0T^*\mathbb{U}^{n+1})$ can be decomposed as $u = \sum_{i=0}^m u_i$ with $u_i = \pi_i(u) \in E_i^{(m)}$ which we can write as

$$u = \sum_{i=0}^m u_i, \quad u_i = \mathcal{S}((Z_0^*)^{\otimes i} \otimes u'_i), \quad u'_i \in E_0^{(m-i)}. \tag{6-6}$$

We can therefore identify $E_k^{(m)}$ with $E_0^{(m-k)}$ and view $E^{(m)}$ as a direct sum $E^{(m)} = \bigoplus_{k=0}^m E_0^{(m-k)}$. The trace-free condition, $\mathcal{T}(u) = 0$, is equivalent to the relations

$$\mathcal{T}(u'_r) = -\frac{(r+2)(r+1)}{(m-r)(m-r-1)} u'_{r+2}, \quad 0 \leq r \leq m-2, \tag{6-7}$$

and, in particular, all u_i are determined by u_0 and u_1 by iterating the trace map \mathcal{T} . The u'_i are related to the elements in the decomposition (4-5) of u_0 and u_1 viewed as a symmetric m -cotensor on the bundle $(Z_0)^\perp$ using the metric $z_0^{-2}h = \sum_i Z_i^* \otimes Z_i^*$. We see that a nonzero trace-free tensor u on \mathbb{U}^{n+1} must have a nonzero u_0 or u_1 component.

The Koszul formula gives us, for $i, j \geq 1$,

$$\nabla_{Z_i} Z_j = \delta_{ij} Z_0, \quad \nabla_{Z_0} Z_j = 0, \quad \nabla_{Z_i} Z_0 = -Z_i, \quad \nabla_{Z_0} Z_0 = 0, \tag{6-8}$$

which implies

$$\nabla Z_0^* = -\sum_{j=1}^n Z_j^* \otimes Z_j^* = -\frac{h}{z_0^2}, \quad \nabla Z_j^* = Z_j^* \otimes Z_0^*. \tag{6-9}$$

We shall use the following notations: If Π_m denotes the set of permutations of $\{1, \dots, m\}$, we write $\sigma(I) := (i_{\sigma(1)}, \dots, i_{\sigma(m)})$ if $\sigma \in \Pi_m$. If $S = S_1 \otimes \dots \otimes S_\ell$ is a tensor in $\otimes^\ell({}^0T^*\mathbb{U}^{n+1})$, we denote by $\tau_{i \leftrightarrow j}(S)$ the tensor obtained by permuting S_i with S_j in S , and by $\rho_{i \rightarrow V}(S)$ the operation of replacing S_i by $V \in {}^0T^*\mathbb{U}^{n+1}$ in S .

The Laplacian and ∇^* acting on $E_0^{(m)}$ and $E_1^{(m)}$. We start by computing the action of Δ on sections of $E_0^{(m)}$ and $E_1^{(m)}$, and we will later deduce from this computation the action on $E_k^{(m)}$. Let us consider the tensor $Z_I^* := Z_{i_1}^* \otimes \dots \otimes Z_{i_m}^* \in E_0^{(m)}$, where $I = (i_1, \dots, i_m) \in \mathcal{A}^m$ and $Z_{\sigma(I)}^* := Z_{i_{\sigma(1)}}^* \otimes \dots \otimes Z_{i_{\sigma(m)}}^*$. The symmetrization of Z_I^* is given by $\mathcal{S}(Z_I^*) = (1/m!) \sum_{\sigma \in \Pi_m} Z_{\sigma(I)}^*$ and those elements form a basis of the space $E_0^{(m)}$ when I ranges over all combinations of m -tuples in $\mathcal{A} = \{1, \dots, n\}$.

Lemma 6.2. *Let $u_0 = \sum_{I \in \mathcal{A}^m} f_I \mathcal{S}(Z_I^*)$ with $f_I \in C^\infty(\mathbb{U}^{n+1})$. Then one has*

$$\Delta u_0 = \sum_{I \in \mathcal{A}^m} ((\Delta + m) f_I) \mathcal{S}(Z_I^*) + 2m \mathcal{S}(\nabla^* u_0 \otimes Z_0^*) + m(m-1) \mathcal{S}(\mathcal{T}(u_0) \otimes Z_0^* \otimes Z_0^*), \tag{6-10}$$

while, denoting $d_z f_I = \sum_{i=1}^n Z_i(f_I) Z_i^*$, the divergence is given by

$$\nabla^* u_0 = -(m-1) \mathcal{S}(\mathcal{T}(u_0) \otimes Z_0^*) - \sum_{I \in \mathcal{A}^m} \iota_{d_z f_I} \mathcal{S}(Z_I^*). \tag{6-11}$$

Proof. Using (6-9), we compute

$$\nabla(f_I \mathcal{S}(Z_I^*)) = \sum_{i=0}^n (Z_i f_I)(z) Z_i^* \otimes \mathcal{S}(Z_I^*) + \frac{f_I(z)}{m!} \sum_{k=1}^m \sum_{\sigma \in \Pi_m} \tau_{1 \leftrightarrow k+1}(Z_0^* \otimes Z_{\sigma(I)}^*).$$

Then, taking the trace of $\nabla(f_I \mathcal{S}(Z_I^*))$ gives

$$\begin{aligned} \nabla^*(f_I \mathcal{S}(Z_I^*)) &= -\frac{f_I}{m!} \sum_{k=2}^m \sum_{\sigma \in \Pi_m} \delta_{i_{\sigma(1)}, i_{\sigma(k)}} \rho_{k-1 \rightarrow Z_0^*}(Z_{i_{\sigma(2)}}^* \otimes \cdots \otimes Z_{i_{\sigma(m)}}^*) \\ &\quad - \sum_{i=1}^n (Z_i f_I) \frac{1}{m!} \sum_{\sigma \in \Pi_m} \delta_{i, i_{\sigma(1)}}(Z_{i_{\sigma(2)}}^* \otimes \cdots \otimes Z_{i_{\sigma(m)}}^*). \end{aligned} \quad (6-12)$$

We notice that $\mathcal{S}(\mathcal{T}(\mathcal{S}(Z_I^*)) \otimes Z_0^*)$ is given by

$$\mathcal{S}(\mathcal{T}(\mathcal{S}(Z_I^*)) \otimes Z_0^*) = \frac{1}{m!(m-1)} \sum_{\sigma \in \Pi_m} \sum_{k=1}^{m-1} \delta_{i_{\sigma(1)}, i_{\sigma(2)}} \tau_{1 \leftrightarrow k}(Z_0^* \otimes Z_{i_{\sigma(3)}}^* \otimes \cdots \otimes Z_{i_{\sigma(m)}}^*),$$

which implies (6-11). Let us now compute $\nabla^2(f_I \mathcal{S}(Z_I^*))$:

$$\begin{aligned} &\nabla^2(f_I \mathcal{S}(Z_I^*)) \\ &= \sum_{i,j=0}^n Z_j Z_i (f_I) Z_j^* \otimes Z_i^* \otimes \mathcal{S}(Z_I^*) - Z_0(f_I) z_0^{-2} h \otimes \mathcal{S}(Z_I^*) \\ &\quad + \sum_{j=1}^n Z_j (f_I) Z_j^* \otimes Z_0^* \otimes \mathcal{S}(Z_I^*) + \frac{Z_0(f_I)}{m!} \sum_{\sigma \in \Pi_m} \sum_{k=1}^m \tau_{1 \leftrightarrow k+2}(Z_0^* \otimes Z_0^* \otimes Z_{\sigma(I)}^*) \\ &\quad + \sum_{i=1}^n \frac{Z_i(f_I)}{m!} \sum_{\sigma \in \Pi_m} \sum_{k=1}^m \tau_{1 \leftrightarrow k+2}(Z_0^* \otimes Z_i^* \otimes Z_{\sigma(I)}^*) + \sum_{i=1}^n \frac{Z_i(f_I)}{m!} \sum_{\sigma \in \Pi_m} \sum_{k=1}^m \tau_{2 \leftrightarrow k+2}(Z_i^* \otimes Z_0^* \otimes Z_{\sigma(I)}^*) \\ &\quad + \frac{Z_0(f_I)}{m!} Z_0^* \otimes \sum_{k=1}^m \sum_{\sigma \in \Pi_m} \tau_{1 \leftrightarrow k+1}(Z_0^* \otimes Z_{\sigma(I)}^*) \\ &\quad - \frac{f_I}{m!} \sum_{j=1}^n Z_j^* \otimes \sum_{\sigma \in \Pi_m} \sum_{k=1}^m \tau_{1 \leftrightarrow k+1}(Z_j^* \otimes Z_{\sigma(I)}^*) + \frac{f_I}{m!} \sum_{k=1}^m \sum_{\substack{\ell=1 \\ \ell \neq k+1}}^{m+1} \tau_{1 \leftrightarrow \ell+1}(Z_0^* \otimes \tau_{1 \leftrightarrow k+1}(Z_0^* \otimes Z_{\sigma(I)}^*)). \end{aligned}$$

We then take the trace: the first line on the right-hand side has trace $-(\Delta f_I) \mathcal{S}(Z_I^*)$, the second and fourth lines have vanishing trace, the first term on the last line has trace $-m f_I \mathcal{S}(Z_I^*)$, the last term has trace

$$\frac{2f_I}{m!} \sum_{\sigma \in \Pi_m} \sum_{1 \leq k < \ell \leq m} \delta_{i_{\sigma(k)}, i_{\sigma(\ell)}} \rho_{k \rightarrow Z_0^*} \rho_{\ell \rightarrow Z_0^*}(Z_{\sigma(I)}^*), \quad (6-13)$$

and the third line has total trace

$$2 \sum_{i=1}^n \frac{Z_i(f_I)}{m!} \sum_{\sigma \in \Pi_m} \sum_{k=1}^m \delta_{i, i_{\sigma(k)}} \rho_{k \rightarrow Z_0^*}(Z_{\sigma(I)}^*). \quad (6-14)$$

Computing $\mathcal{S}(\mathcal{T}(\mathcal{S}(Z_I^*)) \otimes Z_0^* \otimes Z_0^*)$ gives

$$\begin{aligned} &\mathcal{S}(\mathcal{T}(\mathcal{S}(Z_I^*)) \otimes Z_0^* \otimes Z_0^*) \\ &= \frac{2}{m!m(m-1)} \sum_{1 \leq k < \ell \leq m} \sum_{\sigma \in \Pi_m} \delta_{i_{\sigma(1)}, i_{\sigma(2)}} \tau_{1 \leftrightarrow k+2} \tau_{2 \leftrightarrow \ell+2} (Z_0^* \otimes Z_0^* \otimes Z_{i_{\sigma(3)}}^* \otimes \cdots \otimes Z_{i_{\sigma(m)}}^*); \end{aligned}$$

therefore the term (6-13) can be simplified to

$$m(m-1) f_I \mathcal{S}(\mathcal{T}(\mathcal{S}(Z_I^*)) \otimes Z_0^* \otimes Z_0^*).$$

Similarly, to simplify (6-14), we compute

$$\begin{aligned} &\mathcal{S}(\nabla^*(f_I \mathcal{S}(Z_I^*)) \otimes Z_0^*) \\ &= -(m-1) \mathcal{S}(\mathcal{T}(f_I \mathcal{S}(Z_I^*)) \otimes Z_0^* \otimes Z_0^*) - \sum_{i=1}^n (Z_i f_I) \frac{1}{m!m} \sum_{k=1}^m \sum_{\sigma \in \Pi_m} \delta_{i, i_{\sigma(1)}} \tau_{1 \leftrightarrow k} (Z_0^* \otimes Z_{i_{\sigma(2)}}^* \otimes \cdots \otimes Z_{i_{\sigma(m)}}^*), \end{aligned}$$

so that

$$\begin{aligned} &2 \sum_{i=1}^n \frac{Z_i(f_I)}{m!} \sum_{\sigma \in \Pi_m} \sum_{k=1}^m \delta_{i, i_{\sigma(k)}} \rho_{k \rightarrow Z_0^*} (Z_{\sigma(I)}^*) \\ &= -2m \mathcal{S}(\nabla^*(f_I \mathcal{S}(Z_I^*)) \otimes Z_0^*) - 2m(m-1) \mathcal{S}(\mathcal{T}(f_I \mathcal{S}(Z_I^*)) \otimes Z_0^* \otimes Z_0^*), \end{aligned}$$

and this achieves the proof of (6-10). □

A similarly tedious calculation, omitted here, yields:

Lemma 6.3. *Let $u_1 = \mathcal{S}(Z_0^* \otimes u'_1)$, $u'_1 = \sum_{J \in \mathcal{A}^{m-1}} g_J \mathcal{S}(Z_J^*)$ with $g_J \in \mathcal{C}^\infty(\mathbb{U}^{n+1})$; then the $E_0^{(m)} \oplus E_1^{(m)}$ components of the Laplacian of u_1 are*

$$\Delta u_1 = \sum_{J \in \mathcal{A}^{m-1}} ((\Delta + n + 3(m-1))g_J) \mathcal{S}(Z_0^* \otimes Z_J^*) + 2 \sum_{J \in \mathcal{A}^{m-1}} \mathcal{S}(d_z g_J \otimes Z_J^*) + \text{Ker}(\pi_0 + \pi_1) \quad (6-15)$$

and the $E_0^{(m)} \oplus E_1^{(m)}$ components of divergence of u_1 are

$$\nabla^* u_1 = \frac{1}{m} \sum_{J \in \mathcal{A}^{m-1}} ((n+m-1)g_J - Z_0(g_J)) \mathcal{S}(Z_J^*) - \frac{m-1}{m} \sum_{J \in \mathcal{A}^{m-1}} \mathcal{S}(Z_0^* \otimes \iota_{d_z g_J} \mathcal{S}(Z_J^*)) + \text{Ker}(\pi_0 + \pi_1). \quad (6-16)$$

General formulas for Laplacian and divergence. Armed with Lemmas 6.2 and 6.3, we can show the following fact, which, together with (6-7), completely determines the Laplacian on trace-free symmetric tensors.

Lemma 6.4. *Assume that $u \in \mathcal{D}'(\mathbb{U}^{n+1}; \otimes_S^m T^* \mathbb{U}^{n+1})$ satisfies $\mathcal{T}(u) = 0$ and is written in the form (6-6).*

Let

$$u_0 = \sum_{I \in \mathcal{A}^m} f_I \mathcal{S}(Z_I^*), \quad u_1 = \sum_{J \in \mathcal{A}^{m-1}} g_J \mathcal{S}(Z_0 \otimes Z_J^*).$$

Then the projection of Δu onto $E_0^{(m)} \oplus E_1^{(m)}$ can be written

$$\pi_0(\Delta u) = \sum_{I \in \mathcal{I}^m} ((\Delta + m) f_I) \mathcal{S}(Z_I^*) + 2 \sum_{J \in \mathcal{I}^{m-1}} \mathcal{S}(d_z g_J \otimes Z_J^*) + m(m-1) \mathcal{S}(z_0^{-2} h \otimes \mathcal{T}(u_0)), \quad (6-17)$$

$$\begin{aligned} \pi_1(\Delta u) = & \sum_{J \in \mathcal{I}^{m-1}} ((\Delta + n + 3(m-1)) g_J) \mathcal{S}(Z_0^* \otimes Z_J^*) - 2m \sum_{I \in \mathcal{I}^m} \mathcal{S}(Z_0^* \otimes \iota_{d_z f_I} \mathcal{S}(Z_I^*)) \\ & + (m-1)(m-2) \mathcal{S}(Z_0^* \otimes z_0^{-2} h \otimes \mathcal{T}(u_1')) - 2m(m-1) \sum_{I \in \mathcal{I}^m} \mathcal{S}(Z_0^* \otimes d_z f_I \otimes \mathcal{T}(\mathcal{S}(Z_I^*))). \end{aligned} \quad (6-18)$$

Proof. First, it is easily seen from (6-9) that Δu_k is a section of $\bigoplus_{j=k-2}^{k+2} E_j^{(m)}$. From Lemmas 6.2 and 6.3, we have

$$\pi_0(\Delta(u_0 + u_1)) = \sum_{I \in \mathcal{I}^m} ((\Delta + m) f_I) \mathcal{S}(Z_I^*) + 2 \sum_{J \in \mathcal{I}^{m-1}} \mathcal{S}(d_z g_J \otimes Z_J^*). \quad (6-19)$$

Then, for u_2 , using $\mathcal{S}((Z_0^*)^{\otimes 2} \otimes u_2') = \mathcal{S}(g_H \otimes u_2') - \mathcal{S}(z_0^{-2} h \otimes u_2')$ and $\Delta \mathcal{I} = \mathcal{I} \Delta$,

$$\pi_0(\Delta u_2) = \pi_0(\mathcal{S}(z_0^{-2} h \otimes \Delta u_2')) - \pi_0(\Delta(\mathcal{S}(z_0^{-2} h \otimes u_2')))$$

and, writing $u_2' = -\frac{1}{2}m(m-1)\mathcal{T}(u_0)$ by (6-7), we obtain, using (6-10),

$$\pi_0(\Delta u_2) = m(m-1) \mathcal{S}(z_0^{-2} h \otimes \mathcal{T}(u_0)). \quad (6-20)$$

We therefore obtain (6-17).

Now we consider the projection on $E_1^{(m)}$ of the equation $(\Delta - s)T = 0$. We have, from (6-10),

$$\pi_1(\Delta u_0) = -2m \sum_{I \in \mathcal{I}^m} \mathcal{S}(Z_0^* \otimes \iota_{d_z f_I} \mathcal{S}(Z_I^*)),$$

where $\iota_{d_z f_I}$ means $\sum_{j=1}^n Z_j(f_I) \iota_{Z_j}$. Then, from (6-15),

$$\pi_1(\Delta u_1) = \sum_{J \in \mathcal{I}^{m-1}} ((\Delta + n + 3(m-1)) g_J) \mathcal{S}(Z_0^* \otimes Z_J^*).$$

Using again $\mathcal{S}((Z_0^*)^{\otimes 2} \otimes u_2') = \mathcal{S}(g_H \otimes u_2') - \mathcal{S}(z_0^{-2} h \otimes u_2')$ and $\Delta \mathcal{I} = \mathcal{I} \Delta$, (6-10) gives

$$\pi_1(\Delta u_2) = -2m(m-1) \sum_{I \in \mathcal{I}^m} \mathcal{S}(Z_0^* \otimes d_z f_I \otimes \mathcal{T}(\mathcal{S}(Z_I^*))).$$

Finally, we compute $\pi_1(\Delta u_3)$: using the computation (6-15), we get

$$\begin{aligned} \pi_1(\Delta u_3) &= \pi_1(\mathcal{S}(z_0^{-2} h \otimes \Delta \mathcal{S}(Z_0^* \otimes u_3')) - \pi_1(\Delta \mathcal{S}(Z_0^* \otimes z_0^{-2} h \otimes u_3')) \\ &= (m-1)(m-2) \mathcal{S}(Z_0^* \otimes z_0^{-2} h \otimes \mathcal{T}(u_1')). \end{aligned}$$

We conclude that $\pi_1(\Delta u)$ is given by (6-18). □

Similarly, we also have:

Lemma 6.5. *Let u be as in Lemma 6.4. Then the projection onto $E_0^{(m-1)} \oplus E_1^{(m-1)}$ of the divergence of u is given by*

$$\pi_0(\nabla^* u) = - \sum_{I \in \mathcal{A}^m} \iota_{d_z f_I} \mathcal{S}(Z_I^*) + \frac{1}{m} \sum_{J \in \mathcal{A}^{m-1}} ((n+m-1)g_J - Z_0(g_J)) \mathcal{S}(Z_J^*), \tag{6-21}$$

$$\pi_1(\nabla^* u) = (m-1) \sum_{I \in \mathcal{A}^m} (Z_0 f_I - (m+n-1)f_I) \mathcal{S}(\mathcal{T}(\mathcal{S}(Z_I^*)) \otimes Z_0^*) - \frac{m-1}{m} \sum_{J \in \mathcal{A}^{m-1}} \mathcal{S}(Z_0^* \otimes \iota_{d_z g_J} \mathcal{S}(Z_J^*)). \tag{6-22}$$

Proof. The π_0 part follows from (6-11) and (6-16). For the π_1 part, we also use (6-11) and (6-16) but we need to see the contribution from $\nabla^* u_2$ as well. For that, we write $u'_2 = -\frac{1}{2}m(m-1) \sum_{I \in \mathcal{A}^m} f_I \mathcal{T}(\mathcal{S}(Z_I^*))$, as before, and a direct calculation shows that

$$\pi_1(\nabla^* u_2) = (m-1) \sum_{I \in \mathcal{A}^m} (Z_0 f_I - (m+n-2)f_I) \mathcal{S}(\mathcal{T}(\mathcal{S}(Z_I^*)) \otimes Z_0^*),$$

implying the desired result. □

6C. Properties of the Poisson kernel. In this section, we study the Poisson kernel \mathcal{P}_λ^- defined by (5-17).

Pairing on the sphere. We start by proving the following formula:

Lemma 6.6. *Let $\lambda \in \mathbb{C}$ and $w \in \mathcal{D}'(\mathbb{S}^n; \otimes_S^m(T^*\mathbb{S}^n))$. Then*

$$\mathcal{P}_\lambda^- w(x) = \int_{\mathbb{S}^n} P(x, \nu)^{n+\lambda} (\otimes^m (\mathcal{A}_-^{-1}(x, \xi_-(x, \nu)))^T) w(\nu) dS(\nu),$$

where the map ξ_- is as defined in (3-20).

Proof. Making the change of variables $\xi = \xi_-(x, \nu)$ defined in (3-20), and using (3-21) and (3-22), we have

$$\begin{aligned} \mathcal{P}_\lambda^- w(x) &= \int_{S_x \mathbb{H}^{n+1}} \Phi_-(x, \xi)^\lambda (\otimes^m (\mathcal{A}_-^{-1}(x, \xi))^T) w(B_-(x, \xi)) dS(\xi) \\ &= \int_{\mathbb{S}^n} P(x, \nu)^{n+\lambda} (\otimes^m (\mathcal{A}_-^{-1}(x, \xi_-(x, \nu)))^T) w(\nu) dS(\nu), \end{aligned}$$

as required. □

Poisson maps to eigenstates. To show that $\mathcal{P}_\lambda^- w(x)$ is an eigenstate of the Laplacian, we use:

Lemma 6.7. *Assume that $w \in \mathcal{D}'(\mathbb{S}^n; \otimes^m(T^*\mathbb{S}^n))$ is the delta function centered at $e_1 = \partial_{x_1} \in \mathbb{S}^n$ with the value $e_{j_1+1}^* \otimes \dots \otimes e_{j_m+1}^*$, where $1 \leq j_1, \dots, j_m \leq n$. Then, under the identifications (3-2) and (3-5), we have*

$$\mathcal{P}_\lambda^- w(z_0, z) = z_0^{n+\lambda} Z_{j_1}^* \otimes \dots \otimes Z_{j_m}^*.$$

Proof. We first calculate

$$P(z, e_1) = z_0.$$

It remains to show the identity in the half-space model

$$\mathcal{A}_-^{-T}(z, \xi_-(z, \nu)) e_{j+1}^* = Z_j^*, \quad 1 \leq j \leq n. \tag{6-23}$$

One can verify (6-23) by a direct computation; since \mathcal{A}_- is an isometry, one can instead calculate the image of e_{j+1} under \mathcal{A}_- , and then apply to it the differentials of the maps ψ and ψ_1 defined in (3-2) and (3-5).

Another way to show (6-23) is to use the interpretation of \mathcal{A}_- as parallel transport to conformal infinity; see (3-35). Note that under the diffeomorphism $\psi_1 : \mathbb{B}^{n+1} \rightarrow \mathbb{U}^{n+1}$, $\nu = e_1$ is sent to infinity and geodesics terminating at ν to straight lines parallel to the z_0 axis. By (6-9), the covector field Z_j^* is parallel along these geodesics and orthogonal to their tangent vectors. It remains to verify that the limit of the field $\rho_0 Z_j^*$ along these geodesics as $z \rightarrow \infty$, considered as a covector in the ball model, is equal to e_{j+1}^* . \square

Proof of Lemma 5.8. It suffices to show that, for each $\nu \in \mathbb{S}^n$, if w is a delta function centered at ν with value some symmetric trace-free tensor in $\otimes_S^m T_\nu^* \mathbb{S}^n$, then

$$(\Delta + \lambda(n + \lambda) - m) \mathcal{P}_\lambda^- w = 0, \quad \nabla^* \mathcal{P}_\lambda^- w = 0, \quad \mathcal{T}(\mathcal{P}_\lambda^- w) = 0.$$

Since the group of symmetries G of \mathbb{H}^{n+1} acts transitively on \mathbb{S}^n , we may assume that $\nu = \partial_1$. Applying Lemma 6.7, we write in the upper half-plane model

$$\mathcal{P}_\lambda^- w = z_0^{n+\lambda} u_0, \quad u_0 \in E_0^{(m)}, \quad \mathcal{T}(u_0) = 0.$$

It immediately follows that $\mathcal{T}(\mathcal{P}_\lambda^- w) = 0$. To see the other two identities, it suffices to apply Lemma 6.2 together with the formula

$$\Delta z_0^{n+\lambda} = -\lambda(n + \lambda) z_0^{n+\lambda}.$$

Injectivity of Poisson. Notice that \mathcal{P}_λ^- is an analytic family of operators in λ . We define the set

$$\mathcal{R}_m = \begin{cases} -\frac{1}{2}n - \frac{1}{2}\mathbb{N}_0 & \text{if } n > 1 \text{ or } m = 0, \\ -\frac{1}{2}\mathbb{N}_0 & \text{if } n = 1 \text{ and } m > 0, \end{cases} \tag{6-24}$$

and we will prove that, if $\lambda \notin \mathcal{R}_m$ and $w \in \mathcal{D}'(\mathbb{S}^n; \otimes_S^m T^* \mathbb{S}^n)$ is trace-free, then $\mathcal{P}_\lambda^-(w)$ has a weak asymptotic expansion at the conformal infinity with the leading term given by a multiple of w , proving injectivity of \mathcal{P}_λ^- . We shall use the 0-cotangent bundle approach in the ball model and rewrite $\mathcal{A}_\pm^{-1}(x, \xi_\pm(x, \nu))$ as the parallel transport $\tau(y', y)$ in ${}^0T\bar{\mathbb{B}}^{n+1}$ with $\psi(x) = y$ and $y' = \nu$, as explained in (3-35). Let $\rho \in C^\infty(\bar{\mathbb{B}}^{n+1})$ be a smooth boundary defining function which satisfies $\rho > 0$ in \mathbb{B}^{n+1} , $|d\rho|_{\rho^2 g_H} = 1$ near $\mathbb{S}^n = \{\rho = 0\}$, where g_H is the hyperbolic metric on the ball. We can for example take the function $\rho = \rho_0$ defined in (3-34) and smooth it near the center $y = 0$ of the ball. Such a function is called a *geodesic boundary defining function* and induces a diffeomorphism

$$\theta : [0, \epsilon)_t \times \mathbb{S}^n \rightarrow \bar{\mathbb{B}}^{n+1} \cap \{\rho < \epsilon\}, \quad \theta(t, \nu) := \theta_t(\nu), \tag{6-25}$$

where θ_t is the flow at time t of the gradient $\nabla^{\rho^2 g_H} \rho$ of ρ (denoted also ∂_ρ) with respect to the metric $\rho^2 g_H$. For ρ given in (3-34), we have, for t small,

$$\theta(t, \nu) = \frac{2-t}{2+t} \nu, \quad \nu \in \mathbb{S}^n.$$

For a fixed geodesic boundary defining function ρ , one can identify, over the boundary \mathbb{S}^n of $\bar{\mathbb{B}}^{n+1}$, the bundle $T^*\mathbb{S}^n$ and $T\mathbb{S}^n$ with the bundles ${}^0T^*\mathbb{S}^n := {}^0T^*\bar{\mathbb{B}}^{n+1} \cap \ker \iota_{\rho\partial\rho}$ simply by the isomorphism $v \mapsto \rho^{-1}v$ (and we identify their duals $T\mathbb{S}^n$ and ${}^0T\mathbb{S}^n$ as well). Similarly, over \mathbb{S}^n , $E^{(m)} \cap \ker \iota_{\rho\partial\rho}$ identifies with $\otimes_S^m T^*\mathbb{S}^n \cap \ker \mathcal{T}$ by the map $v \mapsto \rho^{-m}v$. We can then view the Poisson operator as an operator

$$\mathcal{P}_\lambda^- : \mathcal{D}'(\mathbb{S}^n; E^{(m)} \cap \ker \iota_{\rho\partial\rho}) \rightarrow \mathcal{C}^\infty(\mathbb{B}^{n+1}; \otimes_S^m ({}^0T^*\bar{\mathbb{B}}^{n+1})).$$

Lemma 6.8. *Let $w \in \mathcal{D}'(\mathbb{S}^n; E^{(m)} \cap \ker \iota_{\rho_0\partial\rho_0})$ and assume that $\lambda \notin \mathcal{R}_m$. Then $\mathcal{P}_\lambda^-(w)$ has a weak asymptotic expansion at \mathbb{S}^n as follows: for each $v \in \mathbb{S}^n$, there exists a neighborhood $V_v \subset \bar{\mathbb{B}}^{n+1}$ of v and a boundary defining function $\rho = \rho_v$ such that, for any $\varphi \in \mathcal{C}^\infty(V_v \cap \mathbb{S}^n; \otimes_S^m ({}^0T\mathbb{S}^n))$, there exist $F_\pm \in \mathcal{C}^\infty([0, \epsilon))$ such that, for $t > 0$ small,*

$$\begin{aligned} \int_{\mathbb{S}^n} \langle \mathcal{P}_\lambda^-(w)(\theta(t, v)), \otimes^m(\tau(\theta(t, v), v)) \cdot \varphi(v) \rangle dS_\rho(v) \\ = \begin{cases} t^{-\lambda} F_-(t) + t^{n+\lambda} F_+(t), & \lambda \notin -\frac{1}{2}n + \mathbb{N}, \\ t^{-\lambda} F_-(t) + t^{n+\lambda} \log(t) F_+(t), & \lambda \in -\frac{1}{2}n + \mathbb{N}. \end{cases} \end{aligned} \tag{6-26}$$

using the product collar neighborhood (6-25) associated to ρ , and, moreover, one has

$$F_-(0) = C \frac{\Gamma(\lambda + \frac{1}{2}n)}{(\lambda + n + m - 1)\Gamma(\lambda + n - 1)} \langle e^{\lambda f} \cdot w, \varphi \rangle \tag{6-27}$$

for some $f \in \mathcal{C}^\infty(\mathbb{S}^n)$ satisfying $\rho = \frac{1}{4}e^f \rho_0 + \mathcal{O}(\rho)$ near $\rho = 0$ and $C \neq 0$ a constant depending only on n . Here dS_ρ is the Riemannian measure for the metric $(\rho^2 g_H)|_{\mathbb{S}^n}$ and the distributional pairing on \mathbb{S}^n is with respect to this measure.

Proof. First we split w into $w_1 + w_2$, where w_1 is supported near $v \in \mathbb{S}^n$ and w_2 is zero near v . For the case where w_2 has support at positive distance from the support of φ , we have, for any geodesic boundary defining function ρ , that

$$t \mapsto t^{-n-\lambda} \int_{\mathbb{S}^n} \langle \mathcal{P}_\lambda^-(w_2)(\theta(t, v)), \otimes^m(\tau(\theta(t, v), v)) \cdot \varphi(v) \rangle dS_\rho(v) \in \mathcal{C}^\infty([0, \epsilon));$$

this is a direct consequence of Lemma 6.6 and the following smoothness properties:

$$\begin{aligned} (y, v) \mapsto \log\left(\frac{P(\psi^{-1}(y), v)}{\rho(y)}\right) \in \mathcal{C}^\infty(\bar{\mathbb{B}}^{n+1} \times \mathbb{S}^n \setminus \text{diag}(\mathbb{S}^n \times \mathbb{S}^n)), \\ \tau(\cdot, \cdot) \in \mathcal{C}^\infty(\bar{\mathbb{B}}^{n+1} \times \bar{\mathbb{B}}^{n+1} \setminus \text{diag}(\mathbb{S}^n \times \mathbb{S}^n); {}^0T^*\bar{\mathbb{B}}^{n+1} \otimes {}^0T\bar{\mathbb{B}}^{n+1}). \end{aligned}$$

This reduces the consideration of the lemma to the case where w is w_1 , supported near v , and to simplify we shall keep the notation w instead of w_1 . We thus consider now w and φ to have support near v . For convenience of calculations and as we did before, we work in the half-space model $\mathbb{R}_{z_0}^+ \times \mathbb{R}_z^n$ by mapping v to $(z_0, z) = (0, 0)$ (using the composition of a rotation on the ball model with the map defined in (3-5)), and we choose a neighborhood V_v of v which is mapped to $z_0^2 + |z|^2 < 1$ in \mathbb{U}^{n+1} and choose the geodesic defining function $\rho = z_0$ (and thus $\theta(z_0, z) = (z_0, z)$). (See Figure 5.) The geodesic boundary defining




Figure 5. The covector $w(z')$, the vector $\varphi(z)$, and their parallel transports to (z_0, z) , viewed in the 0-bundles for the case $m = 1$.

function $\rho_0 = 2(1 - |y|)/(1 + |y|)$ in the ball equals

$$\rho_0(z_0, z) = \frac{4z_0}{1 + z_0^2 + |z|^2} \tag{6-28}$$

in the half-space model. The metric dS_ρ becomes the Euclidean metric dz on \mathbb{R}^n near 0 and w has compact support in \mathbb{R}^n . By (3-5) and (3-19), the Poisson kernel in these coordinates becomes

$$\tilde{P}(z_0, z; z') = e^{f(z')} P(z_0, z; z') \quad \text{with} \quad P(z_0, z; z') := \frac{z_0}{z_0^2 + |z - z'|^2}, \quad f(z') = \log(1 + |z'|^2),$$

where $z, z' \in \mathbb{R}^n$ and $z_0 > 0$. One has $\rho = \frac{1}{4}e^f \rho_0 + \mathcal{O}(\rho)$ near $\rho = 0$.

In [Guillarmou et al. 2010, Appendix], the parallel transport $\tau(z_0, z; 0, z')$ is computed for $z' \in \mathbb{R}^n$ in a neighborhood of 0: in the local orthonormal basis $Z_0 = z_0 \partial_{z_0}$, $Z_i = z_0 \partial_{z_i}$ of the bundle ${}^0T\mathbb{U}^{n+1}$, near ν , the matrix of $\tau(z_0, z; z') := \tau(z_0, z; 0, z')$ is given by

$$\begin{aligned} \tau_{00} &= 1 - 2P(z_0, z; z') \frac{|z - z'|^2}{z_0}, \\ \tau_{0i} &= -\tau_{i0} = -2z_0(z_i - z'_i) \frac{P(z_0, z; z')}{z_0}, \\ \tau_{ij} &= \delta_{ij} - 2P(z_0, z; z') \frac{(z_i - z'_i) \cdot (z_j - z'_j)}{z_0}. \end{aligned}$$

In particular, we see that $\tau(z_0, z; z)$ is the identity matrix in the basis $(Z_i)_i$ and thus $\tau(\theta(z_0, z), z)$ as well. We denote by $(Z_j^*)_j$ the dual basis to $(Z_j)_j$ as before.

Now, we use the correspondence between symmetric tensors and homogeneous polynomials to facilitate computations, as explained in Section 4A. To $\mathcal{S}(Z_I^*)$, we associate the polynomial on \mathbb{R}^n given by

$$P_I(x) = \mathcal{S}(Z_I^*) \left(\sum_{i=1}^n x_i Z_i, \dots, \sum_{i=1}^n x_i Z_i \right) = x_I,$$

where $x_I = \prod_{k=1}^m x_{i_k}$ if $I = (i_1, \dots, i_m)$. We denote by $\text{Pol}^m(\mathbb{R}^n)$ the space of homogeneous polynomials of degree m on \mathbb{R}^n and $\text{Pol}_0^m(\mathbb{R}^n)$ those which are harmonic (thus corresponding to trace-free symmetric

tensors in $E_0^{(m)}$. Then we can write $w = \sum_{\alpha} w_{\alpha} p_{\alpha}(x)$ for some $w_{\alpha} \in \mathcal{D}'(\mathbb{R}^n)$ supported near 0 and $p_{\alpha}(x) \in \text{Pol}_0^m(\mathbb{R}^n)$. Each $p_{\alpha}(x)$ composed with the linear map $\tau(z'; z_0, z)|_{Z_0^{\perp}}$ becomes the homogeneous polynomial in x

$$p_{\alpha}\left(x - 2(z - z')\langle z - z', x \rangle \cdot \frac{P(z_0, z; z')}{z_0}\right),$$

where $\langle \cdot, \cdot \rangle$ just denotes the Euclidean scalar product. To prove the desired asymptotic expansion, it suffices to take $\varphi \in C_0^{\infty}([0, \infty)_{z_0} \times \mathbb{R}^n)$ and to analyze the following homogeneous polynomial in x as $z_0 \rightarrow 0$:

$$\int_{\mathbb{R}^n} \sum_{\alpha} \left\langle e^{(n+\lambda)f} w_{\alpha}, \varphi(z_0, z) P(z_0, z; \cdot)^{n+\lambda} p_{\alpha}\left(x - 2(z - \cdot)\langle z - \cdot, x \rangle \cdot \frac{P(z_0, z; \cdot)}{z_0}\right) \right\rangle dz, \tag{6-29}$$

where the bracket $\langle w_{\alpha}, \cdot \rangle$ means the distributional pairing coming from pairing with respect to the canonical measure dS on S^n , which in \mathbb{R}^n becomes the measure $4^n e^{-nf} dz$, and so the e^{nf} in (6-29) cancels out if one works with the Euclidean measure dz , which we do now. We have a convolution kernel in z and thus apply the Fourier transform in z (denoted \mathcal{F}): writing $P(z_0; |z - z'|)$ for $P(z_0, z; z')$, the integral (6-29) becomes (up to nonzero multiplicative constant)

$$I(z_0, x) := \sum_{\alpha} \left\langle \mathcal{F}^{-1}(e^{\lambda f} w_{\alpha}), \mathcal{F}(\varphi) \cdot \mathcal{F}_{\zeta \rightarrow \cdot} \left(P(z_0; |\zeta|)^{n+\lambda} p_{\alpha}\left(x - 2\frac{\zeta \langle \zeta, x \rangle}{z_0} P(z_0; |\zeta|)\right) \right) \right\rangle_{\mathbb{R}^n}.$$

We can expand $p_{\alpha}(x - (2\zeta \langle \zeta, x \rangle / z_0) P(z_0; |\zeta|))$ so that

$$P(z_0; |\zeta|)^{n+\lambda} p_{\alpha}\left(x - 2\frac{\zeta \langle \zeta, x \rangle}{z_0} P(z_0; |\zeta|)\right) = \sum_{r=0}^m Q_{r,\alpha}(\zeta, x) z_0^{-r} 2^r P(z_0; |\zeta|)^{n+\lambda+r},$$

where $Q_{r,\alpha}(\zeta)$ is homogeneous of degree m in x and degree $2r$ in ζ . Now we have (for some $C \neq 0$ independent of λ, r, α)

$$\frac{2^r}{z_0^r} \mathcal{F}_{\zeta \rightarrow \xi} \left(P^{n+\lambda+r}(z_0; |\zeta|) Q_{r,\alpha}(\zeta, x) \right) = \frac{C 2^{-\lambda} z_0^{-\lambda}}{\Gamma(\lambda + n + r)} \left[Q_{r,\alpha}(i\partial_{\zeta}, x) (|\zeta|^{\lambda+n/2+r} K_{\lambda+n/2+r}(|\zeta|)) \right]_{\zeta=z_0\xi},$$

where $K_{\nu}(\cdot)$ is the modified Bessel function (see [Abramowitz and Stegun 1964, Chapter 9]) defined by

$$K_{\nu}(z) := \frac{\pi (I_{-\nu}(z) - I_{\nu}(z))}{2 \sin(\nu\pi)} \quad \text{if } I_{\nu}(z) := \sum_{\ell=0}^{\infty} \frac{1}{\ell! \Gamma(\ell + \nu + 1)} \left(\frac{z}{2}\right)^{2\ell+\nu}, \tag{6-30}$$

satisfying that $|K_{\nu}(z)| = \mathcal{O}(e^{-z}/\sqrt{z})$ as $z \rightarrow \infty$, and, for $s \notin \mathbb{N}_0$,

$$\mathcal{F}((1 + |z|^2)^{-s})(\xi) = \frac{2^{-s+1} (2\pi)^{n/2}}{\Gamma(s)} |\xi|^{s-n/2} K_{s-n/2}(|\xi|).$$

When $\lambda \notin (-\frac{1}{2}n + \mathbb{Z}) \cup (-n - \frac{1}{2}\mathbb{N}_0)$, we have

$$\begin{aligned}
 & 2^{-\lambda} z_0^{-\lambda} Q_{r,\alpha}(i\partial_\zeta, x)(|\zeta|^{\lambda+n/2+r} K_{\lambda+n/2+r}(|\zeta|))|_{\zeta=z_0\xi} \\
 &= \frac{2^{r+n/2}\pi z_0^{-\lambda}}{2 \sin(\pi(\lambda + \frac{1}{2}n + r))} \\
 & \times \left(\sum_{\ell=0}^{\infty} \frac{z_0^{2(\ell-r)} Q_{r,\alpha}(i\partial_\xi, x)(|\frac{1}{2}\xi|^{2\ell})}{\ell! \Gamma(\ell - \lambda - \frac{1}{2}n - r + 1)} - z_0^{2\lambda+n} \sum_{\ell=0}^{\infty} \frac{z_0^{2\ell} Q_{r,\alpha}(i\partial_\xi, x)(|\frac{1}{2}\xi|^{2(\lambda+r+\ell)+n})}{\ell! \Gamma(\ell + \lambda + \frac{1}{2}n + r + 1)} \right). \tag{6-31}
 \end{aligned}$$

Here the powers of $|\xi|$ are homogeneous distributions (note that, for $\lambda \notin \mathcal{R}_m$, the exceptional powers $|\xi|^{-n-j}$, $j \in \mathbb{N}_0$, do not appear) and the pairing of (6-31) with $\mathcal{F}^{-1}(e^{\lambda f} w_\alpha) \mathcal{F}(\varphi)$ makes sense since this distribution is Schwartz, as w_α has compact support. We deduce from this expansion that, for any $w_\alpha \in \mathcal{D}'(\mathbb{R}^n)$ supported near 0 and $\varphi \in C_0^\infty(\mathbb{R}^n)$, when $\lambda \notin (-\frac{1}{2}n + \mathbb{Z}) \cup (-n - \frac{1}{2}\mathbb{N}_0)$,

$$I(z_0, x) = z_0^{-\lambda} F_-(z_0, x) + z_0^{n+\lambda} F_+(z_0, x)$$

for some smooth functions $F_\pm \in C^\infty([0, \epsilon) \times \mathbb{R}^n)$ homogeneous of degree m in x . We need to analyze $F_-(0, x)$, which is obtained by computing the term of order 0 in ξ in the expansion (6-31) (that is, the terms with $\ell = r$ in the first sum; note that the terms with $\ell < r$ in this sum are zero): we obtain, for some universal constant $C \neq 0$,

$$F_-(0, x) = C \sum_\alpha \langle e^{\lambda f} w_\alpha, \varphi \rangle_{\mathbb{R}^n} \sum_{r=0}^m \frac{(-1)^r 2^{-r} \Gamma(\lambda + \frac{1}{2}n)}{r! \Gamma(\lambda + n + r)} Q_{r,\alpha}(i\partial_\xi, x)(|\xi|^{2r}),$$

where we have used the inversion formula $\Gamma(1-z)\Gamma(z) = \pi / \sin(\pi z)$ and $Q_{r,\alpha}(i\partial_\xi, x)(|\xi|^{2r})$ is constant in ξ . Using the Fourier transform, we notice that

$$Q_{r,\alpha}(i\partial_\xi, x)(|\xi|^{2r}) = \Delta_\zeta^r Q_{r,\alpha}(\zeta, x)|_{\zeta=0} = \Delta_\zeta^r (p_\alpha(x - \zeta \langle \zeta, x \rangle))|_{\zeta=0}.$$

We use Lemma A.5 to deduce that

$$F_-(0, x) = C \sum_\alpha \langle e^{\lambda f} w_\alpha, \varphi \rangle_{\mathbb{R}^n} p_\alpha(x) m! \frac{\Gamma(\lambda + \frac{1}{2}n)}{\Gamma(\lambda + n + m)} \sum_{r=0}^m \frac{(-1)^r \Gamma(\lambda + n + m)}{(m-r)! \Gamma(\lambda + n + r)}.$$

The sum over r is a nonzero polynomial of order m in λ , and, using the binomial formula, we see that its roots are $\lambda = -n - m + 2, \dots, -n + 1$; therefore, we deduce that

$$F_-(0, x) = C \langle e^{\lambda f} w, \varphi \rangle_{\mathbb{R}^n} \frac{\Gamma(\lambda + \frac{1}{2}n)}{(\lambda + n + m - 1) \Gamma(\lambda + n - 1)}.$$

We obtain the claimed result except for $\lambda \in -\frac{1}{2}n + \mathbb{N}$ by using that the volume measure on \mathbb{S}^n is $4^{-n} e^{nf}$.

Now assume that $\lambda = -\frac{1}{2}n + j$ with $j \in \mathbb{N}$. The Bessel function satisfies, for $j \in \mathbb{N}$,

$$|\xi|^j K_j(|\xi|) = - \sum_{\ell=0}^{j-1} \frac{(-1)^\ell 2^{j-1-2\ell} (j - \ell - 1)!}{\ell!} |\xi|^{2\ell} + |\xi|^{2j} (\log(|\xi|) L_j(|\xi|) + H_j(|\xi|))$$

for some function $L_j, H_j \in C^\infty(\mathbb{R}^+) \cap L^2(\mathbb{R}^+)$ with $L_j(0) \neq 0$. Then we apply the same arguments as before, and this implies the desired statement. \square

We obtain as a corollary:

Corollary 6.9. *For $m \in \mathbb{N}_0$ and $\lambda \notin \mathcal{R}_m$, the operator*

$$\mathcal{P}_\lambda^- : \mathcal{D}'(\mathbb{S}^n; \otimes_S^m(T^*\mathbb{S}^n) \cap \ker \mathcal{T}) \rightarrow C^\infty(\mathbb{H}^{n+1}; \otimes_S^m(T^*\mathbb{H}^{n+1}))$$

is injective.

This corollary immediately implies the injectivity part of [Theorem 6](#) in [Section 5B](#).

7. Expansions of eigenstates of the Laplacian

In this section, we show the surjectivity of the Poisson operator \mathcal{P}_λ^- (see [Theorem 6](#) in [Section 5B](#)). For that, we take an eigenstate u of the Laplacian on M and lift it to \mathbb{H}^{n+1} . The resulting tensor is tempered and thus expected to have a weak asymptotic expansion at the conformal boundary \mathbb{S}^n ; a precise form of this expansion is obtained by a careful analysis of both the Laplacian and the divergence-free condition. We then show that $u = \mathcal{P}_\lambda^- w$, where w is some constant times the coefficient of $\rho^{-\lambda}$ in the expansion of u (compare with [Lemma 6.8](#)).

7A. Indicial calculus and general weak expansion. Recall the bundle $E^{(m)}$ defined in [\(6-5\)](#). The operator Δ acting on $C^\infty(\bar{\mathbb{B}}^{n+1}; E^{(m)})$ is an elliptic differential operator of order 2 that lies in the 0-calculus of [\[Mazzeo and Melrose 1987\]](#), which essentially means that it is an elliptic polynomial in elements of the Lie algebra $\mathcal{V}_0(\bar{\mathbb{B}}^{n+1})$ of smooth vector fields vanishing at the boundary of the closed unit ball $\bar{\mathbb{B}}^{n+1}$. Let $\rho \in C^\infty(\bar{\mathbb{B}}^{n+1})$ be a smooth geodesic boundary defining function (see the paragraph preceding [\(6-25\)](#)). The theory developed by Mazzeo [\[1991\]](#) shows that solutions of $\Delta u = su$ which are in $\rho^{-N}L^2(\mathbb{B}^{n+1}; E^{(m)})$ for some N have weak asymptotic expansions at the boundary $\mathbb{S}^n = \partial\bar{\mathbb{B}}^{n+1}$, where ρ is any geodesic boundary defining function. To make this more precise, we introduce the *indicial family* of Δ : if $\lambda \in \mathbb{C}$, $\nu \in \mathbb{S}^n$, then there exists a family $I_{\lambda,\nu}(\Delta) \in \text{End}(E^{(m)}(\nu))$ depending smoothly on $\nu \in \mathbb{S}^n$ and holomorphically on λ such that, for all $u \in C^\infty(\bar{\mathbb{B}}^{n+1}; E^{(m)})$,

$$t^{-\lambda} \Delta(\rho^\lambda u)(\theta(t, \nu)) = I_{\lambda,\nu}(\Delta)u(\theta(0, \nu)) + \mathcal{O}(t)$$

near \mathbb{S}^n , where the remainder is estimated with respect to the metric g_H . Notice that $I_{\lambda,\nu}(\Delta)$ is independent of the choice of boundary defining function ρ .

For $\sigma \in \mathbb{C}$, the *indicial set* $\text{spec}_b(\Delta - \sigma; \nu)$ at $\nu \in \mathbb{S}^n$ of $\Delta - \sigma$ is the set

$$\text{spec}_b(\Delta - \sigma; \nu) := \{\lambda \in \mathbb{C} \mid I_{\lambda,\nu}(\Delta) - \sigma \text{ Id is not invertible}\}.$$

Then [\[Mazzeo 1991, Theorem 7.3\]](#) gives the following:²

²The full power of [\[Mazzeo 1991\]](#) is not needed for this lemma. In fact, it can be proved in a direct way by viewing the equation $(\Delta - \sigma)u = 0$ as an ordinary differential equation in the variable $\log \rho$. The indicial operator gives the constant coefficient principal part and the remaining terms are exponentially decaying; an iterative argument shows the needed asymptotics.

Lemma 7.1. Fix σ and assume that $\text{spec}_b(\Delta - \sigma; \nu)$ is independent of $\nu \in \mathbb{S}^n$. If $u \in \rho^\delta L^2(\bar{\mathbb{B}}^{n+1}; E^{(m)})$ with respect to the Euclidean measure for some $\delta \in \mathbb{R}$, and $(\Delta - \sigma)u = 0$, then u has a weak asymptotic expansion at $\mathbb{S}^n = \{\rho = 0\}$ of the form

$$u = \sum_{\substack{\lambda \in \text{spec}_b(\Delta - \sigma) \\ \text{Re}(\lambda) > \delta - \frac{1}{2}}} \sum_{\substack{\ell \in \mathbb{N}_0 \\ \text{Re}(\lambda) + \ell < \delta - \frac{1}{2} + N}} \sum_{p=0}^{k_{\lambda,\ell}} \rho^{\lambda+\ell} (\log \rho)^p w_{\lambda,\ell,p} + \mathcal{O}(\rho^{\delta+N-\frac{1}{2}-\epsilon})$$

for all $N \in \mathbb{N}$ and all $\epsilon > 0$ small, where $k_{\lambda,\ell} \in \mathbb{N}_0$, and $w_{\lambda,\ell,p}$ are in the Sobolev spaces

$$w_{\lambda,\ell,p} \in H^{-\text{Re}(\lambda)-\ell+\delta-\frac{1}{2}}(\mathbb{S}^n; E^{(m)}).$$

Here the weak asymptotic means that, for any $\varphi \in C^\infty(\mathbb{S}^n)$, as $t \rightarrow 0$,

$$\int_{\mathbb{S}^n} u(\theta(t, \nu))\varphi(\nu) dS_\rho(\nu) = \sum_{\substack{\lambda \in \text{spec}_b(\Delta - \sigma) \\ \text{Re}(\lambda) > \delta - \frac{1}{2}}} \sum_{\substack{\ell \in \mathbb{N}_0 \\ \text{Re}(\lambda) + \ell < \delta - \frac{1}{2} + N}} \sum_{p=0}^{k_{\lambda,\ell}} t^{\lambda+\ell} \log(t)^p \langle w_{\lambda,\ell,p}, \varphi \rangle + \mathcal{O}(t^{\delta+N-\frac{1}{2}-\epsilon}), \tag{7-1}$$

where dS_ρ is the measure on \mathbb{S}^n induced by the metric $(\rho^2 g_H)|_{\mathbb{S}^n}$ and the distributional pairing is with respect to this measure. Moreover, the remainder $\mathcal{O}(t^{\delta+N-1/2-\epsilon})$ is conormal in the sense that it remains $\mathcal{O}(t^{\delta+N-1/2-\epsilon})$ after applying the operator $t\partial_t$ any finite number of times, and it depends on some Sobolev norm of φ .

Remark. The existence of the expansion (7-1) proved by Mazzeo [1991, Theorem 7.3] is independent of the choice of ρ , but the coefficients in the expansion depend on the choice of ρ . Let $\lambda_0 \in \text{spec}_b(\Delta - \sigma)$ with $\text{Re}(\lambda_0) > \delta - \frac{1}{2}$ be an element in the indicial set and assume that $k_{\lambda_0,0} = 0$, which means that the exponent ρ^{λ_0} in the weak expansion (7-1) has no log term. Assume also that there is no element $\lambda \in \text{spec}_b(\Delta - \sigma)$ with $\text{Re}(\lambda_0) > \text{Re}(\lambda) > \delta - \frac{1}{2}$ such that $\lambda \in \lambda_0 - \mathbb{N}$. Then it is direct to see from the weak expansion that, for a fixed function $\chi \in C^\infty(\mathbb{B}^{n+1})$ equal to 1 near \mathbb{S}^n and supported close to \mathbb{S}^n and for each $\varphi \in C^\infty(\mathbb{B}^{n+1})$, the Mellin transform

$$h(\zeta) := \int_{\mathbb{B}^{n+1}} \rho(y)^\zeta \chi(y)\varphi(y)u(y) d\text{Vol}_{g_H}(y), \quad \text{Re } \zeta > n + \frac{1}{2} - \delta,$$

(with values in E^m) has a meromorphic extension to $\zeta \in \mathbb{C}$ with a simple pole at $\zeta = n - \lambda_0$ and residue

$$\text{Res}_{\zeta=n-\lambda_0} h(\zeta) = \langle w_{\lambda_0,0,0}, \varphi|_{\mathbb{S}^n} \rangle. \tag{7-2}$$

As an application, if ρ' is another geodesic boundary defining function, one has $\rho = e^f \rho' + \mathcal{O}(\rho')$ for some $f \in C^\infty(\mathbb{S}^n)$ and we deduce that, if $w'_{\lambda_0,0,0}$ is the coefficient of $(\rho')^{\lambda_0}$ in the weak expansion of u using ρ' , then, as a distribution on \mathbb{S}^n ,

$$w'_{\lambda_0,0,0} = e^{\lambda_0 f} w_{\lambda_0,0,0}. \tag{7-3}$$

In particular, under the assumption above for λ_0 (this assumption can similarly be seen to be independent of the choice of ρ), if one knows the exponents of the asymptotic expansion, then proving that the

coefficient of ρ^{λ_0} term is nonzero can be done locally near any point of \mathbb{S}^n and with any choice of geodesic boundary defining function.

Finally, if $w_{\lambda_0,0,0}$ is the coefficient of $\rho_0^{\lambda_0}$ in the weak expansion with boundary defining function ρ_0 defined in (3-34) and if $\gamma^*u = u$ for some hyperbolic isometry $\gamma \in G$, we can use that $\rho_0 \circ \gamma = N_\gamma^{-1} \cdot \rho_0 + \mathcal{O}(\rho_0^2)$ near \mathbb{S}^n , together with (7-2) to get

$$L_\gamma^* w_{\lambda_0,0,0} = N_\gamma^{\lambda_0} w_{\lambda_0,0,0} \in \mathcal{D}'(\mathbb{S}^n; E^{(m)}) \tag{7-4}$$

as distributions on \mathbb{S}^n (with respect to the canonical measure on \mathbb{S}^n) with values in $E^{(m)}$. Here N_γ, L_γ are as defined in Section 3E. If we view $w_{\lambda_0,0,0}$ as a distribution with values in $\otimes_S^m T^*\mathbb{S}^n$, the covariance becomes

$$L_\gamma^* w_{\lambda_0,0,0} = N_\gamma^{\lambda_0-m} w_{\lambda_0,0,0} \in \mathcal{D}'(\mathbb{S}^n; \otimes_S^m T^*\mathbb{S}^n). \tag{7-5}$$

Using the calculations of Section 6B, we will compute the indicial family of the Laplacian on $E^{(m)}$:

Lemma 7.2. *Let Δ be the Laplacian on sections of $E^{(m)}$. Then the indicial set $\text{spec}_b(\Delta - \sigma, \nu)$ does not depend on $\nu \in \mathbb{S}^n$ and is equal to³*

$$\bigcup_{k=0}^{\lfloor m/2 \rfloor} \{ \lambda \mid -\lambda^2 + n\lambda + m + 2k(2m + n - 2k - 2) = \sigma \} \cup \bigcup_{k=0}^{\lfloor (m-1)/2 \rfloor} \{ \lambda \mid -\lambda^2 + n\lambda + n + 3(m - 1) + 2k(n + 2m - 2k - 4) = \sigma \}.$$

Proof. We consider an isometry mapping the ball model \mathbb{B}^{n+1} to the half-plane model \mathbb{U}^{n+1} which also maps ν to 0 and do all the calculations in \mathbb{U}^{n+1} with the geodesic boundary defining function z_0 near 0. By (6-7), each tensor $u \in E^{(m)}$ is determined uniquely by its $E_0^{(m)}$ and $E_1^{(m)}$ components, which are denoted u_0 and u_1 ; therefore, it suffices to understand how the corresponding components of $I_{\lambda,\nu}(\Delta)u$ are determined by u_0 and u_1 . We can use the geodesic boundary defining function $\rho = z_0$; note that $\Delta z_0^\lambda = \lambda(n - \lambda)z_0^\lambda$ for all $\lambda \in \mathbb{C}$.

Assume first that u satisfies $u_1 = 0$ and u_0 is constant in the frame $\mathcal{S}(Z_I^*)$. Then, by Lemma 6.4,

$$\begin{aligned} \pi_0(z_0^{-\lambda} \Delta(z_0^\lambda u)) &= R_0 u_0 = (\lambda(n - \lambda) + m)u_0 + m(m - 1)\mathcal{S}(z_0^{-2}h \otimes \mathcal{T}(u_0)), \\ \pi_1(z_0^{-\lambda} \Delta(z_0^\lambda u)) &= 0. \end{aligned}$$

Assume now that u satisfies $u_0 = 0$ and u_1 is constant in the frame $\mathcal{S}(Z_0^* \otimes Z_J^*)$. Then, by Lemma 6.4,

$$\begin{aligned} \pi_0(z_0^{-\lambda} \Delta(z_0^\lambda u)) &= 0, \\ \pi_1(z_0^{-\lambda} \Delta(z_0^\lambda u)) &= R_1 u_1 = (\lambda(n - \lambda) + n + 3(m - 1))u_1 + (m - 1)(m - 2)\mathcal{S}(Z_0^* \otimes z_0^{-2}h \otimes \mathcal{T}(u_1)). \end{aligned}$$

We see that the indicial operator does not intertwine the u_0 and u_1 components and it remains to understand for which λ the number s is a root of R_0 or R_1 .

³Our argument in the next section does not actually use the precise indicial roots, as long as they are independent of ν and form a discrete set.

Next, we consider the decomposition (4-5), where we define $\mathcal{I}(u) = \frac{1}{2}(m+2)(m+1)\mathcal{S}(z_0^{-2}h \otimes u)$ for $u \in E_0^{(m)}$; we have

$$u_0 = \sum_{k=0}^{\lfloor m/2 \rfloor} \mathcal{I}^k(\otimes u_0^k), \quad u_1 = \sum_{k=0}^{\lfloor (m-1)/2 \rfloor} \mathcal{S}(Z_0^* \otimes \mathcal{I}^k(u_1^k)),$$

where $u_0^k \in E_0^{(m-2k)}$ and $u_1^k \in E_0^{(m-2k-1)}$ are trace-free tensors. Using (4-4), we calculate

$$\begin{aligned} R_0(\mathcal{I}^k(u_0^k)) &= (\lambda(n-\lambda) + m)\mathcal{I}^k(u_0^k) + 2\mathcal{I}(\mathcal{T}(\mathcal{I}^k(u_0^k))) \\ &= (-\lambda^2 + n\lambda + m + 2k(2m+n-2k-2))\mathcal{I}^k(u_0^k), \\ R_1(\mathcal{S}(Z_0^* \otimes \mathcal{I}^k(u_1^k))) &= (\lambda(n-\lambda) + n + 3(m-1))\mathcal{S}(Z_0^* \otimes \mathcal{I}^k(u_1^k)) + 2\mathcal{S}(Z_0^* \otimes \mathcal{I}(\mathcal{T}(\mathcal{I}^k(u_1^k)))) \\ &= (-\lambda^2 + n\lambda + n + 3(m-1) + 2k(n+2m-2k-4))\mathcal{S}(Z_0^* \otimes \mathcal{I}^k(u_1^k)), \end{aligned}$$

which finishes the proof of the lemma. □

7B. Weak expansions in the divergence-free case. By Lemma 7.1, we now know that solutions of $\Delta u = \sigma u$ that are trace-free symmetric tensors of order m in some weighted L^2 space have weak asymptotic expansions at the boundary of $\bar{\mathbb{B}}^{n+1}$ with exponents obtained from the indicial set of Lemma 7.2. In fact, we can be more precise about the exponents which really appear in the weak asymptotic expansion if we ask that u also be divergence-free:

Lemma 7.3. *Let $u \in \rho^\delta L^2(\bar{\mathbb{B}}^{n+1}; E^{(m)})$ be a trace-free symmetric m -cotensor with ρ a geodesic boundary defining function and $\delta \in (-\infty, \frac{1}{2})$, where the measure is the Euclidean Lebesgue measure on the ball. Assume that u is a nonzero divergence-free eigentensor for the Laplacian on hyperbolic space:*

$$\Delta u = \sigma u, \quad \nabla^* u = 0 \tag{7-6}$$

for some $\sigma = m + \frac{1}{4}n^2 - \mu^2$ with $\text{Re}(\mu) \in [0, \frac{1}{2}(n+1) - \delta)$ and $\mu \neq 0$. Then the following weak expansion holds: for all $r \in [0, m]$, $N > 0$, and $\epsilon > 0$ small,

$$\begin{aligned} &(\iota_{\rho\partial_\rho})^r u \\ &= \sum_{\substack{\ell \in \mathbb{N}_0 \\ \text{Re}(-\mu) + \ell < N - \epsilon}} \rho^{n/2 - \mu + r + \ell} w_{-\mu, \ell}^r + \sum_{\substack{\ell \in \mathbb{N}_0 \\ \text{Re}(\mu) + \ell < N - \epsilon}} \sum_{p=0}^{k_{\mu, \ell}} \rho^{n/2 + \mu + r + \ell} \log(\rho)^p w_{\mu, \ell, p}^r + \mathcal{O}(\rho^{n/2 + N + r - \epsilon}) \end{aligned} \tag{7-7}$$

with $w_{-\mu, \ell}^r \in H^{-n/2 + \text{Re}(\mu) - r - \ell + \delta - 1/2}(\mathbb{S}^n; E^{(m-r)})$, $w_{\mu, \ell, p}^r \in H^{-n/2 - \text{Re}(\mu) - r - \ell + \delta - 1/2}(\mathbb{S}^n; E^{(m-r)})$. Moreover, if $\mu \notin \frac{1}{2}\mathbb{N}_0$, then $k_{\mu, \ell} = 0$.

Remark. (i) If u is the lift to \mathbb{H}^{n+1} of an eigentensor on a compact quotient $M = \Gamma \backslash \mathbb{H}^{n+1}$, then $u \in L^\infty(\mathbb{B}^{n+1}; E^{(m)})$ and so, for all $\epsilon > 0$, the following regularity holds:

$$w_{-\mu, 0} \in H^{-n/2 + \text{Re}(\mu) - \epsilon}(\mathbb{S}^n; E^{(m)}), \quad w_{\mu, 0, 0} \in H^{-n/2 - \text{Re}(\mu) - \epsilon}(\mathbb{S}^n; E^{(m)}).$$

(ii) The existence of the expansion (7-7) does not depend on the choice of ρ . For $r = 0$, this follows from analyzing the Mellin transform of u as in the remark following Lemma 7.1. For $r > 0$, we additionally use that, if ρ' is another geodesic boundary defining function, then $\rho\partial_\rho - \rho'\partial_{\rho'} \in \rho \cdot {}^0T\bar{\mathbb{B}}^{n+1}$

(indeed, the dual covector by the metric is $\rho^{-1}d\rho - (\rho')^{-1}d\rho'$ and we have $\rho' = e^f \rho$ for some smooth function f on $\overline{\mathbb{B}^{n+1}}$). Therefore, $(\iota_{\rho'\partial_{\rho'}})^r u$ is a linear combination of contractions with 0-vector fields of $\rho^{r-r'}(\iota_{\rho\partial_{\rho}})^{r'} u$ for $0 \leq r' \leq r$, which have the desired asymptotic expansion. Moreover, as follows from (7-3), for each $r \in [0, m]$, the condition that $w_{-\mu,0}^{r'} = 0$ for all $r' \in [0, r]$ also does not depend on the choice of ρ , and the same can be said about $w_{\mu,0,0}^{r'}$ when $\mu \notin \frac{1}{2}\mathbb{N}_0$.

Proof. It suffices to describe the weak asymptotic expansion of u near any point $\nu \in \mathbb{S}^n$. For that, we work in the half-space model \mathbb{U}^{n+1} by sending $-\nu$ to ∞ and ν to 0 as we did before (composing a rotation of the ball model with the map (3-5)). Since the choice of geodesic boundary defining function does not change the nature of the weak asymptotic expansion (but only the coefficients), we can take the geodesic boundary defining function ρ to be equal to $\rho(z_0, z) = z_0$ inside $|z| + z_0 < 1$ (which corresponds to a neighborhood of ν in the ball model). Considering the weak asymptotic (7-1) of u near 0 amounts to taking φ supported near ν in \mathbb{S}^n in (7-1); for instance, if we work in the half-space model, we shall consider $\varphi(z)$ supported in $|z| < 1$ in the boundary of \mathbb{U}^{n+1} .

We have the decomposition $u = \sum_{k=0}^m u_k$ with $u_k \in \rho^\delta L^2(\mathbb{U}^{n+1}; E_k^{(m)})$ and we write $u_k = S((Z_0^*)^{\otimes k} \otimes u'_k)$ for some $u'_k \in \rho^\delta L^2(\mathbb{U}^{n+1}; E_0^{(m-k)})$ following what we did in (6-6). Now, since $u \in \rho^\delta L^2(\overline{\mathbb{B}^{n+1}}) = \rho_0^\delta L^2(\overline{\mathbb{B}^{n+1}})$ satisfies $\Delta u = \sigma u$, we deduce from the form of the Laplacian near $\rho = 0$ that u is in $\rho_0^{\delta-2k} H^{2k}(\overline{\mathbb{B}^{n+1}}; E^{(m)})$ for all $k \in \mathbb{N}$, where H^k denotes the Sobolev space of order k associated to the Euclidean Laplacian on the closed unit ball. Then, by Sobolev embedding, one has that, for each $t > 0$, $u|_{z_0=t}$ belongs to $(1 + |z|)^N L^2(\mathbb{R}_z^n; E^{(m)})$ for some $N \in \mathbb{N}$ and we can consider its Fourier transform in z , as a tempered distribution.⁴ Then Fourier transforming the equation $(\pi_0 + \pi_1)(\Delta u - \sigma u) = 0$ in the z variable (recall that π_i is the orthogonal projection on $E_i^{(m)}$), and writing the Fourier variable ξ as $\xi = \sum_{i=1}^n \xi_i dz_i = \sum_{i=1}^n z_0 \xi_i Z_i^*$, with the notations of Lemma 6.4, we get

$$\sum_{I \in \mathcal{A}^m} ((-Z_0)^2 + nZ_0 + z_0^2 |\xi|^2 + m - \sigma) \hat{f}_I S(Z_I^*) + 2i \sum_{J \in \mathcal{A}^{m-1}} \hat{g}_J S(\xi \otimes Z_J^*) + m(m-1) \sum_I \hat{f}_I S(z_0^{-2} h \otimes \mathcal{T}(S(Z_I^*))) = 0. \tag{7-8}$$

and

$$\sum_{J \in \mathcal{A}^{m-1}} ((-Z_0)^2 + nZ_0 + z_0^2 |\xi|^2 + n + 3(m-1) - \sigma) \hat{g}_J S(Z_J^*) - 2im \sum_{I \in \mathcal{A}^m} \hat{f}_I \iota_\xi S(Z_I^*) - 2im(m-1) \sum_{I \in \mathcal{A}^m} \hat{f}_I S(\xi \otimes \mathcal{T}(S(Z_I^*))) + (m-1)(m-2) \sum_{J \in \mathcal{A}^{m-1}} \hat{g}_J S(z_0^{-2} h \otimes \mathcal{T}(S(Z_J^*))) = 0, \tag{7-9}$$

where hat denotes Fourier transform in z and ι_ξ means $\sum_{j=1}^n z_0 \xi_j \iota_{Z_j}$.

⁴Unlike in Lemma 6.8, we only use Fourier analysis here for convenience of notation — all the calculations below could be done with differential operators in z instead.

Similarly, we Fourier transform in z the equation $(\pi_0 + \pi_1)(\nabla^*u) = 0$ using [Lemma 6.5](#) to obtain

$$\begin{aligned} \sum_{I \in \mathcal{A}^m} i \hat{f}_I \iota_\xi \mathcal{S}(Z_I^*) &= \frac{1}{m} \sum_{J \in \mathcal{A}^{m-1}} ((n+m-1)\hat{g}_J - Z_0(\hat{g}_J))\mathcal{S}(Z_J^*), \\ \sum_{I \in \mathcal{A}^m} (Z_0 \hat{f}_I - (n+m-1)\hat{f}_I)\mathcal{T}(\mathcal{S}(Z_I^*)) &= \frac{1}{m} \sum_{J \in \mathcal{A}^{m-1}} i \hat{g}_J \iota_\xi \mathcal{S}(Z_J^*). \end{aligned} \tag{7-10}$$

Now, we use the correspondence between symmetric tensors and homogeneous polynomials to facilitate computations, as explained in [Section 4A](#) and in the proof of [Lemma 6.8](#); that is, to $\mathcal{S}(Z_I^*)$, we associate the polynomial x_I on \mathbb{R}^n . If $\xi \in \mathbb{R}^n$ is a fixed element and $u \in \text{Pol}^m(\mathbb{R}^n)$, we write $\partial_\xi u = du \cdot \xi \in \text{Pol}^{m-1}(\mathbb{R}^n)$ for the derivative of u in the direction of ξ and ξ^*u for the element $\langle \xi, \cdot \rangle_{\mathbb{R}^n} u \in \text{Pol}^{m+1}(\mathbb{R}^n)$. The trace map \mathcal{T} becomes $-(1/(m(m-1)))\Delta_x$. We define $\hat{u}_0 := \sum_{I \in \mathcal{A}^m} \hat{f}_I x_I$ and $\hat{u}_1 = \sum_{J \in \mathcal{A}^{m-1}} \hat{g}_J x_J$. The elements $\hat{f}_I(z_0, \xi)$ and $\hat{g}_I(z_0, \xi)$ belong to the space $\mathcal{C}^\infty(\mathbb{R}_{z_0}^+; \mathcal{S}'(\mathbb{R}_\xi^n))$. We decompose them as

$$\hat{u}_0 = \sum_{j=0}^{\lfloor m/2 \rfloor} |x|^{2j} \hat{u}_0^{2j} \quad \text{and} \quad \hat{u}_1 = \sum_{j=0}^{\lfloor (m-1)/2 \rfloor} |x|^{2j} \hat{u}_1^{2j} \tag{7-11}$$

for some $\hat{u}_i^{2j} \in \text{Pol}_0^{m-i-2j}(\mathbb{R}^n)$ (harmonic in x , that is, trace-free).

Using the homogeneous polynomial description of u_0 , [\(7-8\)](#) becomes

$$-(Z_0)^2 + nZ_0 + z_0^2 |\xi|^2 + m - \sigma) \hat{u}_0 + 2iz_0 \xi^* \hat{u}_1 - |x|^2 \Delta_x \hat{u}_0 = 0. \tag{7-12}$$

First, if W is a harmonic homogeneous polynomial in x of degree j , one has $\Delta_x(\xi^*W) = -2\partial_\xi W$ and $\Delta_x^2(\xi^*W) = 0$; thus one can write

$$\xi^*W = \left(\xi^*W - \frac{\partial_\xi W}{n+2(j-1)} |x|^2 \right) + \frac{\partial_\xi W}{n+2(j-1)} |x|^2 \tag{7-13}$$

for the decomposition [\(4-5\)](#) of ξ^*W . In particular, one can write the decomposition [\(4-5\)](#) of $\xi^*\hat{u}_1$ as

$$\xi^*\hat{u}_1 = \sum_{j=0}^{\lfloor (m-1)/2 \rfloor} |x|^{2j} \left(\xi^*\hat{u}_1^{2j} - \frac{\partial_\xi \hat{u}_1^{2j}}{n+2(m-2-2j)} |x|^2 + \frac{\partial_\xi \hat{u}_1^{2(j-1)}}{n+2(m-2j)} \right).$$

We can write $\Delta_x \hat{u}_0 = \sum_{j=0}^{\lfloor m/2 \rfloor} \lambda_j |x|^{2j-2} \hat{u}_0^{2j}$ for $\lambda_j = -2j(n+2(m-j-1))$. Thus [\(7-12\)](#) gives, for $j \leq \lfloor m/2 \rfloor$,

$$(-(Z_0)^2 + nZ_0 + z_0^2 |\xi|^2 + m - \sigma - \lambda_j) \hat{u}_0^{2j} + 2iz_0 \left(\xi^*\hat{u}_1^{2j} - \frac{|x|^2 \partial_\xi \hat{u}_1^{2j}}{n+2(m-2-2j)} + \frac{\partial_\xi \hat{u}_1^{2(j-1)}}{n+2(m-2j)} \right) = 0. \tag{7-14}$$

Notice that $\iota_\xi(\mathcal{S}(Z_I^*))$ corresponds to the polynomial $(z_0/m) dx_I \cdot \xi = (z_0/m) \partial_{\xi^*} x_I$ if $I \in \mathcal{A}^m$. From [\(7-10\)](#) we thus have, for $c_m := n+m-1$,

$$\begin{aligned} -iz_0 \partial_\xi \hat{u}_0 &= (Z_0 - c_m) \hat{u}_1, \\ -iz_0 \partial_\xi \hat{u}_1 &= (Z_0 - c_m) \Delta_x \hat{u}_0. \end{aligned} \tag{7-15}$$

Next, (7-9) implies

$$-(Z_0)^2 + nZ_0 + z_0^2|\xi|^2 + n + 3(m - 1) - \sigma \hat{u}_1 - 2iz_0\partial_\xi \hat{u}_0 + 2iz_0\xi^* \Delta_x \hat{u}_0 - |x|^2 \Delta_x \hat{u}_1 = 0.$$

Using (7-15), this can be rewritten as

$$-(Z_0)^2 + (n + 2)Z_0 + z_0^2|\xi|^2 - n + m - 1 - \sigma \hat{u}_1 + 2iz_0\xi^* \Delta_x \hat{u}_0 - |x|^2 \Delta_x \hat{u}_1 = 0. \tag{7-16}$$

We can write $\Delta_x \hat{u}_1 = \sum_{j=0}^{(m-1)/2} \lambda'_j |x|^{2j-2} \hat{u}_1^{2j}$ for $\lambda'_j = -2j(n + 2(m - j - 2))$. From (7-16), we get

$$\begin{aligned} & (-(Z_0)^2 + (n + 2)Z_0 + z_0^2|\xi|^2 - n + m - 1 - \sigma - \lambda'_j) \hat{u}_1^{2j} \\ & + 2iz_0 \left(\lambda_{j+1} \xi^* \hat{u}_0^{2(j+1)} - \frac{\lambda_{j+1} \partial_\xi \hat{u}_0^{2(j+1)}}{n + 2(m - 3 - 2j)} |x|^2 + \frac{\lambda_j \partial_\xi \hat{u}_0^{2j}}{n + 2(m - 1 - 2j)} \right) = 0. \end{aligned} \tag{7-17}$$

We shall now partially uncouple the system of equations for \hat{u}_0^{2j} and \hat{u}_1^{2j} . Using (7-13) and applying the decomposition (4-5), we have

$$\begin{aligned} \partial_\xi (|x|^{2j} \hat{u}_0^{2j}) &= |x|^{2j} \partial_\xi \hat{u}_0^{2j} \frac{n + 2(m - j - 1)}{n + 2(m - 2j - 1)} + 2j |x|^{2j-2} \left(\xi^* \hat{u}_0^{2j} - \frac{|x|^2 \partial_\xi \hat{u}_0^{2j}}{n + 2(m - 2j - 1)} \right), \\ \partial_\xi (|x|^{2j} \hat{u}_1^{2j}) &= |x|^{2j} \partial_\xi \hat{u}_1^{2j} \frac{n + 2(m - j - 2)}{n + 2(m - 2j - 2)} + 2j |x|^{2j-2} \left(\xi^* \hat{u}_1^{2j} - \frac{|x|^2 \partial_\xi \hat{u}_1^{2j}}{n + 2(m - 2j - 2)} \right), \end{aligned}$$

and, from (7-15), this implies that, for $j \geq 0$,

$$(Z_0 - c_m) \hat{u}_1^{2j} = -iz_0 \left(\partial_\xi \hat{u}_0^{2j} \frac{n + 2(m - j - 1)}{n + 2(m - 2j - 1)} + 2(j + 1) \left(\xi^* \hat{u}_0^{2(j+1)} - \frac{|x|^2 \partial_\xi \hat{u}_0^{2(j+1)}}{n + 2(m - 2j - 3)} \right) \right), \tag{7-18}$$

and, for $j > 0$,

$$(Z_0 - c_m) \hat{u}_0^{2j} = iz_0 \left(\frac{\partial_\xi \hat{u}_1^{2(j-1)}}{2j(n + 2(m - 2j))} + \frac{1}{n + 2(m - j - 1)} \left(\xi^* \hat{u}_1^{2j} - \frac{|x|^2 \partial_\xi \hat{u}_1^{2j}}{n + 2(m - 2j - 2)} \right) \right). \tag{7-19}$$

Combining with (7-14) and (7-17) we get, for $j \geq 0$,

$$\begin{aligned} & (-(Z_0)^2 + (n + 4j)Z_0 + z_0^2|\xi|^2 + m - \sigma - \lambda_j - 4jc_m) \hat{u}_0^{2j} \\ & + 2iz_0 \frac{n + 2(m - 2j - 1)}{n + 2(m - j - 1)} \left(\xi^* \hat{u}_1^{2j} - \frac{|x|^2 \partial_\xi \hat{u}_1^{2j}}{n + 2(m - 2 - 2j)} \right) = 0, \end{aligned} \tag{7-20}$$

$$\begin{aligned} & (-(Z_0)^2 + (n + 2 + 4j)Z_0 + z_0^2|\xi|^2 - n + m - 1 - \sigma - \lambda'_j - 4jc_m) \hat{u}_1^{2j} \\ & + 2iz_0 (\lambda_{j+1} + 4j(j + 1)) \left(\xi^* \hat{u}_0^{2(j+1)} - \frac{|x|^2 \partial_\xi \hat{u}_0^{2(j+1)}}{n + 2(m - 3 - 2j)} \right) = 0, \end{aligned} \tag{7-21}$$

$$\begin{aligned} & \left(-(Z_0)^2 + \left(n + 2 - \frac{\lambda_{j+1}}{j + 1} \right) Z_0 + z_0^2|\xi|^2 - n + m - 1 - \sigma + \frac{\lambda_{j+1}}{j + 1} (c_m - j) \right) \hat{u}_1^{2j} \\ & + 2iz_0 \frac{(n + 2(m - j - 1))(n + 2(m - 2j - 2))}{n + 2(m - 2j - 1)} \partial_\xi \hat{u}_0^{2j} = 0, \end{aligned} \tag{7-22}$$

and, for $j > 0$,

$$\left(-Z_0^2 + \left(n - \frac{\lambda_j}{j}\right)Z_0 + z_0^2|\xi|^2 + m - \sigma + \frac{\lambda_j}{j}(c_m - j)\right)\hat{u}_0^{2j} - iz_0 \frac{2(m-1-2j)+n}{j(n+2(m-2j))} \partial_\xi \hat{u}_1^{2(j-1)} = 0. \tag{7-23}$$

To prove the lemma, we will show the following weak asymptotic expansion for $i = 0, 1$:

$$\begin{aligned} \langle \hat{u}_i^{2j}(z_0, \cdot), \hat{\varphi} \rangle &= \sum_{\substack{\ell \in \mathbb{N}_0, \\ \operatorname{Re}(-\mu) + \ell < N - \epsilon}} z_0^{n/2 - \mu + 2j + i + \ell} \langle \tilde{w}_{i; -\mu, \ell}^{2j}, \varphi \rangle \\ &+ \sum_{\substack{\ell \in \mathbb{N}_0, \\ \operatorname{Re}(\mu) + \ell < N - \epsilon}} \sum_{p=0}^{k_{\mu, \ell}} z_0^{n/2 + \mu + 2j + i + \ell} \log(z_0)^p \langle \tilde{w}_{i; \mu, \ell, p}^{2j}, \varphi \rangle + \mathcal{O}(z_0^{n/2 + 2j + i + N - \epsilon}), \end{aligned} \tag{7-24}$$

where $\tilde{w}_{i; -\mu, \ell}^{2j}$ and $\tilde{w}_{i; \mu, \ell, p}^{2j}$ are distributions in some Sobolev spaces in $\{|z| < 1\} \subset \mathbb{R}^n$ and, for $\mu \notin \frac{1}{2}\mathbb{N}_0$, we have $k_{\mu, \ell} = 0$.

Define, for $0 \leq r \leq m$ and $\varphi \in C_0^\infty(\mathbb{R}^n)$ supported in $\{|z| < 1\}$,

$$F^r(\varphi)(z_0) := \begin{cases} \langle \hat{u}_0^r(z_0, \cdot), \hat{\varphi} \rangle & \text{if } r \text{ is even,} \\ \langle \hat{u}_1^{r-1}(z_0, \cdot), \hat{\varphi} \rangle & \text{if } r \text{ is odd.} \end{cases}$$

Since \hat{u}_i^{r-i} is the Fourier transform in z of iterated traces of u_i , [Lemma 7.1](#) gives that the function $F^r(\varphi)(z_0)$ satisfies, for all $N \in \mathbb{N}$, $\epsilon > 0$,

$$F^r(\varphi)(z_0) = \sum_{\substack{\lambda \in \operatorname{spec}_b(\Delta - \sigma) \\ \operatorname{Re}(\lambda) > \delta - \frac{1}{2}}} \sum_{\substack{\ell \in \mathbb{N}_0, \\ \operatorname{Re}(\lambda) + \ell < N - \epsilon}} \sum_{p=0}^{k_{\lambda, \ell}^r} z_0^{\lambda + \ell} \log(z_0)^p \langle w_{\lambda, \ell, p}^r, \varphi \rangle + \mathcal{O}(z_0^{N - \epsilon}) \tag{7-25}$$

as $z_0 \rightarrow 0$ for some $w_{\lambda, \ell, p}^r$ in some Sobolev space on $\{|z| < 1\}$. We pair [\(7-20\)](#), [\(7-21\)](#) with $\hat{\varphi}$, and it is direct to see that we obtain a differential equation in z_0 of the form

$$P^r(Z_0)F^r(\varphi)(z_0) = -z_0^2 F^r(\Delta\varphi)(z_0) + z_0 F^{r+1}(Q^r\varphi)(z_0) \tag{7-26}$$

for $Z_0 = z_0 \partial_{z_0}$,

$$P^r(\lambda) := -\lambda^2 + (n + 2r)\lambda - r(n + r) - \frac{1}{4}n^2 + \mu^2 = -\left(\lambda - \frac{1}{2}n - r\right)^2 + \mu^2,$$

and Q^r some differential operator of order 1 with values in homomorphisms on the space of polynomials in x . Here we denote $F^{m+1} = 0$.

We now show the expansion [\(7-24\)](#) by induction on $r = 2j + i = m, m - 1, \dots, 0$. By plugging the expansion [\(7-25\)](#) in [\(7-26\)](#) and using

$$P^r(Z_0)z_0^\lambda \log(z_0)^p = z_0^\lambda (P_0^r(\lambda)(\log z_0)^p + p \partial_\lambda P_0^r(\lambda)(\log z_0)^{p-1} + \mathcal{O}((\log z_0)^{p-2})), \tag{7-27}$$

we see that if, for some p , $z_0^\lambda (\log z_0)^p$ is featured in the asymptotic expansion of $F^r(\varphi)(z_0)$, then either $\lambda \in \frac{1}{2}n + r - \mu + \mathbb{N}_0$, or $\lambda \in \frac{1}{2}n + r + \mu + \mathbb{N}_0$, or $z_0^{\lambda-2} (\log z_0)^p$ is featured in the expansion of $F^r(\Delta\varphi)(z_0)$. Moreover, if $p > 0$ and $\lambda \notin \{\frac{1}{2}n + r \pm \mu\}$, then either $z_0^\lambda (\log z_0)^{p'}$ is featured in $F^r(\varphi)(z_0)$ for

some $p' > p$, or $z_0^{\lambda-2}(\log z_0)^p$ is featured in $F^r(\Delta\varphi)(z_0)$, or $z_0^{\lambda-1}(\log z_0)^p$ is featured in $F^{r+1}(Q^r\varphi)(z_0)$. If $p > 0$ and $\lambda = \frac{1}{2}n + r \pm \mu$, then (since $\mu \neq 0$ and thus $\partial_\lambda P_0^r(\lambda) \neq 0$) either $z_0^\lambda(\log z_0)^{p'}$ is featured in $F^r(\varphi)(z_0)$ for some $p' > p$, or $z_0^{\lambda-2}(\log z_0)^{p-1}$ is featured in $F^r(\Delta\varphi)(z_0)$, or $z_0^{\lambda-1}(\log z_0)^{p-1}$ is featured in $F^{r+1}(Q^r\varphi)(z_0)$, however the latter two cases are only possible when $\lambda = \frac{1}{2}n + r + \mu$ and $\mu \in \frac{1}{2}\mathbb{N}_0$. Together, these facts (applied to φ as well as its images under combinations of Δ and Q^r), imply that the weak expansion of u_i^{2j} has the form (7-24).

The asymptotic expansions (7-7) now follow from (7-24), since $\rho\partial_\rho = Z_0$ for our choice of ρ and, for each $r \in [0, m]$, by (6-7) and (7-11), we see that (identifying symmetric tensors with homogeneous polynomials in (x_0, x))

$$(t_{Z_0})^r u(x_0, x) = \sum_{r'=r}^m \sum_{\substack{s \geq 0 \\ r'+2s \leq m}} c_{m,r,r',s} x_0^{r'-r} |x|^{2s} u_{r'-2[r'/2]}^{2[r'/2]+2s}(x) \tag{7-28}$$

for some constants $c_{m,r,r',s}$; for later use, we also note that $c_{m,r,r,0} \neq 0$. □

7C. Surjectivity of the Poisson operator. In this section, we prove the surjectivity part of [Theorem 6](#) in [Section 5B](#) (together with the injectivity part established in [Corollary 6.9](#), this finishes the proof of that theorem). The remaining essential component of the proof is showing that, unless $u \equiv 0$, a certain term in the asymptotic expansion of [Lemma 7.3](#) is nonzero (in particular we will see that u cannot be vanishing to infinite order on \mathbb{S}^n in the weak sense). We start with:

Lemma 7.4. *Take some u satisfying (7-6). Assume that, for all $r \in [0, m]$, the coefficient $w_{-\mu,0}^r$ of the weak expansion (7-7) is zero. (By [Remark \(ii\)](#) following [Lemma 7.3](#), this condition is independent of the choice of ρ .) Then $u \equiv 0$. If $\mu \notin \frac{1}{2}\mathbb{N}_0$, then we can replace $w_{-\mu,0}^r$ by $w_{\mu,0,0}^r$ in the assumption above.*

Proof. We choose some $v \in \mathbb{S}^n$ and transform \mathbb{B}^{n+1} to the half-space model as explained in the proof of [Lemma 7.3](#), and use the notation of that proof. Define the function $f \in C^\infty(\mathbb{B}^{n+1})$ in the half-space model as follows:

$$f = \begin{cases} z_0^{-m} u_0^{2m} & \text{if } m \text{ is even,} \\ z_0^{-m} u_1^{2m-1} & \text{if } m \text{ is odd.} \end{cases}$$

Here u_0^{2j} and u_1^{2j} are obtained by taking the inverse Fourier transforms of \hat{u}_0^{2j} and \hat{u}_1^{2j} . By (7-20) and (7-21) (see also (7-26)) we have

$$(\Delta_{\mathbb{H}^{n+1}} - \frac{1}{4}n^2 + \mu^2)f = 0. \tag{7-29}$$

Denote by $C_{\text{temp}}^\infty(\mathbb{B}^{n+1})$ the set of smooth functions f in \mathbb{B}^{n+1} which are tempered in the sense that there exists $N \in \mathbb{R}$ such that $\rho_0^N f \in L^2(\mathbb{B}^{n+1})$. Set $\lambda := -\frac{1}{2}n + \mu$; it is proved in [[van den Ban and Schlichtkrull 1987](#); [Ōshima and Sekiguchi 1980](#)] (see also [[Grellier and Otal 2005](#)] for a simpler presentation in the case $|\text{Re}(\lambda) + \frac{1}{2}n| < \frac{1}{2}n$) that the Poisson operator acting on distributions on hyperbolic space is an isomorphism

$$\mathcal{P}_\lambda^- : \mathcal{D}'(\mathbb{S}^n) \rightarrow \ker(\Delta_{\mathbb{H}^{n+1}} + \lambda(n + \lambda)) \cap C_{\text{temp}}^\infty(\mathbb{B}^{n+1})$$

for $\lambda \notin -n - \mathbb{N}_0$, and, if $\operatorname{Re}(\lambda) \geq -\frac{1}{2}n$ with $\lambda \neq 0$, any element $v \in C_{\text{temp}}^\infty(\mathbb{B}^{n+1})$ with $(\Delta_{\mathbb{H}^{n+1}} + \lambda(n + \lambda))v = 0$ and $v \neq 0$ satisfies a weak expansion for any $N \in \mathbb{N}$,

$$v = \mathcal{P}_\lambda^-(v_{-\mu, \ell}) = \sum_{\ell=0}^N \left(\rho_0^{n/2-\mu+\ell} v_{-\mu, \ell} + \sum_{p=1}^{k_{\mu, \ell}} \rho_0^{n/2+\mu+\ell} \log(\rho_0)^p v_{\mu, \ell, p} \right) + \mathcal{O}(\rho_0^{n/2-\mu+N})$$

with $v_{-\mu, 0} \neq 0$; moreover, $k_{\mu, \ell} = 0$ if $\lambda \notin -\frac{1}{2}n + \frac{1}{2}\mathbb{N}_0$, and $v_{\mu, 0, 0} \neq 0$ for such λ (here $v_{-\mu, \ell}, v_{\mu, \ell, p}$ are distributions on \mathbb{S}^n as before).⁵

Next, by (7-28), for some nonzero constant c we have

$$f = c(z_0^{-1} \iota_{z_0})^m u = c \langle u, \otimes^m \partial_{z_0} \rangle.$$

A calculation using (3-5) shows that in the ball model, using the geodesic boundary defining function ρ_0 from (3-34),

$$\partial_{z_0} = -\left(\frac{1}{2}(1 - |y|^2)v + (1 + y \cdot v)y\right) \partial_y \tag{7-30}$$

is a $C^\infty(\bar{\mathbb{B}}^{n+1})$ -linear combination of ∂_{ρ_0} and a 0-vector field. It follows from the form of the expansion (7-7) and the assumption of this lemma that the coefficient of $\rho_0^{n/2-\mu}$ of the weak expansion of f is zero. (If $\mu \notin \frac{1}{2}\mathbb{N}_0$, then we can also consider instead the coefficient of $\rho_0^{n/2+\mu}$.)

By (7-29) and the surjectivity of the scalar Poisson kernel discussed above, we now see that $f \equiv 0$. Now, for each fixed $y \in \mathbb{B}^{n+1}$ and each $\eta \in T_y \mathbb{B}^{n+1}$, we can choose v such that η is a multiple of (7-30) at y ; in fact, it suffices to take v such that the geodesic $\varphi_t(y, \eta)$ converges to $-v$ as $t \rightarrow +\infty$. Therefore, for each y, η , we have $\langle u, \otimes^m \eta \rangle = 0$ at y . Since u is a symmetric tensor, this implies $u \equiv 0$. \square

We now relax the assumptions of Lemma 7.4 to only include the term with $r = 0$:

Lemma 7.5. *Take some u satisfying (7-6). If $n = 1$ and $m > 0$, then we additionally assume that $\mu \neq \frac{1}{2}$. Assume that the coefficient $w_{-\mu, 0}^0$ of the weak expansion (7-7) is zero. (By Remark (ii) following Lemma 7.3, this condition is independent of the choice of ρ .) Then $u \equiv 0$. If $\mu \notin \frac{1}{2}\mathbb{N}_0$, then we can replace $w_{-\mu, 0}^0$ by $w_{\mu, 0, 0}^0$ in our assumption.*

Proof. Assume that $w_{\pm\mu, 0}^0 = 0$; here we consider the case of $w_{\mu, 0}^0 := w_{\mu, 0, 0}^0$ only when $\mu \notin \frac{1}{2}\mathbb{N}_0$. By Lemma 7.4, it suffices to prove that $w_{\pm\mu, 0}^r = 0$ for $r = 0, \dots, m$. This is a local statement and we use the half-plane model and the notation of the proof of Lemma 7.3. By (7-28), it then suffices to show that, if $\tilde{w}_{0; \pm\mu, 0}^0 = 0$ in the expansion (7-24), then $\tilde{w}_{i; \pm\mu, 0}^{2j} = 0$ for all i, j .

We argue by induction on $r = 2j + i = 0, \dots, m$. Assume first that $i = 0, j > 0$, and $\tilde{w}_{1; \pm\mu, 0}^{2(j-1)} = 0$. Then we plug (7-24) into (7-23) and consider the coefficient next to $z_0^{n/2 \pm \mu + 2j}$; this gives $\tilde{w}_{0; \pm\mu, 0}^{2j} = 0$ if, for $\lambda = \frac{1}{2}n \pm \mu + 2j$, the following constant is nonzero:

$$-\lambda^2 + \left(n - \frac{\lambda_j}{j}\right) \lambda + m - \sigma + \frac{\lambda_j}{j} (c_m - j) = (n + 2m - 2 - 4j)(\pm 2\mu - n - 2m + 2 + 4j). \tag{7-31}$$

⁵The existence of the weak expansion with known coefficients for elements in the image of \mathcal{P}_λ^- is directly related to the special case $m = 0$ of Lemma 6.8 and the existence of a weak expansion for scalar eigenfunctions of the Laplacian follows from the $m = 0$ case of Lemma 7.3. However, neither the surjectivity of the scalar Poisson operator nor the fact that eigenfunctions have nontrivial terms in their weak expansions follows from these statements.

We see immediately that (7-31) is nonzero unless $m = 2j$. For the case $m = 2j$, we can use (7-19) directly; taking the coefficient next to $z_0^{n/2 \pm \mu + m}$, we get $\tilde{w}_{0; \pm \mu, 0}^{2j} = 0$ as long as $\frac{1}{2}n \pm \mu + m \neq c_m$, or equivalently $\pm \mu \neq \frac{1}{2}n - 1$; the latter inequality is immediately true unless $n = 1$, and it is explicitly excluded by the statement of the present lemma when $n = 1$.

Similarly, assume that $i = 1$, $0 \leq 2j < m$, and $\tilde{w}_{0; \pm \mu, 0}^{2j} = 0$. Then we plug (7-24) into (7-22) and consider the coefficient next to $z_0^{n/2 \pm \mu + 2j + 1}$; this gives $\tilde{w}_{1; \pm \mu, 0}^{2j} = 0$ if, for $\lambda = \frac{1}{2}n \pm \mu + 2j + 1$, the following constant is nonzero:

$$-\lambda^2 + \left(n + 2 - \frac{\lambda_{j+1}}{j+1}\right)\lambda - n + m - 1 - \sigma + \frac{\lambda_{j+1}}{j+1}(c_m - j) = (n + 2m - 4 - 4j)(\pm 2\mu - n - 2m + 4 + 4j). \tag{7-32}$$

We see immediately that (7-32) is nonzero unless $m = 2j + 1$. For the case $m = 2j + 1$, we can use (7-18) directly; taking the coefficient next to $z_0^{n/2 \pm \mu + m}$, we get $\tilde{w}_{1; \pm \mu, 0}^{2j} = 0$ as long as $\frac{1}{2}n \pm \mu + m \neq c_m$, which we have already established is true. \square

We finish the section by the following statement, which immediately implies the surjectivity part of Theorem 6. Note that, for the lifts of elements of $\text{Eig}^m(-\lambda(n + \lambda) + m)$, we can take any $\delta < \frac{1}{2}$ below. The condition $\text{Re } \lambda < \frac{1}{2} - \delta$ for $m > 0$ follows from Lemma 6.1.

Corollary 7.6. *Let $u \in \rho^\delta L^2(\mathbb{B}^{n+1}; E^{(m)})$ be a trace-free symmetric m -cotensor with ρ a geodesic boundary defining function and $\delta \in (-\infty, \frac{1}{2})$, where the measure is the Euclidean Lebesgue measure on the ball. Assume that u is a nonzero divergence-free eigentensor for the Laplacian on hyperbolic space:*

$$\Delta u = (-\lambda(n + \lambda) + m)u, \quad \nabla^* u = 0, \tag{7-33}$$

with $\text{Re}(\lambda) < \frac{1}{2} - \delta$ and $\lambda \notin \mathcal{R}_m$, where \mathcal{R}_m is as defined in (5-20). Then, $u = \mathcal{P}_\lambda^-(w)$ for some $w \in H^{\text{Re}(\lambda) + \delta - 1/2}(\mathbb{S}^n; \otimes_S^m T^* \mathbb{S}^n)$. Moreover, if $\gamma^* u = u$ for some $\gamma \in G$, then $L_\gamma^* w = N_\gamma^{-\lambda - m} w$.

Proof. For the case $\text{Re}(\lambda) \geq -\frac{1}{2}n$ we set $\mu = \frac{1}{2}n + \lambda$ and apply Lemma 7.3; the distribution w will be given by $C(\lambda)w_{-\mu, 0}$ for some constant $C(\lambda)$ to be chosen, and this has the desired covariance with respect to elements of G by using (7-5) from the remark after Lemma 7.1.

To see that $u = \mathcal{P}_\lambda^-(w)$ for a certain $C(\lambda)$, it suffices to use the weak expansion in Lemma 6.8 and the identity (7-3) from the remark following Lemma 7.1, to deduce that $C(\lambda)B(\lambda)w_{-\mu, 0}$ appears as the leading coefficient of the power $\rho_0^{-\lambda}$ in the expansion of u , where $B(\lambda)$ is a nonzero constant times the factor appearing in (6-27); here ρ_0 is as defined in (3-34). (The factor $B(\lambda)$ does not depend on the point $v \in \mathbb{S}^n$ since the Poisson operator is equivariant under rotations of \mathbb{B}^{n+1} .) Then, choosing $C(\lambda) := B(\lambda)^{-1}$, we observe that u and $\mathcal{P}_\lambda^-(w)$ both satisfy (7-33) and have the same asymptotic coefficient of $\rho_0^{-\lambda}$ in their weak expansion (7-7); thus from Lemma 7.5 we have $u = \mathcal{P}_\lambda^-(w)$. Finally, for $\text{Re}(\lambda) < -\frac{1}{2}n$ with $\lambda \notin -\frac{1}{2}n - \frac{1}{2}\mathbb{N}_0$ we do the same thing but setting $\mu := -\frac{1}{2}n - \lambda$ in Lemma 7.3. \square

Appendix A: Some technical calculations

A1. Asymptotic expansions for certain integrals. In this subsection, we prove the following version of Hadamard regularization:

Lemma A.1. Fix $\chi \in C_0^\infty(\mathbb{R})$ and define for $\operatorname{Re} \alpha > 0$, $\beta \in \mathbb{C}$, and $\varepsilon > 0$,

$$F_{\alpha\beta}(\varepsilon) := \int_0^\infty t^{\alpha-1} (1+t)^{-\beta} \chi(\varepsilon t) dt.$$

If $\alpha - \beta \notin \mathbb{N}_0$, then $F_{\alpha\beta}(\varepsilon)$ has the following asymptotic expansion as $\varepsilon \rightarrow +0$:

$$F_{\alpha\beta}(\varepsilon) = \frac{\Gamma(\alpha)\Gamma(\beta - \alpha)}{\Gamma(\beta)} \chi(0) + \sum_{0 \leq j \leq \operatorname{Re}(\alpha - \beta)} c_j \varepsilon^{\beta - \alpha + j} + o(1) \tag{A-1}$$

for some constants c_j depending on χ .

Proof. We use the following identity obtained by integrating by parts:

$$\varepsilon \partial_\varepsilon F_{\alpha\beta}(\varepsilon) = \int_0^\infty t^\alpha (1+t)^{-\beta} \partial_t (\chi(\varepsilon t)) dt = (\beta - \alpha) F_{\alpha\beta}(\varepsilon) - \beta F_{\alpha, \beta+1}(\varepsilon). \tag{A-2}$$

By using the Taylor expansion of χ at zero, we also see that

$$\chi(\varepsilon t) = \chi(0) + \mathcal{O}(\varepsilon t);$$

given the following formula, obtained by the change of variables $s = (1+t)^{-1}$ and using the beta function,

$$\int_0^\infty t^{\alpha-1} (1+t)^{-\beta} dt = \frac{\Gamma(\alpha)\Gamma(\beta - \alpha)}{\Gamma(\beta)} \quad \text{if } \operatorname{Re} \beta > \operatorname{Re} \alpha > 0,$$

we see that

$$F_{\alpha\beta}(\varepsilon) = \frac{\Gamma(\alpha)\Gamma(\beta - \alpha)}{\Gamma(\beta)} \chi(0) + \mathcal{O}(\varepsilon) \quad \text{if } \operatorname{Re}(\beta - \alpha) > 1.$$

By applying this asymptotic expansion to $F_{\alpha, \beta+M}$ for a large integer M and iterating (A-2), we derive the expansion (A-1). □

For the next result, we need the following two calculations (see Section 4A for some of the notation used):

Lemma A.2. For each $\ell \geq 0$,

$$\int_{\mathbb{S}^{n-1}} (\otimes^{2\ell} \eta) dS(\eta) = \frac{2\pi^{(n-1)/2} \Gamma(\ell + \frac{1}{2})}{\Gamma(\ell + \frac{1}{2}n)} S(\otimes^\ell I),$$

where $I = \sum_{j=1}^n \partial_j \otimes \partial_j$.

Proof. Since both sides are symmetric tensors, it suffices to show that, for each $x \in \mathbb{R}^n$,

$$\int_{\mathbb{S}^{n-1}} (x \cdot \eta)^{2\ell} dS(\eta) = \frac{2\pi^{(n-1)/2} \Gamma(\ell + \frac{1}{2})}{\Gamma(\ell + \frac{1}{2}n)} |x|^{2\ell}.$$

Without loss of generality (using homogeneity and rotational invariance), we may assume that $x = \partial_1$. Then, using polar coordinates and Fubini's theorem, we have

$$\frac{1}{2} \Gamma(\ell + \frac{1}{2}n) \int_{\mathbb{S}^{n-1}} \eta_1^{2\ell} dS(\eta) = \int_{\mathbb{R}^n} e^{-|\eta|^2} \eta_1^{2\ell} d\eta = \pi^{(n-1)/2} \Gamma(\ell + \frac{1}{2}),$$

finishing the proof. □

Lemma A.3. For each $\eta \in \mathbb{R}^n$, define the linear map $\mathcal{C}_\eta : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by

$$\mathcal{C}_\eta(\tilde{\eta}) = \tilde{\eta} - \frac{2}{1 + |\eta|^2}(\tilde{\eta} \cdot \eta)\eta.$$

Then, for each $A_1, A_2 \in \otimes_{\mathbb{S}}^m \mathbb{R}^n$ with $\mathcal{T}(A_1) = \mathcal{T}(A_2) = 0$, and each $r \geq 0$, we have

$$\int_{\mathbb{S}^{n-1}} \langle (\otimes^m \mathcal{C}_{r\eta})A_1, A_2 \rangle dS(\eta) = 2\pi^{n/2} \sum_{\ell=0}^m \frac{m!}{(m-\ell)! \Gamma(\frac{1}{2}n + \ell)} \left(-\frac{r^2}{1+r^2}\right)^\ell \langle A_1, A_2 \rangle.$$

Proof. We have

$$\mathcal{C}_{r\eta} = \text{Id} - \frac{2r^2}{1+r^2} \eta^* \otimes \eta,$$

where $\eta^* \in (\mathbb{R}^n)^*$ is the dual to η by the standard metric. Then

$$\int_{\mathbb{S}^{n-1}} \langle (\otimes^m \mathcal{C}_{r\eta})A_1, A_2 \rangle dS(\eta) = \int_{\mathbb{S}^{n-1}} \left\langle \otimes^m \left(I - \frac{2r^2}{1+r^2} \eta \otimes \eta \right), \sigma(A_1 \otimes A_2) \right\rangle dS(\eta),$$

where σ is the operator defined by

$$\sigma(\eta_1 \otimes \dots \otimes \eta_m \otimes \eta'_1 \otimes \dots \otimes \eta'_m) = \eta_1 \otimes \eta'_1 \otimes \dots \otimes \eta_m \otimes \eta'_m.$$

We use [Lemma A.2](#), a binomial expansion, and the fact that the A_j are symmetric, to calculate

$$\begin{aligned} & \int_{\mathbb{S}^{n-1}} \left\langle \otimes^m \left(I - \frac{2r^2}{1+r^2} \eta \otimes \eta \right), \sigma(A_1 \otimes A_2) \right\rangle dS(\eta) \\ &= \sum_{\ell=0}^m \frac{m!}{\ell!(m-\ell)!} \left(-\frac{2r^2}{1+r^2}\right)^\ell \int_{\mathbb{S}^{n-1}} \langle (\otimes^{2\ell} \eta) \otimes (\otimes^{m-\ell} I), \sigma(A_1 \otimes A_2) \rangle dS(\eta) \\ &= 2\pi^{(n-1)/2} \sum_{\ell=0}^m \frac{m!}{\ell!(m-\ell)!} \cdot \frac{\Gamma(\ell + \frac{1}{2})}{\Gamma(\ell + \frac{1}{2}n)} \left(-\frac{2r^2}{1+r^2}\right)^\ell \langle S(\otimes^\ell I) \otimes (\otimes^{m-\ell} I), \sigma(A_1 \otimes A_2) \rangle. \end{aligned}$$

Since $\mathcal{T}(A_1) = \mathcal{T}(A_2) = 0$, we can compute

$$\langle S(\otimes^\ell I) \otimes (\otimes^{m-\ell} I), \sigma(A_1 \otimes A_2) \rangle = \frac{2^\ell (\ell!)^2}{(2\ell)!} \langle A_1, A_2 \rangle.$$

Here $2^\ell (\ell!)^2 / (2\ell)!$ is the proportion of permutations τ of 2ℓ elements that satisfy, for each j , that $\tau(2j-1) + \tau(2j)$ is odd. It remains to calculate

$$\sum_{\ell=0}^m \frac{m!}{\ell!(m-\ell)!} \cdot \frac{\Gamma(\ell + \frac{1}{2})}{\Gamma(\ell + \frac{1}{2}n)} \cdot \frac{2^\ell (\ell!)^2}{(2\ell)!} r^{2\ell} = \sum_{\ell=0}^m \frac{\sqrt{\pi} m!}{(m-\ell)! \Gamma(\ell + \frac{1}{2}n)} \left(\frac{1}{2}r\right)^\ell. \quad \square$$

We can now state the following asymptotic formula, used in the proof of [Lemma 5.11](#):

Lemma A.4. *Let $\chi \in C_0^\infty(\mathbb{R})$ be equal to 1 near 0, and take $A_1, A_2 \in \otimes_S^m \mathbb{R}^n$ satisfying $\mathcal{T}(A_1) = \mathcal{T}(A_2) = 0$. Then, for $\lambda \in \mathbb{C}, \lambda \notin -(\frac{1}{2}n + \mathbb{N}_0)$, we have, as $\varepsilon \rightarrow +0$,*

$$\int_{\mathbb{R}^n} \chi(\varepsilon|\eta|)(1 + |\eta|^2)^{-\lambda-n} \langle (\otimes^m \mathcal{C}_\eta) A_1, A_2 \rangle d\eta$$

$$= \pi^{n/2} \frac{\Gamma(\frac{1}{2}n + \lambda)}{(n + \lambda + m - 1)\Gamma(n - 1 + \lambda)} \langle A_1, A_2 \rangle + \sum_{0 \leq j \leq -\operatorname{Re} \lambda - n/2} c_j \varepsilon^{n+2\lambda+2j} + o(1)$$

for some constants c_j .

Proof. We write, using the change of variables $\eta = \sqrt{t}\theta, \theta \in \mathbb{S}^n$, and $\chi(s) = \tilde{\chi}(s^2)$, and by Lemma A.3,

$$\int_{\mathbb{R}^n} \chi(\varepsilon|\eta|)(1 + |\eta|^2)^{-\lambda-n} \langle (\otimes^m \mathcal{C}_\eta) A_1, A_2 \rangle d\eta$$

$$= \frac{1}{2} \int_0^\infty \tilde{\chi}(\varepsilon^2 t) t^{\frac{1}{2}n-1} (1+t)^{-\lambda-n} \int_{\mathbb{S}^{n-1}} \langle (\otimes^m \mathcal{C}_{\sqrt{t}\theta}) A_1, A_2 \rangle dS(\theta) dt$$

$$= \pi^{n/2} \sum_{\ell=0}^m \frac{(-1)^\ell m!}{(m-\ell)! \Gamma(\frac{1}{2}n + \ell)} \langle A_1, A_2 \rangle \int_0^\infty \tilde{\chi}(\varepsilon^2 t) t^{n/2+\ell-1} (1+t)^{-\lambda-n-\ell} dt.$$

We now apply Lemma A.1 to get the required asymptotic expansion. The constant term in the expansion is $\langle A_1, A_2 \rangle$ times

$$\pi^{n/2} \Gamma(\frac{1}{2}n + \lambda) \sum_{\ell=0}^m \frac{(-1)^\ell m!}{(m-\ell)! \Gamma(n + \lambda + \ell)} = \pi^{n/2} (-1)^m m! \Gamma(\frac{1}{2}n + \lambda) \sum_{\ell=0}^m \frac{(-1)^\ell}{\ell! \Gamma(n + \lambda + m - \ell)}. \tag{A-3}$$

We now use the binomial expansion

$$\frac{(1-t)^{n+\lambda+m-1}}{\Gamma(n + \lambda + m)} = \sum_{\ell=0}^\infty \frac{(-1)^\ell}{\ell! \Gamma(n + \lambda + m - \ell)} t^\ell$$

and the sum in the last line of (A-3) is the t^m coefficient of

$$(1-t)^{-1} \frac{(1-t)^{n+\lambda+m-1}}{\Gamma(n + \lambda + m)} = \frac{(1-t)^{n+\lambda+m-2}}{\Gamma(n + \lambda + m)} = \frac{1}{n + \lambda + m - 1} \sum_{j=0}^\infty \frac{(-1)^j}{j! \Gamma(n + \lambda + m - j - 1)} t^j;$$

this finishes the proof. □

A2. The Jacobian of Ψ . Here we compute the Jacobian of the map $\Psi : \mathcal{E} \rightarrow S_\Delta^2 \mathbb{H}^{n+1}$ appearing in the proof of Lemma 5.11, proving (5-31). By the G -equivariance of Ψ , we may assume that $x = \partial_0, \xi = \partial_1, \eta = \sqrt{s} \partial_2$ for some $s \geq 0$. We then consider the following volume 1 basis of $T_{(x,\xi,\eta)} \mathcal{E}$:

$$X_1 = (\partial_1, \partial_0, 0), \quad X_2 = (\partial_2, 0, \sqrt{s} \partial_0), \quad X_3 = (0, \partial_2, -\sqrt{s} \partial_1), \quad X_4 = (0, 0, \partial_2);$$

$$\partial_{x_j}, \partial_{\xi_j}, \partial_{\eta_j}, \quad 3 \leq j \leq n + 1.$$

We have $\Psi(x, \xi, \eta) = (y, \eta_-, \eta_+)$, where

$$y = (\sqrt{s+1}, 0, \sqrt{s}, 0, \dots, 0), \quad \eta_{\pm} = \left(\mp \frac{s}{\sqrt{s+1}}, \frac{1}{\sqrt{s+1}}, \mp \sqrt{s}, 0, \dots, 0 \right).$$

Then we can consider the following volume 1 basis for $T_{(y, \eta_-, \eta_+)} S_{\Delta}^2 \mathbb{H}^{n+1}$:

$$\begin{aligned} Y_1 &= \left(\partial_1, \frac{y}{\sqrt{s+1}}, \frac{y}{\sqrt{s+1}} \right), & Y_2 &= \left(\sqrt{s} \partial_0 + \sqrt{s+1} \partial_2, \frac{\sqrt{s}}{\sqrt{s+1}} y, -\frac{\sqrt{s}}{\sqrt{s+1}} y \right), \\ Y_3 &= \frac{(0, \sqrt{s} \partial_0 - \sqrt{s} \partial_1 + \sqrt{s+1} \partial_2, 0)}{\sqrt{s+1}}, & Y_4 &= \frac{(0, 0, \sqrt{s} \partial_0 + \sqrt{s} \partial_1 + \sqrt{s+1} \partial_2)}{\sqrt{s+1}}; \\ & & & \partial_{y_j}, \partial_{v_{-j}}, \partial_{v_{+j}}, \quad 3 \leq j \leq n+1. \end{aligned}$$

Then the differential $d\Psi(x, \xi, \eta)$ maps

$$\begin{aligned} X_1 &\mapsto \sqrt{s+1} Y_1 - \sqrt{s} Y_3 - \sqrt{s} Y_4, \\ X_2 &\mapsto Y_2, \\ X_3 &\mapsto -\sqrt{s} Y_1 + \sqrt{s+1} Y_3 + \sqrt{s+1} Y_4, \\ X_4 &\mapsto \frac{1}{\sqrt{s+1}} Y_2 + \frac{1}{s+1} Y_3 - \frac{1}{s+1} Y_4. \end{aligned}$$

Moreover, for $3 \leq j \leq n+1$, $d\Psi(x, \xi, \eta)$ maps linear combinations of $\partial_{x_j}, \partial_{\xi_j}, \partial_{\eta_j}$ to linear combinations of $\partial_{y_j}, \partial_{v_{-j}}, \partial_{v_{+j}}$ by the matrix $A(s)$. The identity (5-31) now follows by a direct calculation.

A3. An identity for harmonic polynomials. We give a technical lemma which is used in the proof of Lemma 6.8 (injectivity of the Poisson kernel).

Lemma A.5. *Let P be a harmonic homogeneous polynomial of order m in \mathbb{R}^n ; then, for $r \leq m$, we have for all $x \in \mathbb{R}^n$ that*

$$\Delta_{\zeta}^r P(x - \zeta \langle \zeta, x \rangle) |_{\zeta=0} = 2^r \frac{m! r!}{(m-r)!} P(x).$$

Proof. By homogeneity, it suffices to choose $|x| = 1$. We set $t = \langle \zeta, x \rangle$ and $u = \zeta - tx$, and $P(x - \zeta \langle \zeta, x \rangle)$, viewed in the (t, u) coordinates, is the homogeneous polynomial $(t, u) \mapsto P((1-t^2)x - tu)$. Now, we write, for all $u \in (\mathbb{R}x)^{\perp}$ and $t > 0$,

$$P(tx - u) = \sum_{j=0}^m t^{m-j} P_j(u),$$

where P_j is a homogeneous polynomial of degree j in $u \in (\mathbb{R}x)^{\perp}$, and, since the Laplacian Δ_{ζ} written in the t, u coordinates is $-\partial_t^2 + \Delta_u$, the condition $\Delta_x P = 0$ can be rewritten

$$\Delta_u P_j(u) = (m-j+2)(m-j+1)P_{j-2}(u), \quad \Delta_u P_1(u) = \Delta_u P_0 = 0,$$

which gives, for all j and $\ell \geq 1$,

$$\Delta_u^{\ell} P_{2\ell}(u) = m(m-1) \cdots (m-2\ell+1)P_0, \quad \Delta^j P_{2\ell-1}(u) |_{u=0} = 0.$$

We write $\Delta_\zeta^r = \sum_{k=0}^r (r!/(k!(r-k)!))(-1)^k \partial_t^{2k} \Delta_u^{r-k}$ and, using parity and homogeneity considerations, we have

$$\begin{aligned} \Delta_\zeta^r P(x - \zeta \langle \zeta, x \rangle) |_{\zeta=0} &= \sum_{k=0}^r \frac{(-1)^k r!}{k!(r-k)!} \sum_{2j \leq m} [\partial_t^{2k} ((1-t^2)^{m-2j} t^{2j}) \Delta_u^{r-k} P_{2j}(u)] |_{(t,u)=0} \\ &= \sum_{\max(0, r-m/2) \leq k \leq r} \frac{(-1)^k r!}{k!(r-k)!} (\partial_t^{2k} ((1-t^2)^{m-2(r-k)} t^{2(r-k)})) |_{t=0} \Delta_u^{r-k} P_{2(r-k)} \\ &= P_0 \cdot \frac{m!r!}{(m-r)!} \sum_{r/2 \leq k \leq r} \frac{(-1)^{k+r} (2k)!}{k!(r-k)!(2k-r)!} = 2^r \frac{m!r!}{(m-r)!} P_0 \end{aligned}$$

and P_0 is the constant given by $P(x)$. Here we used the identity

$$\sum_{r/2 \leq k \leq r} \frac{(-1)^{k+r} (2k)!}{k!(r-k)!(2k-r)!} = \sum_{0 \leq k \leq r/2} (-1)^k \frac{r!}{k!(r-k)!} \cdot \frac{(2r-2k)!}{r!(r-2k)!} = 2^r,$$

which holds because both sides are equal to the t^r coefficient of the product

$$(1-t^2)^r \cdot (1-t)^{-1-r} = \frac{(1+t)^r}{1-t} :$$

since

$$(1-t)^{-1-r} = \frac{1}{r!} d_t^r (1-t)^{-1} = \sum_{j=0}^{\infty} \frac{(j+r)!}{j!r!} t^j,$$

the t^r coefficient of $(1+t)^r/(1-t)$ equals the sum of the t^0, t^1, \dots, t^r coefficients of $(1+t)^r$, or simply $(1+1)^r = 2^r$. □

Appendix B: The special case of dimension 2

We explain how the argument of [Section 2A](#) fits into the framework of [Sections 3](#) and [4](#). In dimension 2 it is more standard to use the upper half-plane model

$$\mathbf{H}^2 := \{w \in \mathbb{C} \mid \text{Im } w > 0\},$$

which is related to the half-space model of [Section 3A](#) by the formula $w = -z_1 + iz_0$.

The group of all isometries of \mathbf{H}^2 is $\text{PSL}(2; \mathbb{R})$, the quotient of $\text{SL}(2; \mathbb{R})$ by the group generated by the matrix $-\text{Id}$, and the action of $\text{PSL}(2; \mathbb{R})$ on \mathbf{H}^2 is by Möbius transformations:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot z = \frac{az+b}{cz+d}, \quad z \in \mathbf{H}^2 \subset \mathbb{C}.$$

Under the identifications (3-2) and (3-5), this action corresponds to the action of $\text{PSO}(1, 2)$ on $\mathbb{H}^2 \subset \mathbb{R}^{1,2}$ by the group isomorphism $\text{PSL}(2; \mathbb{R}) \rightarrow \text{PSO}(1, 2)$ defined by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{pmatrix} \frac{1}{2}(a^2 + b^2 + c^2 + d^2) & \frac{1}{2}(a^2 - b^2 + c^2 - d^2) & -ab - cd \\ \frac{1}{2}(a^2 + b^2 - c^2 - d^2) & \frac{1}{2}(a^2 - b^2 - c^2 + d^2) & cd - ab \\ -ac - bd & bd - ac & ad + bc \end{pmatrix}. \tag{B-1}$$

The induced Lie algebra isomorphism maps the vector fields X, U_-, U_+ of (2-1) to the fields X, U_1^-, U_1^+ of (3-6), (3-7).

The horocyclic operators $U_{\pm} : \mathcal{D}'(S\mathbb{H}^2) \rightarrow \mathcal{D}'(S\mathbb{H}^2; \mathcal{E}^*)$ of Section 4B (and analogously horocyclic operators of higher orders) then take the form

$$U_{\pm}u = (U_{\pm}u)\eta^*,$$

where η^* is the dual to the section $\eta \in C^\infty(S\mathbb{H}^2; \mathcal{E})$ defined as follows: for $(x, \xi) \in S\mathbb{H}^2$, $\eta(x, \xi)$ is the unique vector in $T_x\mathbb{H}^2$ such that (ξ, η) is a positively oriented orthonormal frame. Note also that $\eta(x, \xi) = \pm \mathcal{A}_{\pm}(x, \xi) \cdot \zeta(B_{\pm}(x, \xi))$, where $\mathcal{A}_{\pm}(x, \xi)$ is as defined in Section 3F and $\zeta(v) \in T_v\mathbb{S}^1, v \in \mathbb{S}^1$, is the result of rotating v counterclockwise by $\frac{1}{2}\pi$; therefore, if we use η and ζ to trivialize the relevant vector bundles, then the operators Q_{\pm} of (4-26) are simply the pullback operators by B_{\pm} , up to multiplication by ± 1 .

Appendix C: Eigenvalue asymptotics for symmetric tensors

C1. Weyl law. In this section, we prove the following asymptotic of the counting function for trace-free, divergence-free tensors (see Sections 4A and 6A for the notation):

Proposition C.1. *If (M, g) is a compact Riemannian manifold of dimension $n + 1$ and constant sectional curvature -1 , and if*

$$\text{Eig}^m(\sigma) = \{u \in C^\infty(M; \otimes_S^m T^*M) \mid \Delta u = \sigma u, \nabla^*u = 0, \mathcal{T}(u) = 0\},$$

then the following Weyl law holds as $R \rightarrow \infty$:

$$\sum_{\sigma \leq R^2} \dim \text{Eig}^m(\sigma) = c_0(n)(c_1(n, m) - c_1(n, m - 2)) \text{Vol}(M)R^{n+1} + \mathcal{O}(R^n),$$

where $c_0(n) = ((2\sqrt{\pi})^{-n-1} / \Gamma(\frac{1}{2}(n + 3)))$ and $c_1(n, m) = (m + n - 1)! / (m!(n - 1)!)$ is the dimension of the space of homogeneous polynomials of order m in n variables. (We put $c_1(n, m) := 0$ for $m < 0$.)

Remark. The constant $c_2(n, m) := c_1(n, m) - c_1(n, m - 2)$ is the dimension of the space of harmonic homogeneous polynomials of order m in n variables. We have

$$c_2(n, 0) = 1, \quad c_2(n, 1) = n.$$

For $m \geq 2$, we have $c_2(n, m) > 0$ if and only if $n > 1$.

The proof of Proposition C.1 uses the following two technical lemmas:

Lemma C.2. Take $u \in \mathcal{D}'(M; \otimes_S^m T^*M)$. Then, denoting $D = \mathcal{S} \circ \nabla$ as in Section 6A,

$$[\Delta, \nabla^*]u = (2 - 2m - n)\nabla^*u - 2(m - 1)D(\mathcal{T}(u)), \tag{C-1}$$

$$[\Delta, D]u = (2m + n)Du + 2m\mathcal{S}(g \otimes \nabla^*u). \tag{C-2}$$

Proof. We have

$$\Delta \nabla^*u = \mathcal{T}^2(\nabla^3u), \quad \nabla^* \Delta u = \mathcal{T}^2(\tau_{1 \leftrightarrow 3} \nabla^3u),$$

where $\tau_{j \leftrightarrow k}v$ denotes the result of swapping the j -th and k -th indices in a cotensor v . We have

$$\text{Id} - \tau_{1 \leftrightarrow 3} = (\text{Id} - \tau_{1 \leftrightarrow 2}) + \tau_{1 \leftrightarrow 2}(\text{Id} - \tau_{2 \leftrightarrow 3}) + \tau_{1 \leftrightarrow 2}\tau_{2 \leftrightarrow 3}(\text{Id} - \tau_{1 \leftrightarrow 2});$$

therefore (using that $\mathcal{T}\tau_{1 \leftrightarrow 2} = \mathcal{T}$)

$$[\Delta, \nabla^*]u = \mathcal{T}^2(\nabla(\text{Id} - \tau_{1 \leftrightarrow 2})\nabla^2u + \tau_{2 \leftrightarrow 3}(\text{Id} - \tau_{1 \leftrightarrow 2})\nabla^3u).$$

Since M has sectional curvature -1 , we have, for any cotensor v of rank m ,

$$(\text{Id} - \tau_{1 \leftrightarrow 2})\nabla^2v = \sum_{\ell=1}^m (\tau_{1 \leftrightarrow \ell+2} - \tau_{2 \leftrightarrow \ell+2})(g \otimes v).$$

Then we compute (using that $\mathcal{T}(\tau_{2 \leftrightarrow 3}\tau_{1 \leftrightarrow 3}) = \mathcal{T}(\tau_{2 \leftrightarrow 3})$)

$$[\Delta, \nabla^*]u = \mathcal{T}^2\left(\tau_{2 \leftrightarrow 3} - \text{Id} + \sum_{\ell=1}^m ((\tau_{2 \leftrightarrow \ell+3} - \tau_{3 \leftrightarrow \ell+3})\tau_{1 \leftrightarrow 3} + \tau_{2 \leftrightarrow 3}(\tau_{1 \leftrightarrow \ell+3} - \tau_{2 \leftrightarrow \ell+3}))\right)(g \otimes \nabla u).$$

Now,

$$\begin{aligned} \mathcal{T}^2(g \otimes \nabla u) &= \mathcal{T}^2(\tau_{2 \leftrightarrow 4}\tau_{1 \leftrightarrow 3}(g \otimes \nabla u)) = \mathcal{T}^2(\tau_{2 \leftrightarrow 3}\tau_{1 \leftrightarrow 4}(g \otimes \nabla u)) = -(n + 1)\nabla^*u, \\ \mathcal{T}^2(\tau_{2 \leftrightarrow 3}(g \otimes \nabla u)) &= \mathcal{T}^2(\tau_{3 \leftrightarrow 4}\tau_{1 \leftrightarrow 3}(g \otimes \nabla u)) = \mathcal{T}^2(\tau_{2 \leftrightarrow 3}\tau_{2 \leftrightarrow 4}(g \otimes \nabla u)) = -\nabla^*u, \end{aligned}$$

and, since u is symmetric, for $1 < \ell \leq m$,

$$\begin{aligned} \mathcal{T}^2(\tau_{2 \leftrightarrow \ell+3}\tau_{1 \leftrightarrow 3}(g \otimes \nabla u)) &= \mathcal{T}^2(\tau_{2 \leftrightarrow 3}\tau_{1 \leftrightarrow \ell+3}(g \otimes \nabla u)) = -\nabla^*u, \\ \mathcal{T}^2(\tau_{3 \leftrightarrow \ell+3}\tau_{1 \leftrightarrow 3}(g \otimes \nabla u)) &= \mathcal{T}^2(\tau_{2 \leftrightarrow 3}\tau_{2 \leftrightarrow \ell+3}(g \otimes \nabla u)) = \tau_{1 \leftrightarrow \ell-1}\nabla(\mathcal{T}(u)). \end{aligned}$$

We then compute

$$[\Delta, \nabla^*]u = (2 - 2m - n)\nabla^*u - 2 \sum_{\ell=1}^{m-1} \tau_{1 \leftrightarrow \ell} \nabla(\mathcal{T}(u)),$$

finishing the proof of (C-1). The identity (C-2) follows from (C-1) by taking the adjoint on the space of symmetric tensors. □

Lemma C.3. Denote by $\tilde{\pi}_m : \otimes_S^m T^*M \rightarrow \otimes_S^m T^*M$ the orthogonal projection onto the space $\ker \mathcal{T}$ of trace-free tensors. Then, for each m , the space

$$F^m := \{v \in C^\infty(M; \otimes_S^m T^*M) \mid \mathcal{T}(v) = 0, \tilde{\pi}_{m+1}(Dv) = 0\} \tag{C-3}$$

is finite-dimensional.

Proof. The space F^m is contained in the kernel of the operator

$$P_m := \nabla^* \tilde{\pi}_{m+1} D$$

acting on trace-free sections of $\otimes_S^m T^*M$. By [Dairbekov and Sharafutdinov 2010, Lemma 5.2], the operator P_m is elliptic; therefore, its kernel is finite-dimensional. \square

We now prove Proposition C.1. For each $m \geq 0$ and $s \in \mathbb{R}$, denote

$$W^m(\sigma) := \{u \in \mathcal{D}'(M; \otimes_S^m T^*M) \mid \Delta u = \sigma u, \mathcal{T}(u) = 0\}.$$

The operator Δ acting on trace-free symmetric tensors is elliptic and, in fact, its principal symbol coincides with that of the scalar Laplacian: $p(x, \xi) = |\xi|_g^2$. It follows that the $W^m(\sigma)$ are finite-dimensional and consist of smooth sections. By the general argument of [Hörmander 1994, Section 17.5] (see also [Dimassi and Sjöstrand 1999, Theorem 10.1; Zworski 2012, Theorem 6.8] — all of these arguments adapt straightforwardly to the case of operators with diagonal principal symbols acting on vector bundles), we have the following Weyl law:

$$\sum_{\sigma \leq R^2} \dim W^m(\sigma) = c_0(n)(c_1(n+1, m) - c_1(n+1, m-2)) \text{Vol}(M)R^{n+1} + \mathcal{O}(R^n); \tag{C-4}$$

here $c_1(n+1, m) - c_1(n+1, m+2)$ is the dimension of the vector bundle on which we consider the operator Δ .

By (C-1), for $m \geq 1$ the divergence operator acts as

$$\nabla^* : W^m(\sigma) \rightarrow W^{m-1}(\sigma + 2 - 2m - n). \tag{C-5}$$

This operator is surjective except at finitely many points σ :

Lemma C.4. *Let $C_1 = \dim F^{m-1}$, where F^{m-1} is as defined in (C-3). Then the number of values σ such that (C-5) is not surjective does not exceed C_1 .*

Proof. Assume that (C-5) is not surjective for some σ . Then there exists nonzero $v \in W^{m-1}(\sigma + 2 - 2m - n)$ which is orthogonal to $\nabla^*(W^m(\sigma))$. Since the spaces $W^{m-1}(\sigma)$ are mutually orthogonal, we see from (C-5) that v is also orthogonal to $\nabla^*(W^m(\sigma))$ for all $\sigma \neq \sigma$. It follows that, for each σ and each $u \in W^m(\sigma)$, we have $\langle Dv, u \rangle_{L^2} = 0$. Since $\bigoplus_{\sigma} W^m(\sigma)$ is dense in the space of trace-free tensors, we see that, for each $u \in C^\infty(M; \otimes_S^m T^*M)$ with $\mathcal{T}(u) = 0$, we have $\langle Dv, u \rangle_{L^2} = 0$, which implies $v \in F^{m-1}$. It remains to note that F^{m-1} can have a nontrivial intersection with at most C_1 of the spaces $W^{m-1}(\sigma + 2 - 2m - n)$. \square

Since $\text{Eig}^m(\sigma)$ is the kernel of (C-5), we have

$$\dim \text{Eig}^m(\sigma) \geq \dim W^m(\sigma) - \dim W^{m-1}(\sigma + 2 - 2m - n),$$

and this inequality is an equality if (C-5) is surjective. We then see that, for some constant C_2 independent of R ,

$$\begin{aligned} \sum_{\sigma \leq R^2} \dim W^m(\sigma) - \sum_{\sigma \leq R^2+2-2m-n} \dim W^{m-1}(\sigma) &\leq \sum_{\sigma \leq R^2} \dim \text{Eig}^m(\sigma) \\ &\leq C_2 + \sum_{\sigma \leq R^2} \dim W^m(\sigma) - \sum_{\sigma \leq R^2+2-2m-n} \dim W^{m-1}(\sigma), \end{aligned}$$

and Proposition C.1 now follows from (C-4) and the identity $c_1(n + 1, m) - c_1(n + 1, m - 1) = c_1(n, m)$.

C2. The case $m = 1$. In this section, we describe the space $\text{Eig}^1(\sigma)$ in terms of Hodge theory; see, for instance, [Petersen 2006, Section 7.2] for the notation used. Note that symmetric cotensors of order 1 are exactly differential 1-forms on M . Since the operator $\nabla : C^\infty(M) \rightarrow C^\infty(M; T^*M)$ is equal to the operator d on 0-forms, we have

$$\text{Eig}^1(\sigma) = \{u \in \Omega^1(M) \mid \Delta u = \sigma u, \delta u = 0\}.$$

Here $\Delta = \nabla^* \nabla$; using that M has sectional curvature -1 , we write Δ in terms of the Hodge Laplacian $\Delta_\Omega := d\delta + \delta d$ on 1-forms using the following Weitzenböck formula [Petersen 2006, Corollary 7.21]:

$$\Delta u = (\Delta_\Omega + n)u, \quad u \in \Omega^1(M).$$

We then see that

$$\text{Eig}^1(\sigma) = \{u \in \Omega^1(M) \mid \Delta_\Omega u = (\sigma - n)u, \delta u = 0\}. \tag{C-6}$$

Finally, let us consider the case $n = 1$. The Hodge star operator acts from $\Omega^1(M)$ to itself, and we see that, for $\sigma \neq 1$,

$$\begin{aligned} \text{Eig}^1(\sigma) &= \{ *u \mid u \in \Omega^1(M), \Delta_\Omega u = (\sigma - 1)u, du = 0 \} \\ &= \{ *(df) \mid f \in C^\infty(M), \Delta f = (\sigma - 1)f \}. \end{aligned} \tag{C-7}$$

Note that $*(df)$ can be viewed as the Hamiltonian field of f with respect to the naturally induced symplectic form (that is, volume form) on M .

Acknowledgements

We would like to thank Maciej Zworski, Richard Melrose, Steve Zelditch, Rafe Mazzeo, and Kiril Datchev for many useful discussions and suggestions regarding this project, and Tobias Weich for corrections and references. Dyatlov would like to thank MSRI (NSF grant 0932078 000, Fall 2013), where part of this work was done; another part was supported by the NSF grant DMS-1201417, and part of this project was completed during the period Dyatlov served as a Clay Research Fellow. Guillarmou and Faure have been supported by “Agence Nationale de la Recherche” under the grant ANR-13-BS01-0007-01.

References

- [Abramowitz and Stegun 1964] M. Abramowitz and I. A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards Applied Mathematics Series **55**, U.S. Government Printing Office, Washington, D.C., 1964. [MR 29 #4914](#) [Zbl 0171.38503](#)
- [Anantharaman and Zelditch 2007] N. Anantharaman and S. Zelditch, “Patterson–Sullivan distributions and quantum ergodicity”, *Ann. Henri Poincaré* **8**:2 (2007), 361–426. [MR 2008f:58028](#) [Zbl 1187.81175](#)
- [Anantharaman and Zelditch 2012] N. Anantharaman and S. Zelditch, “Intertwining the geodesic flow and the Schrödinger group on hyperbolic surfaces”, *Math. Ann.* **353**:4 (2012), 1103–1156. [MR 2944025](#) [Zbl 06065228](#)
- [van den Ban and Schlichtkrull 1987] E. P. van den Ban and H. Schlichtkrull, “Asymptotic expansions and boundary values of eigenfunctions on Riemannian symmetric spaces”, *J. Reine Angew. Math.* **380** (1987), 108–165. [MR 89g:43010](#) [Zbl 0631.58028](#)
- [Bony and Häfner 2008] J.-F. Bony and D. Häfner, “Decay and non-decay of the local energy for the wave equation on the de Sitter–Schwarzschild metric”, *Comm. Math. Phys.* **282**:3 (2008), 697–719. [MR 2010h:58041](#) [Zbl 1159.35007](#)
- [Bunke and Olbrich 1995] U. Bunke and M. Olbrich, *Selberg zeta and theta functions: a differential operator approach*, Mathematical Research **83**, Akademie-Verlag, Berlin, 1995. [MR 97c:11088](#) [Zbl 0831.58001](#)
- [Bunke and Olbrich 1997] U. Bunke and M. Olbrich, “Fuchsian groups of the second kind and representations carried by the limit set”, *Invent. Math.* **127**:1 (1997), 127–154. [MR 98e:11065](#) [Zbl 0880.30035](#)
- [Bunke and Olbrich 1999] U. Bunke and M. Olbrich, “Group cohomology and the singularities of the Selberg zeta function associated to a Kleinian group”, *Ann. of Math. (2)* **149**:2 (1999), 627–689. [MR 2000f:11110](#) [Zbl 0969.11019](#)
- [Bunke and Olbrich 2001] U. Bunke and M. Olbrich, “Regularity of invariant distributions”, preprint, 2001. [arXiv 0103144](#)
- [Butterley and Liverani 2007] O. Butterley and C. Liverani, “Smooth Anosov flows: correlation spectra and stability”, *J. Mod. Dyn.* **1**:2 (2007), 301–322. [MR 2008a:37035](#) [Zbl 1144.37011](#)
- [Dairbekov and Sharafutdinov 2010] N. S. Dairbekov and V. A. Sharafutdinov, “Conformal Killing symmetric tensor fields on Riemannian manifolds”, *Mat. Tr.* **13**:1 (2010), 85–145. In Russian; translated in *Sib. Adv. Math.* **21**:1 (2011), 1–41. [MR 2011j:53052](#) [Zbl 1249.53050](#)
- [Datchev et al. 2014] K. Datchev, S. Dyatlov, and M. Zworski, “Sharp polynomial bounds on the number of Pollicott–Ruelle resonances”, *Ergodic Theory Dynam. Systems* **34**:4 (2014), 1168–1183. [MR 3227152](#) [Zbl 06359408](#)
- [Dimassi and Sjöstrand 1999] M. Dimassi and J. Sjöstrand, *Spectral asymptotics in the semi-classical limit*, London Mathematical Society Lecture Note Series **268**, Cambridge University Press, 1999. [MR 2001b:35237](#) [Zbl 0926.35002](#)
- [Dolgopyat 1998] D. Dolgopyat, “On decay of correlations in Anosov flows”, *Ann. of Math. (2)* **147**:2 (1998), 357–390. [MR 99g:58073](#) [Zbl 0911.58029](#)
- [Dyatlov 2012] S. Dyatlov, “Asymptotic distribution of quasi-normal modes for Kerr–de Sitter black holes”, *Ann. Henri Poincaré* **13**:5 (2012), 1101–1166. [MR 2935116](#) [Zbl 1246.83111](#)
- [Dyatlov 2015] S. Dyatlov, “Asymptotics of linear waves and resonances with applications to black holes”, *Comm. Math. Phys.* **335**:3 (2015), 1445–1485. [MR 3320319](#) [Zbl 06421372](#)
- [Dyatlov and Zworski 2015] S. Dyatlov and M. Zworski, “Dynamical zeta functions for Anosov flows via microlocal analysis”, preprint, 2015. To appear in *Ann. Sci. École Norm. Sup.* [arXiv 1306.4203](#)
- [Faure and Sjöstrand 2011] F. Faure and J. Sjöstrand, “Upper bound on the density of Ruelle resonances for Anosov flows”, *Comm. Math. Phys.* **308**:2 (2011), 325–364. [MR 2851145](#) [Zbl 1260.37016](#)
- [Faure and Tsujii 2013a] F. Faure and M. Tsujii, “Prequantum transfer operator for symplectic Anosov diffeomorphism”, preprint, 2013. To appear in the Astérisque series. [arXiv 1206.0282](#)
- [Faure and Tsujii 2013b] F. Faure and M. Tsujii, “Band structure of the Ruelle spectrum of contact Anosov flows”, *C. R. Math. Acad. Sci. Paris* **351**:9–10 (2013), 385–391. [MR 3072166](#) [Zbl 06192619](#)
- [Faure and Tsujii 2014] F. Faure and M. Tsujii, “The semiclassical zeta function for geodesic flows on negatively curved manifolds”, preprint, 2014. [arXiv 1311.4932](#)
- [Flaminio and Forni 2003] L. Flaminio and G. Forni, “Invariant distributions and time averages for horocycle flows”, *Duke Math. J.* **119**:3 (2003), 465–526. [MR 2004g:37039](#) [Zbl 1044.37017](#)

- [Fried 1986] D. Fried, “Analytic torsion and closed geodesics on hyperbolic manifolds”, *Invent. Math.* **84**:3 (1986), 523–540. [MR 87g:58118](#) [Zbl 0621.53035](#)
- [Fried 1995] D. Fried, “Meromorphic zeta functions for analytic flows”, *Comm. Math. Phys.* **174** (1995), 161–190. [MR 97a:58148](#) [Zbl 0841.58053](#)
- [Gel’fand and Fomin 1952] I. M. Gel’fand and S. V. Fomin, “Geodesic flows on manifolds of constant negative curvature”, *Uspehi Matem. Nauk (N.S.)* **7**:1(47) (1952), 118–137. In Russian; translated in *Amer. Math. Soc. Transl. (2)* **1** (1955), 49–65. [MR 14,660f](#) [Zbl 0048.09203](#)
- [Giulietti et al. 2013] P. Giulietti, C. Liverani, and M. Pollicott, “Anosov flows and dynamical zeta functions”, *Ann. of Math. (2)* **178**:2 (2013), 687–773. [MR 3071508](#) [Zbl 06203676](#)
- [Grellier and Otal 2005] S. Grellier and J.-P. Otal, “Bounded eigenfunctions in the real hyperbolic space”, *Int. Math. Res. Not.* **2005**:62 (2005), 3867–3897. [MR 2007a:58035](#) [Zbl 1139.58024](#)
- [Guillarmou et al. 2010] C. Guillarmou, S. Moroianu, and J. Park, “Eta invariant and Selberg zeta function of odd type over convex co-compact hyperbolic manifolds”, *Adv. Math.* **225**:5 (2010), 2464–2516. [MR 2011m:58038](#) [Zbl 1209.58017](#)
- [Guillemin 1977] V. Guillemin, “Lectures on spectral theory of elliptic operators”, *Duke Math. J.* **44**:3 (1977), 485–517. [MR 56 #6758](#) [Zbl 0463.58024](#)
- [Hansen et al. 2012] S. Hansen, J. Hilgert, and M. Schröder, “Patterson–Sullivan distributions in higher rank”, *Math. Z.* **272**:1–2 (2012), 607–643. [MR 2968245](#) [Zbl 1252.53062](#)
- [Helgason 1970] S. Helgason, “A duality for symmetric spaces with applications to group representations”, *Advances in Math.* **5** (1970), 1–154. [MR 41 #8587](#) [Zbl 0209.25403](#)
- [Helgason 1974] S. Helgason, “Eigenspaces of the Laplacian; integral representations and irreducibility”, *J. Functional Analysis* **17** (1974), 328–353. [MR 51 #3353](#) [Zbl 0303.43021](#)
- [Hörmander 1983] L. Hörmander, *The analysis of linear partial differential operators, I: Distribution theory and Fourier analysis*, Grundlehren der Math. Wissenschaften **256**, Springer, Berlin, 1983. [MR 85g:35002a](#) [Zbl 0521.35001](#)
- [Hörmander 1994] L. Hörmander, *The analysis of linear partial differential operators, III: Pseudo-differential operators*, Grundlehren der Math. Wissenschaften **274**, Springer, Berlin, 1994. [MR 95h:35255](#) [Zbl 0601.35001](#)
- [Juhl 2001] A. Juhl, *Cohomological theory of dynamical zeta functions*, Progress in Mathematics **194**, Birkhäuser, Basel, 2001. [MR 2001j:37046](#) [Zbl 1099.37017](#)
- [Leboeuf 2004] P. Leboeuf, “Periodic orbit spectrum in terms of Ruelle–Pollicott resonances”, *Phys. Rev. E* (3) **69**:2 (2004), 026204, 13. [MR 2005g:37120](#)
- [Liverani 2004] C. Liverani, “On contact Anosov flows”, *Ann. of Math. (2)* **159**:3 (2004), 1275–1312. [MR 2005k:37048](#) [Zbl 1067.37031](#)
- [Mazzeo 1991] R. Mazzeo, “Elliptic theory of differential edge operators, I”, *Comm. Partial Differential Equations* **16**:10 (1991), 1615–1664. [MR 93d:58152](#) [Zbl 0745.58045](#)
- [Mazzeo and Melrose 1987] R. R. Mazzeo and R. B. Melrose, “Meromorphic extension of the resolvent on complete spaces with asymptotically constant negative curvature”, *J. Funct. Anal.* **75**:2 (1987), 260–310. [MR 89c:58133](#) [Zbl 0636.58034](#)
- [Melrose 1993] R. B. Melrose, *The Atiyah–Patodi–Singer index theorem*, Research Notes in Mathematics **4**, A K Peters, Ltd., Wellesley, MA, 1993. [MR 96g:58180](#) [Zbl 0796.58050](#)
- [Minemura 1975] K. Minemura, “Eigenfunctions of the Laplacian on a real hyperbolic space”, *J. Math. Soc. Japan* **27**:1 (1975), 82–105. [MR 52 #3432](#) [Zbl 0292.35023](#)
- [Moore 1987] C. C. Moore, “Exponential decay of correlation coefficients for geodesic flows”, pp. 163–181 in *Group representations, ergodic theory, operator algebras, and mathematical physics* (Berkeley, CA, 1984), edited by C. C. Moore, Math. Sci. Res. Inst. Publ. **6**, Springer, New York, 1987. [MR 89d:58102](#) [Zbl 0625.58023](#)
- [Nonnenmacher and Zworski 2015] S. Nonnenmacher and M. Zworski, “Decay of correlations for normally hyperbolic trapping”, *Invent. Math.* **200**:2 (2015), 345–438. [MR 3338007](#)
- [Olbrich 1995] M. Olbrich, *Die Poisson-Transformation für homogene Vektorbündel*, Ph.D. thesis, Humboldt-Universität zu Berlin, 1995, <http://www.uni-math.gwdg.de/olbrich/>.

- [Ōshima and Sekiguchi 1980] T. Ōshima and J. Sekiguchi, “Eigenspaces of invariant differential operators on an affine symmetric space”, *Invent. Math.* **57**:1 (1980), 1–81. [MR 81k:43014](#) [Zbl 0434.58020](#)
- [Otal 1998] J.-P. Otal, “Sur les fonctions propres du laplacien du disque hyperbolique”, *C. R. Acad. Sci. Paris Sér. I Math.* **327**:2 (1998), 161–166. [MR 99e:35161](#) [Zbl 0919.31005](#)
- [Petersen 2006] P. Petersen, *Riemannian geometry*, 2nd ed., Graduate Texts in Mathematics **171**, Springer, New York, 2006. [MR 2007a:53001](#) [Zbl 1220.53002](#)
- [Pollicott 1986] M. Pollicott, “Meromorphic extensions of generalised zeta functions”, *Invent. Math.* **85**:1 (1986), 147–164. [MR 87k:58218](#) [Zbl 0604.58042](#)
- [Pollicott 1989] M. Pollicott, “Distributions at infinity for Riemann surfaces”, pp. 91–100 in *Dynamical systems and ergodic theory* (Warsaw, 1986), edited by K. Krzyżewski, Banach Center Publ. **23**, PWN, Warsaw, 1989. [MR 92e:58162](#) [Zbl 0696.58037](#)
- [Ratner 1987] M. Ratner, “The rate of mixing for geodesic and horocycle flows”, *Ergodic Theory Dynam. Systems* **7**:2 (1987), 267–288. [MR 88j:58103](#) [Zbl 0623.22008](#)
- [Ruelle 1986] D. Ruelle, “Resonances of chaotic dynamical systems”, *Phys. Rev. Lett.* **56**:5 (1986), 405–407. [MR 87f:58087](#)
- [Tsuji 2010] M. Tsuji, “Quasi-compactness of transfer operators for contact Anosov flows”, *Nonlinearity* **23**:7 (2010), 1495–1545. [MR 2012e:37040](#) [Zbl 1200.37025](#)
- [Tsuji 2012] M. Tsuji, “Contact Anosov flows and the Fourier–Bros–Iagolnitzer transform”, *Ergodic Theory Dynam. Systems* **32**:6 (2012), 2083–2118. [MR 2995886](#) [Zbl 1276.37020](#)
- [Zelditch 1987] S. Zelditch, “Uniform distribution of eigenfunctions on compact hyperbolic surfaces”, *Duke Math. J.* **55**:4 (1987), 919–941. [MR 89d:58129](#) [Zbl 0643.58029](#)
- [Zworski 2012] M. Zworski, *Semiclassical analysis*, Graduate Studies in Mathematics **138**, American Mathematical Society, Providence, RI, 2012. [MR 2952218](#) [Zbl 1252.58001](#)

Received 28 Oct 2014. Revised 30 Jan 2015. Accepted 6 Mar 2015.

SEMYON DYATLOV: dyatlov@math.mit.edu

Department of Mathematics, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, United States

FRÉDÉRIC FAURE: frederic.faure@ujf-grenoble.fr

Institut Fourier, Université Joseph Fourier, 100, rue des Maths, BP74, 38402 St Martin d’Hères, France

COLIN GUILLARMOU: cguillar@dma.ens.fr

DMA, U.M.R. 8553 CNRS, École Normale Supérieure, 45 rue d’Ulm, DMA, 75230 Paris, France

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2015 is US \$205/year for the electronic version, and \$390/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 8 No. 4 2015

Inequality for Burkholder's martingale transform PAATA IVANISVILI	765
Classification of blowup limits for SU(3) singular Toda systems CHANG-SHOU LIN, JUN-CHENG WEI and LEI ZHANG	807
Ricci flow on surfaces with conic singularities RAFE MAZZEO, YANIR A. RUBINSTEIN and NATASA SESUM	839
Growth of Sobolev norms for the quintic NLS on T^2 EMANUELE HAUS and MICHELA PROCESI	883
Power spectrum of the geodesic flow on hyperbolic manifolds SEMYON DYATLOV, FRÉDÉRIC FAURE and COLIN GUILLARMOU	923
Paving over arbitrary MASAs in von Neumann algebras SORIN POPA and STEFAAN VAES	1001