

# Asymptotic spectral gap and Weyl law for Ruelle resonances of open partially expanding maps

Jean Francois Arnoldi<sup>\*</sup>, Frédéric Faure<sup>†</sup>, Tobias Weich<sup>‡</sup>

v1:04-06-2013, v2:27-03-2015

## Abstract

We consider a simple model of an open partially expanding map. Its trapped set  $\mathcal{K}$  in phase space is a fractal set. We first show that there is a well defined discrete spectrum of Ruelle resonances which describes the asymptotic of correlation functions for large time and which is parametrized by the Fourier component  $\nu$  on the neutral direction of the dynamics. We introduce a specific hypothesis on the dynamics that we call “minimal captivity”. This hypothesis is stable under perturbations and means that the dynamics is univalued on a neighborhood of  $\mathcal{K}$ . Under this hypothesis we show the existence of an asymptotic spectral gap and a fractal Weyl law for the upper bound of density of Ruelle resonances in the semiclassical limit  $\nu \rightarrow \infty$ . Some numerical computations with the truncated Gauss map and Bowen-Series maps illustrate these results.

1

---

<sup>\*</sup>Institut Fourier, UMR 5582, 100 rue des Maths, BP74 38402 St Martin d’Hères.

<sup>†</sup>Institut Fourier, UMR 5582, 100 rue des Maths, BP74 38402 St Martin d’Hères. frederic.faure@ujf-grenoble.fr <http://www-fourier.ujf-grenoble.fr/~faure>

<sup>‡</sup>Fachbereich Mathematik, Philipps-Universität Marburg, a Hans-Meerwein-Straße, 35032 Marburg, Germany. weich@mathematik.uni-marburg.de <http://www.mathematik.uni-marburg.de/~weich/>

<sup>1</sup>2010 Mathematics Subject Classification: 37D20 Uniformly hyperbolic systems (expanding, Anosov, Axiom A, etc.) 37D35 Thermodynamic formalism, variational principles, equilibrium states 37C30 Zeta functions, (Ruelle-Frobenius) transfer operators, and other functional analytic techniques in dynamical systems 81Q20 Semiclassical techniques, including WKB and Maslov methods 81Q50 Quantum chaos

Keywords: Transfer operator; Ruelle resonances; decay of correlations; Semi-classical analysis.

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
<b>2</b>	<b>Basic definitions and statement of the main results</b>	<b>4</b>
2.1	Iterated function scheme (I.F.S.) . . . . .	4
2.2	Model of dynamics and transfer operators . . . . .	7
2.3	Discrete spectrum . . . . .	9
2.4	Asymptotic spectral radius . . . . .	10
2.5	Expansion of correlations for partially expanding maps . . . . .	13
2.6	Upper bound for the density of resonances (fractal Weyl law) . . . . .	15
<b>3</b>	<b>Proof of Theorem 2.6 about the discrete spectrum</b>	<b>17</b>
3.1	Extension of the transfer operator to distributions on $\mathbb{R}$ . . . . .	18
3.2	Dynamics on the cotangent space $T^*I$ . . . . .	20
3.3	The escape function . . . . .	21
3.4	Use of the Egorov Theorem . . . . .	22
<b>4</b>	<b>Dynamics of the canonical map <math>F : T^*I \rightarrow T^*I</math></b>	<b>24</b>
4.1	The trapped set $\mathcal{K}$ in phase space . . . . .	26
4.2	Symbolic dynamics . . . . .	29
4.3	Relation to the non-local integrability condition of Dolgopyat . . . . .	34
4.4	Dimension of the trapped set $\mathcal{K}$ . . . . .	35
<b>5</b>	<b>Proof of Theorem 2.9 for the spectral gap in the semiclassical limit</b>	<b>37</b>
5.1	The escape function . . . . .	38
5.2	Using the Egorov Theorem . . . . .	38
<b>6</b>	<b>Proof of Theorem 2.16 about the fractal Weyl law</b>	<b>41</b>
6.1	A refined escape function . . . . .	41
6.2	Weyl law . . . . .	45
<b>7</b>	<b>Numerical results for the truncated Gauss map and Bowen-Series maps</b>	<b>47</b>
7.1	The truncated Gauss map . . . . .	48
7.2	Bowen-Series maps for Schottky surfaces . . . . .	51
7.3	Proof of minimal captivity for both models . . . . .	55
<b>A</b>	<b>General lemmas on singular values of compact operators</b>	<b>57</b>
<b>B</b>	<b>Symbol classes of local <math>\hbar</math>-order</b>	<b>59</b>
B.1	Standard semiclassical symbol classes and their quantization . . . . .	59
B.2	Definition of the symbol classes $S_\mu(A_\hbar)$ . . . . .	60
B.3	Composition of symbols . . . . .	63
B.4	Ellipticity and inverses . . . . .	63

B.5 Egorov’s theorem for diffeomorphisms . . . . .	65
--	----

<b>C Adapted Weyl type estimates</b>	<b>69</b>
--------------------------------------	-----------

## 1 Introduction

The study of “Ruelle resonances” has been initiated in the seventies by D. Ruelle, R. Bowen in order to study the decay of correlations in dynamical systems. In a modern approach these Ruelle resonances show up as the discrete spectrum of transfer operators in suitable Banach spaces. While for analytic expanding maps, such function spaces were already known in the early works of Ruelle [48] for hyperbolic systems they were constructed much later by the work of Kitaev, Blank-Keller-Liverani, Baladi-Tsujii and Gouëzel-Liverani [34, 6, 28, 3, 4]. In a series of papers by the second author together with N. Roy and J. Sjöstrand it has been shown, that semiclassical techniques provide a natural approach for the construction of such suitable function spaces. Up to now this semiclassical approach to the transfer operators has been established for expanding [22] and partially expanding maps [21], Anosov diffeomorphisms [23], and Anosov flows [24]. All these systems have in common that they are closed dynamical systems, i.e. systems where the non-wandering set equals the full manifold.

The purpose of this work is to establish the semiclassical approach to “iterated function schemes” (I.F.S.) [19, 33]. In these dynamical systems the non-wandering set consists of a fractal subset of the whole system and they can thus be considered as a simple model of an open dynamical system with trapped set (i.e. non closed). Beside being a toy-model for such an open system they also appear naturally in various contexts, for example in the reduction of the geodesic flow on convex co-compact hyperbolic surfaces via Bowen-Series maps [7, 33] or in complex dynamics in the analysis of Julia sets [33]. We will study the spectral behavior of a certain family of transfer operators that are associated to these I.F.S. and using semiclassical techniques we are able to prove the existence of discrete spectrum in Sobolev spaces as well as a spectral gap and a fractal Weyl law in a certain semiclassical limit. The concrete form of the transfer operators which we study, as well as the semiclassical limit which we consider is again motivated from two sides. First of all, these families of transfer operators naturally arise from a decomposition of an open partially expanding map, which has a neutral direction. The existence of discrete spectrum together with the result on the spectral gap enables us to prove exponential decay of correlations for these systems. Secondly these transfer operators appear in the dynamical approach for Selberg zeta functions on convex co-compact surfaces and a famous result of Patterson and Perry connects the spectrum of these

transfer operators to the resonances of the Laplace operator on these surfaces. The article is organized as follows. In Section 2 we will introduce some basic definitions, state the main theorems and discuss their relation to previously known results in the literature. We also show how these transfer operators arise from open partially expanding maps and we obtain a result on the decay of correlations in such systems. Section 3 is dedicated to the semiclassical construction of the Sobolev spaces as well as to the proof of the existence of the discrete spectrum in these spaces. In Section 4 we provide a detailed study of the dynamics on the cotangent space that appears in our semiclassical approach and we are led to an important assumption on this dynamics which we call minimally captivity. In particular in Section 4.3 we show that this “minimally captive assumption” implies the “non local integrability assumption” of Dolgopyat [16] and F. Naud [40]. Section 5 and 6 are then dedicated to the proof of the spectral gap estimate and the fractal Weyl law. Finally in Section 7 we provide two important examples, show that they fulfill the minimally captive assumption and compare numerical results with the predictions of our theorems.

*Acknowledgments:* We would like to thank Stéphane Nonnenmacher and Anke Pohl for helpful discussions. This work has been supported by the “Agence National de la Recherche” via the project 2009-12 METHCHAOS. T.W. acknowledged financial support of the German National Academic foundation.

## 2 Basic definitions and statement of the main results

### 2.1 Iterated function scheme (I.F.S.)

The transfer operator studied in this paper is constructed from a simple model of chaotic dynamics called “an iterated function scheme, I.F.S.” [20, chap.9]. We give the definition below and refer to Section 7 where several standard examples are presented.

**Definition 2.1.** “An iterated function scheme (I.F.S.)”. Let  $N \in \mathbb{N}$ ,  $N \geq 1$ . Let  $I_1, \dots, I_N \subset \mathbb{R}$  be a finite collection of **disjoint bounded and closed** intervals. Let  $A$  be a  $N \times N$  matrix, called adjacency matrix, with  $A_{i,j} \in \{0, 1\}$ . We will note  $i \rightsquigarrow j$  if  $A_{i,j} = 1$ . Assume that for each pair  $i, j \in \{1, \dots, N\}$  such that  $i \rightsquigarrow j$ , we have a smooth invertible map  $\phi_{i,j} : I_i \rightarrow \phi_{i,j}(I_i) \subset \text{Int}(I_j)$ . Assume that the map  $\phi_{i,j}$  is a **strict contraction**, i.e. there exists  $0 < \theta < 1$  such that for every  $x \in I_i$ ,

$$|\phi'_{i,j}(x)| \leq \theta \quad (2.1)$$

We suppose that different images of the maps  $\phi_{i,j}$  do not intersect (this is the “strong separation condition” in [19, p.35]):

$$\phi_{i,j}(I_i) \cap \phi_{k,l}(I_k) \neq \emptyset \quad \Rightarrow \quad i = k \text{ and } j = l. \quad (2.2)$$

Note that in general the derivatives  $\phi'_{i,j}(x)$  may be negative. Notice also that we assume smoothness of the maps for our results (see Remark 2.7).

As a first illustration we will give the following example of a truncated Gauss map. Further examples will be given in Section 7.

**Example 2.2.** The Gauss map is

$$G : \begin{cases} ]0, 1[ & \rightarrow ]0, 1[ \\ y & \rightarrow \left\{ \frac{1}{y} \right\} \end{cases} \quad (2.3)$$

where  $\{a\} := a - [a] \in [0, 1[$  denotes the fractional part of  $a \in \mathbb{R}$ . Let  $j \in \mathbb{N} \setminus \{0\}$ , and  $y \in \mathbb{R}$  such that  $\frac{1}{j+1} < y \leq \frac{1}{j}$  then  $G(y) = G_j(y) := \frac{1}{y} - j$ . Notice that  $dG/dy < 0$ . The inverse map is  $y = G_j^{-1}(x) = \frac{1}{x+j}$ .

Let  $N \geq 1$ . We will consider only the first  $N$  “branches”  $(G_j)_{j=1, \dots, N}$ . In order to have a well defined I.F.S according to Definition 2.1, for  $1 \leq i \leq N$ , let  $\alpha_i := G_i^{-1}\left(\frac{1}{N+1}\right)$ ,  $a_i = \frac{1}{1+i}$ ,  $b_i$  such that  $\alpha_i < b_i < \frac{1}{i}$ , and intervals  $I_i := [a_i, b_i]$ . On these intervals  $(I_i)_{i=1, \dots, N}$ , we define the maps

$$\phi_{i,j}(x) = G_j^{-1}(x) = \frac{1}{x+j}, \quad j = 1, \dots, N. \quad (2.4)$$

See Figure 2.1. The adjacency matrix is  $A = (A_{i,j})_{i,j}$ , the full  $N \times N$  matrix with all entries  $A_{i,j} = 1$ . The values  $a_i, b_i$  are somewhat arbitrary but satisfy hypothesis (2.2). We will show below that the spectral results are independent on the intervals  $I_i = [a_i, b_i]$  and depend only on the set of branches, here  $\{1, \dots, N\}$ , as soon as the intervals  $I_i$  are large enough to contain the trapped set  $K$  defined below. See Remark 2.7(3). So we call this model the “**truncated Gauss map with  $N$  branches**”.




Figure 2.1: The iterated functions scheme (IFS) defined from the truncated Gauss map (2.3). Here we have  $N = 3$  branches. The maps  $\phi: \phi_{i,j} : I_i \rightarrow I_j$ ,  $i, j = 1 \dots N$  are contracting and given by  $\phi_{i,j}(x) = \frac{1}{x+j}$ . The trapped set  $K$  defined in (2.7) is a  $N$ -adic Cantor set. It is obtained as the limit of the sets  $K_0 = (I_1 \cup I_2 \dots \cup I_N) \supset K_1 = \phi(K_0) \supset K_2 = \phi(K_1) \supset \dots \supset K$ .

In order to shorten the notation we write

$$I := \bigcup_{i=1}^N I_i \quad (2.5)$$

and introduce the multivalued map:

$$\phi : I \rightarrow I, \quad \phi = (\phi_{i,j})_{i,j}.$$

The map  $\phi$  can be iterated and generates a multivalued<sup>2</sup> map  $\phi^n : I \rightarrow I$  for  $n \geq 1$ . From hypothesis (2.2) the inverse map

$$\phi^{-1} : \phi(I) \rightarrow I$$

is uni-valued. If we define  $K_0 := I$  and

$$K_n := \phi^n(I) \subset I \quad (2.6)$$

for all  $n \in \mathbb{N}$  then we have  $K_{n+1} \subset K_n$  and we can define the limit set

$$K := \bigcap_{n \in \mathbb{N}} K_n \quad (2.7)$$

called the **trapped set**. On this set the map

$$\phi^{-1} : K \rightarrow K \quad (2.8)$$

is well defined and uni-valued.

<sup>2</sup>For any  $x \in I$ , we have  $\#\{\phi^n(x)\} \leq N^n$ .

## 2.2 Model of dynamics and transfer operators

From the I.F.S. defined above we first define a dynamical map  $f$  that is partially expanding and introduce the transfer operator  $\hat{\mathcal{F}}$  associated to it. We first recall the following notation: we denote by  $C_0^\infty(\mathbb{R})$  the space of smooth functions on  $\mathbb{R}$  with compact support. If  $B \subset \mathbb{R}$  is a compact set, we denote  $C_0^\infty(B) \subset C_0^\infty(\mathbb{R})$  the space of smooth functions on  $\mathbb{R}$  with compact support included in  $B$ . If not further specified, we will consider complex valued functions. If we want to specify the values we write e.g.  $C_0^\infty(B; \mathbb{R})$  for real valued functions.

### 2.2.1 Partially expanding maps and transfer operators

Let  $\phi$  be an iterated function scheme as defined in Definition 2.1. Recall that the map  $\phi^{-1} : \phi(I) \rightarrow I$  is univalued and expanding. Let  $\tau \in C^\infty(\phi(I); \mathbb{R})$  be a smooth, real valued function called **roof function**. We define the map

$$f : \begin{cases} \phi(I) \times S^1 & \rightarrow I \times S^1 \\ (x, y) & \rightarrow (\phi^{-1}(x), y + \tau(x)) \end{cases} \quad (2.9)$$

with  $S^1 := \mathbb{R}/\mathbb{Z}$ . Notice that the map  $f$  is expanding in the  $x$  variable whereas it is neutral in the  $y$  variable in the sense that  $\frac{\partial f}{\partial y} = 1$ . This is called a partially expanding map and may serve as a very simple model for the general study of partially hyperbolic dynamics [44] such as Axiom A flows. Let  $V \in C^\infty(\phi(I); \mathbb{C})$  called a **potential** function.

**Definition 2.3.** The transfer operator of the map  $f$  with potential  $V$  is

$$\hat{\mathcal{F}} : \begin{cases} C_0^\infty(I \times S^1) & \rightarrow C_0^\infty(\phi(I) \times S^1) \\ \psi(x, y) & \mapsto e^{V(x)} \psi(f(x, y)) \end{cases}. \quad (2.10)$$

Notice that  $\psi(x, y)$  can be decomposed into Fourier modes in the  $y$  direction. For  $\nu \in \mathbb{Z}$ , a Fourier mode is

$$\psi_\nu(x, y) = \varphi(x) e^{i2\pi\nu y}$$

and we have

$$\begin{aligned} \left(\hat{\mathcal{F}}\psi_\nu\right)(x, y) &= e^{V(x)} \psi_\nu(f(x, y)) = e^{V(x)} \varphi(\phi^{-1}(x)) e^{i2\pi\nu(y+\tau(x))} \\ &= \left(\hat{F}_{1/(2\pi\nu)}\varphi\right)(x) e^{i2\pi\nu y} \end{aligned}$$

with a “**reduced transfer operator**”  $\hat{F}_{1/(2\pi\nu)} : C_0^\infty(I) \rightarrow C_0^\infty(\phi(I))$  given by

$$\left(\hat{F}_{1/(2\pi\nu)}\varphi\right)(x) := e^{V(x)} e^{i2\pi\nu\tau(x)} \varphi(\phi^{-1}(x)). \quad (2.11)$$




Figure 2.2: Action of the transfer operator  $\hat{F}_h$  on a function  $\varphi$  as defined in (2.12). In this schematic figure we have  $V = 0$  and  $\tau = 0$ . In general the factor  $e^{V(x)}$  changes the amplitude and  $e^{i\frac{1}{h}\tau(x)}$  creates some fast oscillations if  $\hbar \ll 1$ .

So the operator  $\hat{\mathcal{F}}$  is the direct sum of operators  $\bigoplus_{\nu \in \mathbb{Z}} \hat{F}_{1/(2\pi\nu)}$ . In this paper we are interested by the spectral properties of the operators  $\hat{F}_{1/(2\pi\nu)}$  in the limit of high Fourier modes  $\nu \rightarrow \infty$  which corresponds to strong oscillations in the neutral direction  $y$ .

### 2.2.2 Reduced transfer operators

Let us consider a direct definition for these reduced transfer operators like (2.11) which does not restrict  $\nu$  to integers.

**Definition 2.4.** Let  $\tau \in C^\infty(\phi(I); \mathbb{R})$  and  $V \in C^\infty(\phi(I); \mathbb{C})$  be smooth functions called **roof function** and **potential function**, respectively. Let  $\hbar > 0$ . We define the **transfer operator**:

$$\hat{F}_h : \begin{cases} C_0^\infty(I) & \rightarrow C_0^\infty(\phi(I)) \\ \varphi & \mapsto \begin{cases} e^{V(x)} e^{i\frac{1}{h}\tau(x)} \varphi(\phi^{-1}(x)) & \text{if } x \in \phi(I) \\ 0 & \text{otherwise} \end{cases} \end{cases} \quad (2.12)$$

See figure 2.2.

*Remark 2.5.*

- To be more precise, we consider (2.12) as a family of transfer operators depending on the parameter  $\hbar > 0$ . We will be interested in the spectrum of these operators in the “semiclassical limit”  $\hbar \rightarrow 0$ .



- For any  $\varphi \in C_0^\infty(I)$ ,  $n \geq 0$  we have

$$\text{supp} \left( \hat{F}_h^n \varphi \right) \subset K_n \quad (2.13)$$

with  $K_n$  defined in (2.6).

- In the definition (2.12) we can write  $e^{V(x)} e^{i\frac{1}{\hbar}\tau(x)} = \exp \left( i\frac{1}{\hbar}\mathcal{V}(x) \right)$  with  $\mathcal{V}(x) := \tau(x) + \hbar(-iV(x))$ . More generally we may consider a finite series  $\mathcal{V}(x) = \sum_{j=0}^n \hbar^j \mathcal{V}_j(x)$  with leading term  $\mathcal{V}_0(x) = \tau(x)$  and complex valued sub-leading terms  $\mathcal{V}_j : I \rightarrow \mathbb{C}$ ,  $j \geq 1$ .

### 2.3 Discrete spectrum

The transfer operator  $\hat{F}_h$  has been defined on smooth functions  $C_0^\infty(I)$  in (2.12). For the proof of the discrete spectrum we will need to extend it to the space of distributions (in Section 3.1) and it will turn out for technical reasons that we need to compose  $\hat{F}_h$  with a cutoff function. We thus introduce a cut-off function  $\chi \in C_0^\infty(I)$  such that  $\chi(x) = 1$  for every  $x \in K_1 = \phi(I)$ , i.e.  $\chi(\phi_{i,j}(x)) = 1$  for every  $x \in I_i$  and  $j$  such that  $i \rightsquigarrow j$ . We denote  $\hat{\chi}$  the multiplication operator by the function  $\chi$  and define:

$$\hat{F}_{h,\chi} := \hat{F}_h \circ \hat{\chi}. \quad (2.14)$$

The first main Theorem 2.6 below states that the transfer operator  $\hat{F}_{h,\chi}$  (for any  $\hbar$ ) has discrete spectrum called ‘‘Ruelle resonances’’ in ordinary Sobolev spaces with negative order and that the spectrum does not depend on the choice of  $\chi$ . Recall that for  $m \in \mathbb{R}$ , the **Sobolev space**  $H^{-m}(\mathbb{R}) \subset \mathcal{D}'(\mathbb{R})$  is defined by ([52] p.271)

$$H^{-m}(\mathbb{R}) := \left\langle \hat{\xi} \right\rangle^m (L^2(\mathbb{R})) \quad (2.15)$$

with the differential operator  $\hat{\xi} := -i\frac{d}{dx}$  and the notation  $\langle x \rangle := (1+x^2)^{1/2}$ . We also recall that a compact operator  $\hat{K}$  has discrete spectrum on  $\mathbb{C} \setminus \{0\}$  (i.e. isolated generalized eigenvalues with finite multiplicities) and if  $\hat{R}$  is an operator with norm  $\|\hat{R}\| \leq \epsilon$  then  $(\hat{K} + \hat{R})$  has still discrete spectrum on the domain  $|z| > \epsilon$ , because the essential spectrum is invariant under compact perturbations.

**Theorem 2.6. "Discrete spectrum of resonances".** For any fixed  $\hbar$ , any  $m \in \mathbb{R}$ , the transfer operator  $\hat{F}_{\hbar,\chi}$  in (2.14) can be extended to a bounded operator on the Sobolev space  $H^{-m}(\mathbb{R})$  and can be written as

$$\hat{F}_{\hbar,\chi} = \hat{K} + \hat{R} \quad (2.16)$$

where  $\hat{K}$  is a compact operator and  $\hat{R}$  is such that:

$$\left\| \hat{R} \right\|_{H^{-m}(\mathbb{R})} \leq r_m, \quad \text{with } r_m := c(\theta + \epsilon)^m \quad (2.17)$$

where  $0 < \theta < 1$  is given in (2.1), with any  $\epsilon > 0$  (taken so that  $\theta + \epsilon < 1$ ) and  $c$  that does not depend on  $m$ . This implies that the operator  $\hat{F}_{\hbar,\chi}$  has discrete spectrum on the domain  $|z| > r_m$  and that  $r_m \rightarrow 0$  as  $m \rightarrow +\infty$ . These eigenvalues of  $\hat{F}_{\hbar,\chi}$  and their eigenspace do not depend on  $m$  nor on  $\chi$ . The support of the eigendistributions is contained in the trapped set  $K$ . These discrete eigenvalues are denoted

$$\text{Res}(\hat{F}_{\hbar}) := \{\lambda_i^{\hbar}\}_i \subset \mathbb{C}^* \quad (2.18)$$

and are called **Ruelle resonances**. (See figure 7.2).

*Remark 2.7.*

1. In this paper we assume for simplicity that the maps  $\phi_{i,j}$  are  $C^\infty$ . This assumption allows to consider the limit  $m \rightarrow \infty$  in Theorem 2.6. It may be possible to assume weaker regularity, say  $C^k$ . Then Theorem 2.6 would be valid only for  $m \leq k - 1$ .
2. In the case of an I.F.S. with analytic branches and with analytic potential and roof function, it has even been shown that these transfer operators are trace class in Banach spaces of analytic functions[33, 48]. However we will proof this result with completely different techniques (microlocal or semiclassical analysis) by the construction of an escape function in the cotangent bundle  $T^*I$ . In Section 7 we will show on different examples that these techniques are also useful for concrete numerical calculations of the spectrum for Ruelle resonances.
3. The Independence of the spectrum on  $\chi$  implies that the spectral properties of the truncated Gauss map in Example 2.2 do not depend on the explicit choice of boundary points  $[a_i, b_i]$ .

## 2.4 Asymptotic spectral radius

Next we want to state a result on an asymptotic bound for the spectral radius  $r_s(\hat{F}_{\hbar,\chi})$  of the operators  $\hat{F}_{\hbar,\chi}$  in the limit  $\hbar \rightarrow \infty$ . A well known general bound on the spectral radius

of transfer operators is given in terms of the topological pressure that we recall now [49]. The topological pressure can be defined from the periodic points which are points  $x \in K$  such that  $x = \phi^{-n}(x)$ , as follows.

**Definition 2.8.** [19, p.72] The **topological pressure** of a continuous function  $\varphi \in C(I)$  is

$$\Pr(\varphi) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left( \sum_{x=\phi^{-n}(x)} e^{\varphi_n(x)} \right) \quad (2.19)$$

where  $\varphi_n(x)$  is the Birkhoff sum of  $\varphi$  along the periodic orbit:

$$\varphi_n(x) := \sum_{k=0}^{n-1} \varphi(\phi^{-k}(x)).$$

It is particularly interesting to consider the topological pressure of the “**unstable Jacobian function**” which is defined by

$$J(x) := \log \left| \frac{d\phi^{-1}}{dx}(x) \right| \quad (2.20)$$

for  $x \in \phi(I)$ . From (2.1) we obtain

$$\forall x, \quad J(x) \geq \log \frac{1}{\theta} > 0. \quad (2.21)$$

One has the general bound for every  $\hbar > 0$ , [49],

$$\log \left( r_s \left( \hat{F}_{\hbar, \chi} \right) \right) \leq \gamma_{max} := \Pr(\operatorname{Re}(V) - J). \quad (2.22)$$

In order to give the asymptotic bound (for  $\hbar \rightarrow 0$ ) below let us first introduce the so called “**damping function**”  $D \in C^\infty(\phi(I))$

$$D := \operatorname{Re}(V) - \frac{1}{2}J. \quad (2.23)$$

(This function appears naturally in different models [26, 25]). We can then give the following result on the spectral radius of the operator  $\hat{F}_{\hbar, \chi}$  in the limit  $\hbar \rightarrow \infty$ .

**Theorem 2.9. Asymptotic spectral gap.** *If the roof function  $\tau$  is “minimal captive” (see Assumption 4.7 for a precise definition) and if  $m$  is sufficiently large so that  $r_m$  (2.17) fulfills  $\log(r_m) < \gamma_+$  then the spectral radius  $r_s(\hat{F}_{\hbar,\chi})$  of the operators  $\hat{F}_{\hbar,\chi} : H^{-m}(\mathbb{R}) \rightarrow H^{-m}(\mathbb{R})$  satisfies in the semi-classical limit  $\hbar \rightarrow 0$ :*

$$\gamma_{asympt} := \log \left( \limsup_{\hbar \rightarrow 0} \left( r_s \left( \hat{F}_{\hbar,\chi} \right) \right) \right) \leq \gamma_+ \quad (2.24)$$

with

$$\gamma_+ := \lim_{n \rightarrow \infty} \left( \sup_{x \in \phi^n(I)} \frac{1}{n} \sum_{k=1}^n D(\phi^{-k}(x)) \right), \quad (2.25)$$

that corresponds to the worst Birkhoff average of the damping function  $D$  along the dynamics of the IFS. Moreover the norm of the resolvent is controlled uniformly with respect to  $\hbar$ : for any  $\rho > e^{\gamma_+}$ , there exist  $C_\rho > 0$ ,  $\hbar_\rho > 0$  such that  $\forall \hbar < \hbar_\rho$ ,  $\forall |z| > \rho$  we have

$$\left\| \left( z - \hat{F}_{\hbar} \right)^{-1} \right\|_{H^{-m}(\mathbb{R})} \leq C_\rho. \quad (2.26)$$

*Remark 2.10.*

1. The limit on the right hand side of (2.25) exists: the sequence  $a_n := \sup_{x \in \phi^n(I)} \sum_{k=1}^n D(\phi^{-k}(x))$  is subadditive (i.e.  $a_n + a_m \geq a_{n+m}$ ) and Fekete’s Lemma guaranties existence of the limit  $\gamma_+ = \lim_{n \rightarrow \infty} a_n/n$ .
2. From the general bound (2.22) valid every  $\hbar$  we have  $\gamma_{asympt} \leq \gamma_{max} := \Pr(\text{Re}(V) - J)$  and we may define precisely  $g_{asympt} := \gamma_{max} - \gamma_{asympt} \geq 0$  to be the **asymptotic spectral gap**. In many cases (but not always), see the concrete examples in Section 7, we have  $\gamma^+ < \gamma_{max}$ . In particular for closed systems treated in [21] and for  $V = 0$ , one has always  $\gamma_{max} = 0$  and  $D = -\frac{1}{2}J < 0$  hence  $\gamma_+ < \gamma_{max}$ .
3. F. Naud obtained in [40] (using techniques of D. Dolgopyat), an asymptotic bound on the spectral radius under a so called “non-local integrability” condition weaker than the “minimal captive assumption” and which is discussed below. Translated to our setting he showed the existence of  $\epsilon > 0$  such that  $\gamma_{asympt} \leq \gamma_{max} - \epsilon < \gamma_{max}$ , i.e. that  $g_{asympt} > 0$ . For systems where  $\gamma^+ < \gamma_{max}$  the result (2.24) improves this bound as it gives an explicit estimate  $g_{asympt} > \gamma_{max} - \gamma_+ > 0$ . However for a general system one may have  $\gamma^+ > \gamma_{max}$  and the result (2.24) gives no asymptotic spectral gap whereas F. Naud’s result always gives one.
4. Notice that Theorem 2.9 depends on the roof function  $\tau$  only implicitly through Assumption 4.7. The value of the upper bound (2.25) does not depend on  $\tau$ . It

is however known that such results cannot hold for a general roof function  $\tau$  (for example it does not hold for roof functions that are cohomologous to a constant, see [21, Appendix A]).

5. Dolgopyat [16, 17] and Naud [40] used the so called “non-local integrability” condition (NLI) and we will see in Section 4.3 that our minimally captive assumption implies this non-local integrability condition. The “minimally captivity” condition which we use arises naturally in the semiclassical approach used in the proof (see Section 4 for a detailed introduction and definition). It is a similar, but stronger assumption than the condition which appeared in [54, 21] and which was coined “partially captive” in the latter reference. With only moderate effort Theorem 2.9 could be proven also under the weaker assumption of “partial captivity” but it will turn out that minimally captivity makes the phase space dynamics on  $T^*I$  particularly easy and is essential in the proof of the fractal Weyl law. This is why we decided to put this condition in the center of attention in this article.
6. In Section 7 we will illustrate with numerical results on the example of the truncated Gauss map, that the bound (2.24) does not seem to be optimal. Also other related numerical and physical experiments [5] have supported the conjecture, that the rigorously known spectral gap estimates are not sharp. The question of finding sharp estimates of asymptotic spectral gaps is an important open question (see e.g. [41] for an overview and further references).

## 2.5 Expansion of correlations for partially expanding maps

In this section we present a quite immediate consequence of the existence of an asymptotic spectral radius  $e^{\gamma+}$  obtained in Theorem 2.9: we obtain a finite expansion for correlation functions  $\langle v | \hat{\mathcal{F}}^n u \rangle$  of the extended transfer operator  $\hat{\mathcal{F}}$  defined in (2.10).

We first introduce some notation: for a given  $\nu \in \mathbb{Z}$ , we have seen in Theorem 2.6 that the transfer operator  $\hat{F}_{1/(2\pi\nu), \chi}$  has a discrete spectrum of resonances. For  $\rho > 0$  such that there is no eigenvalue on the circle  $|z| = \rho$  and for any  $\nu \in \mathbb{Z}$ , we denote by  $\Pi_{\rho, \nu}$  the spectral projector of the operator  $\hat{F}_{1/(2\pi\nu), \chi}$  on the domain  $\{z \in \mathbb{C}, |z| > \rho\}$ . These projection operators have obviously finite rank and each commutes with  $\hat{F}_{1/(2\pi\nu), \chi}$ .

**Theorem 2.11. "Expansion of correlations".** For any  $\rho > e^{\gamma+}$ ,  $m$  large enough (such that  $r_m < e^{\gamma+}$  in Th. 2.6), there exists  $\nu_0 \in \mathbb{N}$  and  $C_\rho > 0$  such that for any  $u \in H^{-m}(I) \otimes L^2(S^1)$ ,  $v \in H^m(I) \otimes L^2(S^1)$ , in the limit  $n \rightarrow \infty$ ,

$$\left| \langle v | \hat{\mathcal{F}}^n u \rangle - \sum_{|\nu| \leq \nu_0} \langle v_\nu | \left( \hat{F}_{1/(2\pi\nu), \chi} \Pi_{\rho, \nu} \right)^n u_\nu \rangle \right| \leq C_\rho \rho^n \|u\|_{H^{-m} \otimes L^2(S^1)}^2 \|v\|_{H^m \otimes L^2(S^1)}^2 \quad (2.27)$$

Here  $u_\nu \in H^{-m}(I)$ ,  $v_\nu \in H^m(I)$  stand for the Fourier components in  $S^1$  direction of  $u, v$  and  $\langle v | u \rangle = \int_{I \times S^1} \bar{v}(x) u(x) dx$  (extended to distributions).

*Remark 2.12.*

- The second term in Eq.(2.27) is a finite sum and each operator  $\hat{F}_{1/(2\pi\nu), \chi} \Pi_{\rho, \nu}$  has finite rank. Using the spectral decomposition of  $\hat{F}_\nu$  we get an expansion of the correlation function  $\langle v | \hat{\mathcal{F}}^n u \rangle$  with a finite number of terms which involve the leading Ruelle resonances (i.e. those with modulus greater than  $\rho$ ) and an error term that is  $O(\rho^n)$ .
- The novelty of this Theorem is that the correlations can be expanded up to this error term  $O(\rho^n)$  for any  $\rho > e^{\gamma+}$ . As discussed in Remark 2.10, previous results are restricted to error terms  $(e^{\text{Pr}(\text{Re}(V)-J)-\varepsilon})^n$  with some non explicit  $\varepsilon > 0$  and more restrictive function spaces for  $u, v$ .

*Proof.* Let  $\rho > e^{\gamma+}$ . Recall that  $\hbar = \frac{1}{2\pi\nu}$  and that  $|\nu| \rightarrow \infty$  corresponds to  $\hbar \rightarrow 0$ . In Theorem 2.6 we have for  $\hbar \rightarrow 0$  that  $r_s \left( \hat{F}_{1/(2\pi\nu), \chi} \right) \leq e^{\gamma+} + o(1)$ . Let the value of  $\nu_0$  be such that  $r_s \left( \hat{F}_{1/(2\pi\nu), \chi} \right) < \rho$  for every  $|\nu| > \nu_0$ . Then for any  $u \in H^{-m}(I) \otimes L^2(S^1)$ ,  $v \in H^m(I) \otimes L^2(S^1)$

$$\langle v | \hat{\mathcal{F}}^n u \rangle = \sum_{|\nu| \leq \nu_0} \langle v_\nu | \left( \hat{F}_{1/(2\pi\nu), \chi} \Pi_{\rho, \nu} \right)^n u_\nu \rangle + \sum_{|\nu| \leq \nu_0} \langle v_\nu | \left( \hat{F}_{1/(2\pi\nu), \chi} (\text{Id} - \Pi_{\rho, \nu}) \right)^n u_\nu \rangle \quad (2.28)$$

$$+ \sum_{|\nu| > \nu_0} \langle v_\nu | \hat{F}_{1/(2\pi\nu), \chi}^n u_\nu \rangle. \quad (2.29)$$

We have  $\left| \langle v_\nu | \hat{F}_{1/(2\pi\nu), \chi}^n u_\nu \rangle \right| \leq \|v_\nu\|_{H^m} \|u_\nu\|_{H^{-m}} \left\| \hat{F}_{1/(2\pi\nu), \chi}^n \right\|_{H^{-m}}$ . For  $|\nu| \leq \nu_0$  we have

$$\left\| \left( \hat{F}_{1/(2\pi\nu), \chi} (\text{Id} - \Pi_{\rho, \nu}) \right)^n \right\| \leq C_{\nu_0} \rho^n$$

where  $C_{\nu_0}$  depends on  $\nu_0$ , hence

$$\left| \sum_{|\nu| \leq \nu_0} \langle v_\nu | \left( \hat{F}_{1/(2\pi\nu), \chi} (\text{Id} - \Pi_{\rho, \nu}) \right)^n u_\nu \rangle \right| \leq C_{\nu_0} \rho^n \sum_{|\nu| \leq \nu_0} \|v_\nu\|_{H^m} \|u_\nu\|_{H^{-m}}.$$

On one hand, as a sequence with respect to  $\nu \in \mathbb{Z}$ , one has  $(\|u_\nu\|_{H^m})_\nu, (\|v_\nu\|_{H^{-m}})_\nu \in l^2(\mathbb{Z})$  and  $\sum_{\nu \in \mathbb{Z}} \|u_\nu\|_{H^m}^2 = \|u\|_{H^m \otimes L^2(S^1)}^2$ . Additionally the resolvent bound (2.26) gives us the existence of a constant  $C_\rho$  such that

$$\left\| \left( z - \hat{F}_{1/(2\pi\nu)} \right)^{-1} \right\|_{H^{-m}} \leq C_\rho$$

uniformly in  $|z| > \rho$  and  $|\nu| > \nu_0$ . From the Cauchy formula  $\hat{F}_{1/(2\pi\nu), \chi}^n = \frac{1}{2\pi i} \oint_\gamma z^n \left( z - \hat{F}_{1/(2\pi\nu), \chi} \right)^{-1} dz$  where  $\gamma$  is the circle of radius  $\rho$  one deduces that  $\left\| \hat{F}_{1/(2\pi\nu), \chi}^n \right\|_{H^{-m}} \leq C_\rho \rho^n$ . So

$$\left| \sum_{|\nu| > \nu_0} \langle v_\nu | \hat{F}_{1/(2\pi\nu), \chi}^n u_\nu \rangle \right| \leq C_\rho \rho^n \sum_{|\nu| > \nu_0} \|v_\nu\|_{H^m} \|u_\nu\|_{H^{-m}} \leq C_\rho \rho^n \|u\|_{H^m \otimes L^2(S^1)} \|v\|_{H^m \otimes L^2(S^1)}.$$

Then (2.28) gives (2.27). □

## 2.6 Upper bound for the density of resonances (fractal Weyl law)

We will finally formulate a fractal Weyl law on the number of Ruelle resonances and therefore introduce the following definition of fractal dimension.

**Definition 2.13.** [38, p.76],[19, p.20] If  $B \subset \mathbb{R}^d$  is a non empty bounded set, its **upper Minkowski dimension** (or box dimension) is

$$\dim_M B := d - \text{codim}_M B \tag{2.30}$$

with

$$\text{codim}_M B := \sup \left\{ s \in \mathbb{R} \mid \limsup_{\delta \downarrow 0} \delta^{-s} \cdot \text{Leb}(B_\delta) < +\infty \right\}. \tag{2.31}$$

where  $B_\delta := \{x \in \mathbb{R}^d, \text{dist}(x, B) \leq \delta\}$  and  $\text{Leb}(\cdot)$  is the Lebesgue measure.

*Remark 2.14.* In general

$$\limsup_{\delta \downarrow 0} \delta^{-\text{codim}_M B} \cdot \text{Leb}(B_\delta) < +\infty \tag{2.32}$$

does not hold, but if it does,  $B$  is said to be of **pure dimension**<sup>3</sup>. It is known that the trapped set  $K$  defined in (2.7) has pure dimension and that the above definition of Minkowski dimension coincides with the more usual **Hausdorff dimension** of  $K$  [19, p.68]:

$$\dim_M K = \dim_H K \in [0, 1[ \tag{2.33}$$

<sup>3</sup>see [51] for comments and further references.




Figure 2.3:  $\text{Pr}_N(-\beta J)$  is the Topological Pressure (2.19) for the truncated Gauss map (Example 2.2), with  $N = 1, 2, 3 \dots$  being the number of branches. The black points mark the zero of  $\text{Pr}_N(-\beta J) = 0$  giving the fractal dimension of the trapped set  $K_N$  for each value of  $N$ :  $\dim_H K_1 = 0$ ,  $\dim_H K_2 = 0.531 \dots$ ,  $\dim_H K_3 = 0.705 \dots$ , and  $\dim_H K_N \xrightarrow{N \rightarrow \infty} 1$ .

Using the following Lemma, the topological pressure defined in (2.19) provides an efficient way to calculate the Hausdorff dimension of the trapped set numerically (see Figure 2.3 for an illustration on the example of the truncated Gauss map).

**Lemma 2.15.** [19, p.77] *If  $\beta > 0$  is a real parameter and  $J$  the unstable Jacobian defined in (2.20), then  $\text{Pr}(-\beta J)$  is continuous and strictly decreasing as a function of  $\beta$  and its unique zero is given by  $\beta = \dim_H K$ .*

We can finally formulate

**Theorem 2.16. "Fractal Weyl upper bound".** *Suppose that the assumption of minimal captivity 4.7 holds and that the adjacency matrix  $A$  is symmetric. For any  $\varepsilon > 0$ , any  $\eta > 0$ , we have for  $\hbar \rightarrow 0$*

$$\#\left\{ \lambda_i^{\hbar} \in \text{Res}(\hat{F}_{\hbar}) \mid |\lambda_i^{\hbar}| \geq \varepsilon \right\} = \mathcal{O}(\hbar^{-\dim_H(K) - \eta}). \quad (2.34)$$

The first result of a fractal Weyl law upper bound has been obtained by J. Sjöstrand [51] for a wide class of semiclassical operators with analytic coefficients. This pioneering



work has triggered as well theoretical and experimental studies in physics [46, 36, 50, 35] as well as an extension of this theorem to various other settings like convex co-compact surfaces [30, 56], manifolds with hyperbolic ends [14] and the scattering at several convex bodies [42]. To our knowledge there are not yet any rigorous result of a fractal Weyl law upper bound for classical Ruelle resonances, however in physics literature the existence of such laws has been observed [18]. The minimal captivity assumption however allows to interpret the transfer operator as a quantization of a bijective map (see [26] for discussions about this interpretation) and it should also be possible to obtain this result using the recent work of Nonnenmacher, Sjöstrand and Zworski [42]. We will however provide an independent proof which is directly based on the semiclassical approach for the transfer operators, by further refining the escape function that appears in the proofs of Theorem 2.6 and 2.9.

The upper bound on the exponent in terms of the Hausdorff dimension is conjectured to be sharp [36], meaning that it is also a lower bound (see also [41] for an overview and further references). This conjecture has been supported by several numerical experiments for example for quantum  $n$ -disk systems [36] or convex co-compact surfaces [8]. Also in the case of iterated function schemes, the bound seems to be also a lower bound as suggested by the numerical results shown in Figure 7.3.

### 3 Proof of Theorem 2.6 about the discrete spectrum

For this proof we follow closely the proof<sup>4</sup> of Theorem 2 in the paper [21] which uses semiclassical analysis. However we have to deal with an additional difficulty associated with the “openness” of the system. This will be taken into account by the introduction of the cut off function  $\chi$  (c.f. (2.14)). In this Section 3 the parameter  $\hbar$  is fixed.

Here is the strategy. We first show in Section 3.1 that the transfer operator  $\hat{F}_{\hbar,\chi}$  has a well defined and unique extension to distributions on  $\mathbb{R}$ . Then in Section 3.2 we explain that the transfer operator is a Fourier integral operator and compute its associated symplectic map on the cotangent space  $\mathfrak{F} : T^*I \rightarrow T^*I$ . We observe that under this map  $\mathfrak{F}$ , the trajectory of a point  $(x, \xi) \in T^*I$  escape towards infinity if  $|\xi| > 0$ . In Section 3.3 we construct an escape function (or Lyapounov function)  $A_m(x, \xi)$  that decreases strictly along the trajectories. We consider the corresponding pseudodifferential operator  $\hat{A}_m := \text{Op}(A_m)$  and in Section 3.4 we show that the conjugated operator  $\hat{A}_m \circ \hat{F}_{\hbar,\chi} \circ \hat{A}_m^{-1}$  has discrete spectrum in  $L^2(\mathbb{R})$ . Equivalently the transfer operator  $\hat{F}_{\hbar,\chi}$  has discrete spectrum in the Sobolev space  $\hat{A}_m^{-1}(L^2(\mathbb{R})) = H^{-m}(\mathbb{R})$ .

---

<sup>4</sup>see also Theorem 4 in [23] which concerns hyperbolic maps and anisotropic Sobolev spaces.

### 3.1 Extension of the transfer operator to distributions on $\mathbb{R}$

Recall that in (2.14) we have introduced the cut off function  $\chi \in C_0^\infty(I)$  with  $\chi(x) = 1$  for every  $x \in K_1 = \phi(I)$  as well as the truncated transfer operators

$$\hat{F}_{h,\chi} := \hat{F}_h \circ \hat{\chi}$$

Note that for any  $\varphi \in C_0^\infty(K_1)$  we have  $\hat{\chi}\varphi = \varphi$  hence  $(\hat{F}_h \hat{\chi})\varphi = \hat{F}_h\varphi$ . One has  $\hat{\chi} : C_0^\infty(\mathbb{R}) \rightarrow C_0^\infty(I)$  hence  $\hat{F}_{h,\chi}$  is defined on  $C_0^\infty(\mathbb{R})$ .

The formal adjoint operator  $\hat{F}_{h,\chi}^* : C_0^\infty(\mathbb{R}) \rightarrow C_0^\infty(\mathbb{R})$  is defined by

$$\langle \varphi | \hat{F}_{h,\chi}^* \psi \rangle = \langle \hat{F}_{h,\chi} \varphi | \psi \rangle, \quad \forall \varphi \in C_0^\infty(\mathbb{R}), \psi \in C_0^\infty(\mathbb{R}), \quad (3.1)$$

with the  $L^2$ -scalar product  $\langle u | v \rangle := \int \bar{u}(x) v(x) dx$ . Note that for any test function  $\varphi \in C_0^\infty(\mathbb{R})$  with  $\text{supp}(\varphi) \cap I = \emptyset$  we have  $\hat{F}_{h,\chi}\varphi = 0$  which directly implies that  $\hat{F}_{h,\chi}^* : C_0^\infty(\mathbb{R}) \rightarrow C_0^\infty(I)$ .

**Lemma 3.1.** *Let  $\psi \in C_0^\infty(\mathbb{R})$  and  $y \in I_i$ . Then the adjoint operator  $\hat{F}_{h,\chi}^* : C_0^\infty(\mathbb{R}) \rightarrow C_0^\infty(I)$  is given by*

$$\left( \hat{F}_{h,\chi}^* \psi \right) (y) = \chi(y) \sum_{j \text{ s.t. } i \rightsquigarrow j} |\phi'_{i,j}(y)| e^{\overline{V(\phi_{i,j}(y))}} e^{-\frac{i}{\hbar} \tau(\phi_{i,j}(y))} \psi(\phi_{i,j}(y)). \quad (3.2)$$

*Proof.* Using the definition (3.2), we calculate

$$\begin{aligned} \langle \varphi | \hat{F}_{h,\chi}^* \psi \rangle &= \int_I \bar{\varphi}(y) \left( \hat{F}_{h,\chi}^* \psi \right) (y) dy \\ &= \sum_i \int_{I_i} \bar{\varphi}(y) \left( \hat{F}_{h,\chi}^* \psi \right) (y) dy \\ &= \sum_i \sum_{j \text{ s.t. } i \rightsquigarrow j} \int_{I_i} \bar{\varphi}(y) \chi(y) |\phi'_{i,j}(y)| e^{\overline{V(\phi_{i,j}(y))}} e^{-\frac{i}{\hbar} \tau(\phi_{i,j}(y))} \psi(\phi_{i,j}(y)) dy. \end{aligned}$$

Now we can perform a change of variables  $x = \phi_{i,j}(y)$  in each of the integrals and obtain

$$\begin{aligned} \langle \varphi | \hat{F}_{h,\chi}^* \psi \rangle &= \sum_i \sum_{j \text{ s.t. } i \rightsquigarrow j} \int_{\phi_{i,j}(I_i)} \overline{e^{V(x)} e^{i \frac{1}{\hbar} \tau(x)} \varphi(\phi^{-1}(x)) \chi(\phi^{-1}(x))} \psi_j(x) dx \\ &= \int_I \overline{\hat{F}_{h,\chi} \varphi(x)} \psi(x) dx \\ &= \langle \hat{F}_{h,\chi} \varphi | \psi \rangle. \end{aligned}$$

□

**Proposition 3.2.** *By duality the transfer operator (2.14) extends to distributions:*

$$\hat{F}_{h,\chi} : \mathcal{D}'(\mathbb{R}) \rightarrow \mathcal{D}'(\mathbb{R}) \quad (3.3)$$

$$\hat{F}_{h,\chi}^* : \mathcal{D}'(\mathbb{R}) \rightarrow \mathcal{D}'(\mathbb{R}).$$

Similarly to (2.13) we have that for any  $n \geq 1$ , any  $\alpha \in \mathcal{D}'(\mathbb{R})$ ,

$$\text{supp} \left( \hat{F}_{h,\chi}^n \alpha \right) \subset K_n \quad (3.4)$$

with  $K_n$  defined in (2.6).

*Proof.* As  $\hat{F}_{h,\chi}^*$  is continuous on the space of test functions  $C_0^\infty(\mathbb{R})$  the extension can directly be defined by

$$\hat{F}_{h,\chi}(\alpha)(\psi) = \alpha \left( \overline{\hat{F}_{h,\chi}^* \psi} \right), \quad \alpha \in \mathcal{D}'(\mathbb{R}), \psi \in C_0^\infty(\mathbb{R}), \quad (3.5)$$

If  $\psi(\phi_{i,j}(y)) = 0$  for all  $i \rightsquigarrow j$  and all  $y \in I_i$  then (3.2) shows that  $\hat{F}_{h,\chi}^* \psi \equiv 0$ . More generally let  $\psi \in C_0^\infty(\mathbb{R})$  with  $\text{supp}(\psi) \cap K_n = \emptyset$  with  $n \geq 1$  and  $K_n$  defined in (2.6). Then

$$\left( \hat{F}_{h,\chi}^* \right)^n \psi \equiv 0. \quad (3.6)$$

For any  $\alpha \in \mathcal{D}'(\mathbb{R})$ , we deduce that  $\left( \hat{F}_{h,\chi}^n \alpha \right)(\bar{\psi}) = \alpha \left( \overline{\left( \hat{F}_{h,\chi}^* \right)^n \psi} \right) = 0$ . By definition, this means that  $\text{supp} \left( \hat{F}_{h,\chi}^n \alpha \right) \subset K_n$ .  $\square$

*Remark 3.3.*

- Without the cut-off function  $\chi$  the image of  $\hat{F}_h^*$  may not be continuous on the boundary of  $I$  and the extension to distribution space in Proposition 3.2 would not have been possible.
- An other more general possibility would have been to consider  $\chi \in C_0^\infty(I)$  such that  $0 < \chi(x)$  for  $x \in \text{Int}(I)$  (without assumption that  $\chi \equiv 1$  on  $K_1$ ) and define

$$\hat{F}_{h,\chi} := \hat{\chi}^{-1} \hat{F}_h \hat{\chi} : C_0^\infty(\mathbb{R}) \rightarrow C_0^\infty(\mathbb{R}) \quad (3.7)$$

which is well defined since  $\text{supp} \left( \hat{F}_h \hat{\chi} \varphi \right) \subset \text{Int}(I)$  where  $\chi$  does not vanish. This more general definition (3.7) may be more useful in some cases, e.g. we use it in numerical computation. We recover the previous definition (2.14) if we make the additional assumption that  $\chi \equiv 1$  on  $K_1$ .

### 3.2 Dynamics on the cotangent space $T^*I$

The remark of fundamental importance given in Proposition 3.4 below is that each operator  $\hat{F}_{\hbar,\chi}$ , although it is a simple composition operator, can be considered as a “Fourier integral operator” whose associate canonical map  $\mathfrak{F}$  is the map  $\phi : I \rightarrow I$  lifted on the cotangent space  $T^*I$ . The definition of Fourier integral operator and its associated canonical map will be given in a more general context in the beginning of Section 4 and the following proposition can be considered as a particular case of Lemma 4.2. Then, according to the “semiclassical approach” we know that in order to study the spectral properties of the transfer operator  $\hat{F}_{\hbar,\chi}$  we have first to study the dynamics of its canonical map  $\mathfrak{F} : T^*I \rightarrow T^*I$ . It is not necessary to know what is a Fourier integral operator to read the main part of this paper.

**Proposition 3.4.** *Considering  $\hbar > 0$  fixed, the transfer operator  $\hat{F}_{\hbar,\chi}$  restricted to  $C_I^\infty(\mathbb{R})$  is a Fourier integral operator (FIO). Its **canonical transform** is a multi-valued symplectic map  $\mathfrak{F} : T^*I \rightarrow T^*I$  on the cotangent space  $T^*I \equiv I \times \mathbb{R}$  given by:*

$$\mathfrak{F} : \begin{cases} T^*I & \rightarrow T^*I \\ (x, \xi) & \mapsto \{\mathfrak{F}_{i,j}(x, \xi), \text{ with } i, j \text{ s.t. } x \in I_i, i \rightsquigarrow j\} \end{cases}$$

with

$$\mathfrak{F}_{i,j} : \begin{cases} x' & = \phi_{i,j}(x) \\ \xi' & = \frac{1}{\phi'_{i,j}(x)}\xi \end{cases}. \quad (3.8)$$

#### Remarks

- For the proof we refer to the proof of Lemma 4.2 with the following remark. Here  $\hbar$  is fixed (it is not a semiclassical parameter) hence the term  $e^{i\frac{1}{\hbar}\tau(x')}$  in (2.12) contributes to the amplitude and not to the phase function. That explains that the canonical map  $\mathfrak{F}$  differs from the canonical map  $F$  which will be introduced in (4.5).
- In [21, Section 3.2] we explain the action of the FIO  $\hat{F}_{\hbar,\chi}$  in terms of wave packets and the clear relation with the symplectic map  $\mathfrak{F}$ .
- For short, we can write

$$\mathfrak{F} : \begin{cases} T^*I & \rightarrow T^*I \\ (x, \xi) & \mapsto \left( \phi(x), \frac{1}{\phi'(x)}\xi \right) \end{cases}. \quad (3.9)$$

- Observe from (3.9) that the dynamics of the map  $\mathfrak{F}$  on  $T^*I$  has a quite simple property: the zero section  $\{(x, \xi) \in I \times \mathbb{R}, \xi = 0\}$  is globally invariant and any other

point  $(x, \xi)$  with  $\xi \neq 0$  escapes towards infinity ( $\xi \rightarrow \pm\infty$ ) in a controlled manner, because  $|\phi'_{i,j}(x)| < \theta < 1$ , with  $\theta$  given in (2.1), hence:

$$|\xi'| \geq \frac{1}{\theta} |\xi| \quad (3.10)$$

- Due to hypothesis (2.2) the map  $\phi_{i,j}^{-1}$  is uni-valued (when it is defined). Therefore the map  $\mathfrak{F}^{-1}$  is also uni-valued and one has

$$\mathfrak{F}^{-1} \circ \mathfrak{F} = \text{Id}_{T^*I}. \quad (3.11)$$

### 3.3 The escape function

**Definition 3.5.** [53, p.2] For  $m \in \mathbb{R}$ , the **class of symbols**  $S^{-m}(T^*\mathbb{R})$ , with **order**  $m$ , is the set of functions on the cotangent space  $A \in C^\infty(T^*\mathbb{R})$  such that for any  $\alpha, \beta \in \mathbb{N}$ , there exists  $C_{\alpha,\beta} > 0$  such that

$$\forall (x, \xi) \in T^*\mathbb{R}, \quad \left| \partial_x^\alpha \partial_\xi^\beta A(x, \xi) \right| \leq C_{\alpha,\beta} \langle \xi \rangle^{-m-|\beta|}, \quad \text{with } \langle \xi \rangle = (1 + \xi^2)^{1/2}. \quad (3.12)$$

**Lemma 3.6.** *Let  $m > 0$  and let*

$$A_m(x, \xi) := \langle \xi \rangle^{-m} \in S^{-m}(T^*\mathbb{R}).$$

*We have*

$$\forall R > 0, \forall |\xi| > R, \quad \forall i \rightsquigarrow j, \forall x \in I_i, \quad \frac{A_m(\mathfrak{F}_{i,j}(x, \xi))}{A_m(x, \xi)} \leq C^m, \quad (3.13)$$

*with  $C = \sqrt{\frac{R^2+1}{R^2/\theta^2+1}} < 1$ . Eq.(3.13) shows that  $A_m$  decreases strictly along the trajectories of  $\mathfrak{F}$  outside the zero section. We say that  $A_m$  is an **escape function**.*

*Proof.* From Eq. (3.8) and (3.10) we have

$$\frac{A_m(\mathfrak{F}_{i,j}(x, \xi))}{A_m(x, \xi)} = \frac{(1 + \xi^2)^{m/2}}{(1 + (\xi')^2)^{m/2}} \leq \frac{(1 + \xi^2)^{m/2}}{(1 + \xi^2/\theta^2)^{m/2}} \leq \left( \frac{1 + R^2}{1 + R^2/\theta^2} \right)^{m/2} = C^m.$$

The last inequality is because the function decreases with  $|\xi|$ .  $\square$

Using the standard quantization rule [53, p.2] the symbol  $A_m$  can be quantized into a pseudodifferential operator  $\hat{A}_m$  (PDO for short) which is self-adjoint and invertible on  $C_0^\infty(\mathbb{R})$ :

$$\left(\hat{A}_m\varphi\right)(x) = \frac{1}{2\pi} \int A_m(x, \xi) e^{i(x-y)\xi} \varphi(y) dy d\xi. \quad (3.14)$$

Conversely  $A_m$  is called the symbol of the PDO  $\hat{A}_m$ . In our simple case, this is very explicit: in Fourier space,  $\hat{A}_m$  is simply the multiplication by  $\langle \xi \rangle^m$ . Its inverse  $\hat{A}_m^{-1}$  is the multiplication by  $\langle \xi \rangle^{-m}$ .

### 3.4 Use of the Egorov Theorem

Let

$$\hat{Q}_m := \hat{A}_m \hat{F}_{h,\chi} \hat{A}_m^{-1} : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}),$$

which is unitarily equivalent to  $\hat{F}_{h,\chi} : H^{-m}(\mathbb{R}) \rightarrow H^{-m}(\mathbb{R})$  (from the definition of  $H^{-m}(\mathbb{R})$ , Eq.(2.15)). This is expressed by the following commutative diagram

$$\begin{array}{ccc} L^2(\mathbb{R}) & \xrightarrow{\hat{Q}_m} & L^2(\mathbb{R}) \\ \downarrow \hat{A}_m^{-1} & & \downarrow \hat{A}_m^{-1} \\ H^{-m}(\mathbb{R}) & \xrightarrow{\hat{F}_{h,\chi}} & H^{-m}(\mathbb{R}) \end{array}. \quad (3.15)$$

We will therefore study the operator  $\hat{Q}_m$  on  $L^2(\mathbb{R})$ . Notice that  $\hat{Q}_m$  is defined a priori on a dense domain  $C_0^\infty(\mathbb{R})$ . Define

$$\hat{P} := \hat{Q}_m^* \hat{Q}_m = \hat{A}_m^{-1} \left( \hat{F}_{h,\chi}^* \hat{A}_m^2 \hat{F}_{h,\chi} \right) \hat{A}_m^{-1} = \hat{A}_m^{-1} \hat{B} \hat{A}_m^{-1}, \quad (3.16)$$

with

$$\hat{B} := \hat{F}_{h,\chi}^* \hat{A}_m^2 \hat{F}_{h,\chi} = \hat{\chi} \hat{F}_h^* \hat{A}_m^2 \hat{F}_h \hat{\chi}. \quad (3.17)$$

Now, the crucial step in the proof is to use the Egorov Theorem.

**Lemma 3.7. (Egorov theorem).**  $\hat{B}$  defined in (3.17) is a pseudo-differential operator with symbol in  $S^{-2m}(T^*\mathbb{R})$  given by:

$$B(x, \xi) = \left( \chi^2(x) \sum_{j \text{ s.t. } i \rightsquigarrow j} |\phi'_{i,j}(x)| e^{2\text{Re}(V(\phi_{i,j}(x)))} A_m^2(\mathfrak{F}_{i,j}(x, \xi)) \right) + R \quad (3.18)$$

where  $R \in S^{-2m-1}(T^*\mathbb{R})$  has a lower order,  $x \in I_i$ ,  $\xi \in \mathbb{R}$ .

*Proof.*  $\hat{F}_h$  and  $\hat{F}_h^*$  are Fourier integral operators (FIO) whose canonical maps are respectively  $\mathfrak{F}$  and  $\mathfrak{F}^{-1}$ . The pseudodifferential operator (PDO)  $\hat{A}_m$  can also be considered as a FIO whose canonical map is the identity. By composition we deduce that  $\hat{B} = \hat{\chi}\hat{F}_h^*\hat{A}_m^2\hat{F}_h\hat{\chi}$  is a FIO whose canonical map is the identity since  $\mathfrak{F}^{-1} \circ \mathfrak{F} = I$  from (3.11). Therefore  $\hat{B}$  is a PDO. Using the precise expressions for  $\hat{F}_h$  (Eq. (2.12)) and  $\hat{F}_h^*$ , (Eq. (3.2)) as well as the behavior of PDOs under a change of variables (see [29, Theorem 3.9]), we obtain that the principal symbol of  $\hat{B}$  is the first term of (3.18).  $\square$

Remark: Contrary to (3.17),  $\hat{F}_h\hat{A}_m\hat{F}_h^*$  is not a PDO, but a FIO whose canonical map  $\mathfrak{F} \circ \mathfrak{F}^{-1}$  is multivalued.

Now by **theorem of composition of PDO** [53, p.11], Eq.(3.16) and Eq.(3.18) imply that  $\hat{P}$  is a PDO with symbol in  $S^0(\mathbb{R})$  and for  $x \in I, \xi \in \mathbb{R}$  the principal symbol is given by

$$P(x, \xi) = \frac{B(x, \xi)}{A_m^2(x, \xi)} = \left( \chi^2(x) \sum_{j \text{ s.t. } i \rightsquigarrow j} |\phi'_{i,j}(x)| e^{2\text{Re}(V(\phi_{i,j}(x)))} \frac{A_m^2(\mathfrak{F}_{i,j}(x, \xi))}{A_m^2(x, \xi)} \right). \quad (3.19)$$

The estimate (3.13) gives the following upper bound for any  $R > 0, x \in I$  and  $|\xi| > R$ :

$$|P(x, \xi)| \leq \chi^2(x) C^{2m} \sum_{j, i \rightsquigarrow j} |\phi'_{i,j}(x)| e^{2\text{Re}(V(\phi_{i,j}(x)))} \leq C^{2m} N \theta e^{2V_{\max}}$$

with  $V_{\max} = \max_{x \in I} \text{Re}(V(x))$ .

We apply the  **$L^2$ -continuity theorem for PDO** to  $\hat{P}$  as given<sup>5</sup> in [29, th 4.5 p.42]. The result is that for any  $\varepsilon > 0$ ,

$$\hat{P} = \hat{k}_\varepsilon + \hat{p}_\varepsilon$$

with  $\hat{k}_\varepsilon$  a smoothing operator (hence compact) and  $\|\hat{p}_\varepsilon\| \leq C^{2m} N \theta e^{2V_{\max}} + \varepsilon$ .

If  $\hat{Q}_m = \hat{U} \left| \hat{Q}_m \right|$  is the polar decomposition of  $\hat{Q}_m$ , with  $\hat{U}$  unitary, then from (3.16),  $\hat{P} = \left| \hat{Q}_m \right|^2$ , hence  $\left| \hat{Q}_m \right| = \sqrt{\hat{P}}$  and the spectral theorem [53, p.75] gives that  $\left| \hat{Q}_m \right|$  has a similar decomposition

$$\left| \hat{Q}_m \right| = \hat{k}'_\varepsilon + \hat{q}_\varepsilon$$

with  $\hat{k}'_\varepsilon$  compact and  $\|\hat{q}_\varepsilon\| \leq \sqrt{C^{2m} N \theta e^{2V_{\max}} + \varepsilon}$ , with any  $\varepsilon > 0$ . Since  $\|\hat{U}\| = 1$  we deduce a similar decomposition for  $\hat{Q}_m = \hat{U} \left| \hat{Q}_m \right| : L^2(I) \rightarrow L^2(I)$ , i.e.  $\hat{Q}_m = \hat{k}''_\varepsilon + \hat{q}'_\varepsilon$ . We also use the fact that  $C \rightarrow \theta$  as  $R \rightarrow \infty$  in (3.13) to get that  $\|\hat{q}'_\varepsilon\| \leq r_m := c(\theta + \varepsilon)^m$  with  $c$  independent on  $m$  and any  $\varepsilon > 0$ . Equivalently from the diagram (3.15), this gives that

---

<sup>5</sup>Actually, we can not apply directly the  $L^2$ -continuity theorem [29, th 4.5 p.42] for PDO to  $\hat{P}$  because  $\hat{P}$  doesn't have a compactly supported Schwartz kernel. However  $\hat{B}$  obviously has a compactly supported Schwartz kernel due to the presence of  $\hat{\chi}$  in Eq.(3.17). The trick is to approximate  $\hat{A}_m^{-1}$  by a properly supported operator  $\Lambda_m$  as it is done in [29, p.45] and then apply the  $L^2$ -continuity theorem to  $\hat{\Lambda}_m \hat{B} \hat{\Lambda}_m$ .

$\hat{F}_{\hbar,\chi} : H^{-m}(\mathbb{R}) \rightarrow H^{-m}(\mathbb{R})$  can be written  $\hat{F}_{\hbar,\chi} = \hat{K} + \hat{R}$  with  $\hat{K}$  compact and  $\|\hat{R}\| \leq r_m$ . We have obtained (2.16) and (2.17).

The fact that the eigenvalues  $\lambda_i$  and their generalized eigenspaces do not depend on the choice of space  $H^{-m}(\mathbb{R})$  is due to density of  $C_0^\infty(\mathbb{R})$  in Sobolev spaces. We refer to the argument given in the proof of corollary 1 in [23].

Finally, if  $\varphi$  is an eigendistribution of  $\hat{F}_{\hbar,\chi}$ , i.e.  $\hat{F}_{\hbar,\chi}\varphi = \lambda\varphi$  with  $\lambda \neq 0$ , we deduce that  $\varphi = \frac{1}{\lambda^n} \hat{F}_{\hbar,\chi}^n \varphi$  for any  $n \geq 1$ , and (3.4) implies that  $\text{supp}(\varphi) \subset K = \bigcap_{n \in \mathbb{N}} K_n$ . On the trapped set we have  $\chi = 1$  hence the eigendistribution and eigenvalues of  $\hat{F}_{\hbar,\chi}$  do not depend on  $\chi$ . This finishes the proof of Theorem 2.6.

## 4 Dynamics of the canonical map $F : T^*I \rightarrow T^*I$

In Theorem 2.6 the operator  $\hat{F}_{\hbar,\chi}$  is considered for a fixed  $\hbar$ . On the contrary, in Theorems 2.9 and 2.16 they are considered as a family of  $\hbar$ -Fourier integral operators (FIOs) and we give partial results on the distribution of their Ruelle resonances in the semiclassical limit  $\hbar \rightarrow 0$ . As a consequence the oscillations of the phase multiplication by  $e^{\frac{i}{\hbar}\tau(x)}$  are not uniformly bounded anymore and contrary to Proposition 3.4 this multiplication operator is not a pseudodifferential operator anymore but contributes to the phase space dynamics of the canonical map. In this section we will introduce this canonical map and study its dynamics in phase space which becomes significantly more complicated compared to the map  $\mathfrak{F}$  which appeared in Proposition 3.4. In Section 4.1 we will introduce the trapped set in phase space and will naturally be led to the minimal captive property. Then we will introduce the symbolic dynamics in Section 4.2 which nicely describes the dynamics in the cotangent space and which is a central technical ingredient in the proofs of Theorem 2.9 and 2.16. Finally in Section 4.4 we will see that under the assumption of minimal captivity, the trapped set in phase space has also a fractal structure and that its Hausdorff dimension equals twice the Hausdorff dimension of the trapped set  $K$  of the underlying I.F.S.

We first recall that on  $\mathbb{R}^n$ , a  $\hbar$ -**Fourier integral operator** (F.I.O) is a linear operator  $\hat{F} : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$  of the form [57, thm 10.4]

$$\left(\hat{F}\varphi\right)(x') = \frac{1}{(2\pi\hbar)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{\frac{i}{\hbar}(\Phi(x',\xi) - x \cdot \xi)} b(x', \xi; \hbar) \varphi(x) dx d\xi \quad (4.1)$$

where  $\Phi(x', \xi)$  is real valued and called the “phase function” and  $b(x', \xi; \hbar)$  is the amplitude. The Fourier integral operator  $\hat{F}$  has an associated canonical map which is the symplectic map  $F : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ ,  $(x', \xi') = F(x, \xi)$  given by [57, Lemma 10.5]

$$\xi' = (\partial_{x'}\Phi)(x', \xi), \quad x = (\partial_\xi\Phi)(x', \xi) \quad (4.2)$$

*Remark 4.1.* As explained in [12] one interpretation of the canonical map is the following. Since we are interested in the situation of high frequencies we write  $\xi/\hbar$  for the frequency with  $\hbar \ll 1$ . In particular the  $\hbar$ -Fourier transform of a function  $u$  is  $(\mathcal{F}\varphi)(\xi) := \frac{1}{(2\pi\hbar)^{n/2}} \int_{\mathbb{R}^n} e^{-i\frac{\xi}{\hbar} \cdot x} \varphi(x) dx$ . If a  $\hbar$ -family of functions  $\varphi_\hbar$  is *micro-localized* at



point  $x \in \mathbb{R}^n$  and its  $\hbar$ -Fourier transform is *micro-localized* at point  $\xi \in T_x^*\mathbb{R}^n$ , which means that these functions decay fast outside these points as  $\hbar \rightarrow 0$ , then the operator  $\hat{F}$  transforms these functions  $\varphi_\hbar$  to functions  $\hat{F}\varphi_\hbar$  micro-localized in another point  $(x', \xi') = F(x, \xi) \in T^*\mathbb{R}^n$  where  $F$  is the associated canonical map.

According to this previous definition, we give now the canonical map  $F$  for the family of  $\hbar$ -FIOs  $(\hat{F}_\hbar)_\hbar$  that concern us and that were defined in (2.12).

**Lemma 4.2.** *The family of operators  $(\hat{F}_\hbar)_\hbar$  restricted to  $C_0^\infty(I)$  is a  $\hbar$ -Fourier integral operator (FIO). Its **canonical map** is a multi-valued symplectic map  $F : T^*I \rightarrow T^*I$  (with  $T^*I \cong I \times \mathbb{R}$ ) given by:*

$$F : \begin{cases} T^*I & \rightarrow T^*I \\ (x, \xi) & \mapsto \{F_{i,j}(x, \xi) \quad \text{with } i, j \text{ s.t. } x \in I_i, i \rightsquigarrow j\} \end{cases} \quad (4.3)$$

with

$$F_{i,j} : \begin{cases} x' & = \phi_{i,j}(x) \\ \xi' & = \frac{1}{\phi'_{i,j}(x)}\xi + \frac{d\tau}{dx}(x') \end{cases} \quad (4.4)$$

*Proof.* From (2.12),

$$\left(\hat{F}_{i,j}\varphi\right)(x') = e^{V(x')} e^{i\frac{1}{\hbar}\tau(x')} \varphi\left(\phi_{i,j}^{-1}(x')\right) = \left(\hat{F}_2 \circ \hat{F}_1\varphi\right)(x')$$

that we have decomposed into a first operator (we will use  $\delta(x) = \frac{1}{(2\pi\hbar)} \int_{\mathbb{R}} e^{\frac{i}{\hbar}x \cdot \xi} d\xi$ )

$$\begin{aligned} \left(\hat{F}_1\varphi\right)(x') &:= \varphi\left(\phi_{i,j}^{-1}(x')\right) = \int_{\mathbb{R}} \delta\left(\phi_{i,j}^{-1}(x') - x\right) \varphi(x) dx \\ &= \frac{1}{(2\pi\hbar)} \int_{\mathbb{R}^2} e^{\frac{i}{\hbar}(\phi_{i,j}^{-1}(x') \cdot \xi - x \cdot \xi)} \varphi(x) dx d\xi \end{aligned}$$

which shows from (4.1) that  $\hat{F}_1$  is a F.I.O. with amplitude  $b = 1$  and phase function  $\Phi(x', \xi) = \phi_{i,j}^{-1}(x') \cdot \xi$ . Its canonical map is then  $(x', \xi') = F_1(x, \xi)$  given from (4.2) by

$$\xi' = (\partial_{x'}\Phi)(x', \xi) = \frac{1}{\phi'_{i,j}(x)}\xi, \quad x = (\partial_\xi\Phi)(x', \xi) = \phi_{i,j}^{-1}(x').$$

Similarly for the second operator we write:

$$\begin{aligned} \left(\hat{F}_2\varphi\right)(x') &:= e^{V(x')} e^{i\tau(x')/\hbar} \varphi(x') = \int_{\mathbb{R}} e^{V(x')} e^{i\tau(x')/\hbar} \delta(x' - x) \varphi(x) dx \\ &= \frac{1}{(2\pi\hbar)} \int_{\mathbb{R}^2} e^{\frac{i}{\hbar}(x' \cdot \xi + \tau(x') - x \cdot \xi)} e^{V(x')} \varphi(x) dx d\xi \end{aligned}$$

which shows from (4.1) that  $\hat{F}_2$  is a F.I.O. with amplitude  $b(x', \xi; \hbar) = e^{V(x')}$  and phase function  $\Phi(x', \xi) = x' \cdot \xi + \tau(x')$ . Its canonical map is then  $(x', \xi') = F_2(x, \xi)$  given from (4.2) by

$$\xi' = (\partial_{x'} \Phi)(x', \xi) = \xi + \tau'(x'), \quad x = (\partial_\xi \Phi)(x', \xi) = x'$$

By composition [29, th.11.12] we deduce that  $\hat{F}_{i,j} = \hat{F}_2 \circ \hat{F}_1$  is a FIO with canonical map  $F_{i,j} = F_2 \circ F_1$  given by (4.4).  $\square$

*Remark 4.3.* For short, we can write

$$F : \begin{cases} T^*I & \rightarrow T^*I \\ (x, \xi) & \rightarrow \left( \phi(x), \frac{1}{\phi'(x)}\xi + \tau'(\phi(x)) \right) \end{cases}. \quad (4.5)$$

We will study the dynamics of  $F$  in detail in later sections, but we can already make some remarks. The term  $\frac{dx}{dx}(x')$  in the expression of  $\xi'$ , Eq.(4.4), complicates significantly the dynamics near the zero section  $\xi = 0$ . However the next Lemma shows that a trajectory from an initial point  $(x, \xi)$  with  $|\xi|$  large enough, escapes towards infinity:

**Lemma 4.4.** *For any  $1 < \kappa < 1/\theta$ , with  $\theta$  defined in (2.1), there exists  $R \geq 0$  such that for any  $(x, \xi)$ , with  $|\xi| > R$  and any  $i \rightsquigarrow j$ , with  $x \in I_i$ , we have*

$$|\xi'| > \kappa |\xi| \quad (4.6)$$

where  $(x', \xi') = F_{i,j}(x, \xi)$ .

*Proof.* From (4.4), one has  $\xi' = \frac{1}{\phi'_{i,j}(x)}\xi + \tau'(x')$ . Also  $\left| \frac{1}{\phi'_{i,j}(x)} \right| \geq \theta^{-1}$  hence

$$\begin{aligned} |\xi'| - \kappa |\xi| &= \left| \frac{1}{\phi'_{i,j}(x)}\xi + \tau'(x') \right| - \kappa |\xi| \geq \left| \frac{1}{\phi'_{i,j}(x)}\xi \right| - |\tau'(x')| - \kappa |\xi| \\ &\geq \left( \frac{1}{\theta} - \kappa \right) |\xi| - \max_x |\tau'(x)| > 0. \end{aligned}$$

The last inequality holds true if  $|\xi| > R := \left( \frac{1}{\theta} - \kappa \right)^{-1} \max_x |\tau'|$ .  $\square$

## 4.1 The trapped set $\mathcal{K}$ in phase space

**Definition 4.5.** The **trapped set** in phase space  $T^*I$  is defined as

$$\mathcal{K} = \{ (x, \xi) \in T^*I, \exists C \Subset T^*I \text{ compact}, \forall n \in \mathbb{Z}, F^n(x, \xi) \cap C \neq \emptyset \}. \quad (4.7)$$

*Remark 4.6.* Since the map  $F : T^*I \rightarrow T^*I$  is a lift of the map  $\phi$ , we have  $\mathcal{K} \subset (K \times \mathbb{R})$ . We can precise this: for any  $R$  given from Lemma 4.4,

$$\mathcal{K} \subset (K \times [-R, R]).$$

For  $\varepsilon > 0$ , let  $\mathcal{K}_\varepsilon$  denote a  $\varepsilon$ -neighborhood of the trapped set  $\mathcal{K}$ , namely

$$\mathcal{K}_\varepsilon := \{(x, \xi) \in T^*I, \quad \exists (x_0, \xi_0) \in \mathcal{K}, \quad \max(|x - x_0|, |\xi - \xi_0|) \leq \varepsilon\}.$$

Recall that the canonical map  $F$  is multivalued. The definition of the trapped set requires that at least one of the future trajectories of points in  $\mathcal{K}$  stays bounded. The following assumption on the map basically demands, that exactly one trajectory stays bounded.

**Assumption 4.7.** *We assume the following property called “minimal captivity”:*

$$\exists \varepsilon > 0, \quad \forall (x, \xi) \in \mathcal{K}_\varepsilon, \quad \#\left\{F(x, \xi) \cap \mathcal{K}_\varepsilon\right\} \leq 1. \quad (4.8)$$

*This means that the dynamics of  $F$  is univalued on the trapped set  $\mathcal{K}$ .*

**Remarks:**

- In the paper [21] the second author introduced the property of “**partial captivity**” which is weaker than “**minimal captivity**”: partial captivity roughly states that most of trajectories escape from the trapped set  $\mathcal{K}$  whereas minimal captivity states that every trajectory except one, escapes from the trapped set  $\mathcal{K}$ .
- Note that the complexity of the dynamics of the map  $F$  in (4.5) is due to the term  $\tau'(\phi_{i,j}(x))$ , so the minimal captive property can also be considered as a condition on the behavior of the roof function along the trajectories of the I.F.S. In particular for trivial (i.e. constant) roof functions the condition can not be fulfilled. In this case, the canonical map  $F$  equals the simpler map  $\mathfrak{F}$  of Proposition 3.4. Then the trapped set is given by  $\mathcal{K} = K \times \{0\}$  and all trajectories in the trapped set stay on the trapped set. The same holds for all roof functions, that are cohomologous to a constant (c.f. [21, Appendix A]).

We will give now a more precise description of the trapped set  $\mathcal{K}$ . Recall that the inverse maps  $\phi^{-1}$  and  $F^{-1}$  are uni-valued. For any integer  $m \geq 0$ , let

$$\tilde{K}_m := F^{-m}(K_m \times [-R, R])$$

where  $K_m = \phi^m(I)$  has been defined in (2.6) and  $R$  is given by Lemma 4.4. In particular  $\tilde{K}_0 = I \times [-R, R]$ . Let  $\pi : (x, \xi) \in T^*I \rightarrow x \in I$  be the projection map. From (4.5) we have:

$$\pi(\tilde{K}_m) = I,$$

and a short computation<sup>6</sup> gives

$$\tilde{K}_{m+1} \subset \tilde{K}_m. \quad (4.9)$$

Let us define

$$\tilde{K} := \bigcap_m \tilde{K}_m. \quad (4.10)$$

Now we combine the sets  $K_n$  defined in (2.6) with the sets  $\tilde{K}_m$  and define for any integers  $a, b \geq 0$

$$\mathcal{K}_{a,b} := \pi^{-1}(K_a) \bigcap \tilde{K}_b. \quad (4.11)$$

We have

$$\mathcal{K}_{a+1,b} \subset \mathcal{K}_{a,b}, \quad \mathcal{K}_{a,b+1} \subset \mathcal{K}_{a,b} \quad (4.12)$$

and

$$F^{-1}(\mathcal{K}_{a,b}) = \mathcal{K}_{a-1,b+1}. \quad (4.13)$$

*Remark 4.8.* We can interpret the trapped set  $K \subset I$  with respect to the lifted map  $F : T^*I \rightarrow T^*I$ , as follows. The trapped set  $\pi^{-1}(K) \subset T^*I$  is characterized by

$$\pi^{-1}(K) = \{(x, \xi) \in T^*I, \exists \text{compact } C \Subset T^*I, \forall n \geq 0, F^{-n}(x, \xi) \in C\}$$

i.e.  $\pi^{-1}(K)$  can be considered as the “trapped set of the map  $F$  in the past”. Similarly  $\tilde{K} \subset T^*I$  can be interpreted as the “trapped set of the map  $F$  in the future” and  $\mathcal{K} \subset T^*I$  as the full trapped set (past and future) since they are characterized by

$$\tilde{K} = \{(x, \xi) \in T^*I, \exists \text{compact } C \Subset T^*I, \forall n \geq 0, F^n(x, \xi) \cap C \neq \emptyset\}$$

$$\begin{aligned} \mathcal{K} &= \{(x, \xi) \in T^*I, \exists \text{compact } C \Subset T^*I, \forall n \in \mathbb{Z}, F^n(x, \xi) \cap C \neq \emptyset\} \\ &= \pi^{-1}(K) \cap \tilde{K}. \end{aligned} \quad (4.14)$$

From this previous remark, we have the following expression for the trapped set equivalent to (4.7).

**Proposition 4.9.** *The trapped set  $\mathcal{K} \subset T^*I$  of the map  $F$  is*

$$\mathcal{K} = \bigcap_{a=0}^{\infty} \mathcal{K}_{a,a} \quad (4.15)$$

---

<sup>6</sup>From Lemma 4.4 we have

$$(K_{m+1} \times [-R, R]) \subset F(K_m \times [-R, R])$$

hence

$$\tilde{K}_{m+1} = F^{-m}(F^{-1}(K_{m+1} \times [-R, R])) \subset F^{-m}(K_m \times [-R, R]) = \tilde{K}_m.$$

The hypothesis of minimal captivity has been defined in Assumption 4.7. The following proposition gives an equivalent and a slightly weaker definition of minimal captivity that will be used in Section 5.2.

**Proposition 4.10.**

1. The map  $F$  is minimally captive (i.e. Eq.(4.8) holds true) if and only if the map  $F$  satisfies

$$\exists a, \quad \forall (x, \xi) \in \mathcal{K}_{a,a}, \quad \# \left\{ F(x, \xi) \cap \mathcal{K}_{a,a} \right\} \leq 1. \quad (4.16)$$

2. If map  $F$  is minimally captive then

$$\exists a, \exists C, \text{ s.t. } \forall (x, \xi) \in \mathcal{K}_{a,0}, \forall n, \quad \# \left\{ F^n(x, \xi) \cap \mathcal{K}_{a,0} \right\} \leq C. \quad (4.17)$$

where  $\mathcal{K}_{a,0} := (\pi^{-1}(K_a) \cap [-R, R])$  has been defined in (4.11).

*Proof.* The fact that (4.16) is equivalent to (4.8) is because

$$\begin{aligned} \forall \varepsilon > 0, \exists a \text{ s.t. } \mathcal{K}_{a,a} \subset \mathcal{K}_\varepsilon \\ \forall a, \exists \varepsilon > 0 \text{ s.t. } \mathcal{K}_\varepsilon \subset \mathcal{K}_{a,a} \end{aligned}$$

Now we prove (4.17). Let  $a > 0$  be an even integer. Let  $(x, \xi) \in \mathcal{K}_{a,0}$  and  $n > \frac{a}{2}$ . We write  $F^n(x, \xi) = F^{a/2}(F^{n-a/2}(x, \xi))$ . Let  $(x', \xi') \in F^{n-a/2}(x, \xi)$ . From (4.13), we have  $F^{-a/2}(\mathcal{K}_{a,0}) = \mathcal{K}_{\frac{a}{2}, \frac{a}{2}}$ , hence if  $(x', \xi') \notin \mathcal{K}_{\frac{a}{2}, \frac{a}{2}}$  then  $F^{a/2}(x', \xi') \notin \mathcal{K}_{a,0}$ . On the contrary, for  $(x', \xi') \in \mathcal{K}_{\frac{a}{2}, \frac{a}{2}}$  then the set  $F^{a/2}(x', \xi')$  has cardinal less than  $N^{a/2}$ , so we obtain

$$\# \{ F^n(x, \xi) \cap \mathcal{K}_{a,0} \} \leq N^{a/2} \cdot \# \{ F^{n-a/2}(x, \xi) \cap \mathcal{K}_{\frac{a}{2}, \frac{a}{2}} \}.$$

So if  $\# \{ F^{n-a/2}(x, \xi) \cap \mathcal{K}_{\frac{a}{2}, \frac{a}{2}} \} = \emptyset$  we are done with the proof. Suppose on the contrary that  $\# \{ F^{n-a/2}(x, \xi) \cap \mathcal{K}_{\frac{a}{2}, \frac{a}{2}} \} \neq \emptyset$ . From (4.13) we have  $(x, \xi) \in \mathcal{K}_{0,n} \cap \mathcal{K}_{a,0} \subset \mathcal{K}_{\frac{a}{2}, \frac{a}{2}}$ . Finally we suppose that assumption that (4.16) is fulfilled for  $\frac{a}{2}$ . This gives that

$$\# \{ F^{n-a/2}(x, \xi) \cap \mathcal{K}_{\frac{a}{2}, \frac{a}{2}} \} \leq 1$$

and  $\# \{ F^n(x, \xi) \cap \mathcal{K}_{a,0} \} \leq N^{a/2}$ . We have obtained (4.17) with the bound  $C = N^{a/2}$ .  $\square$

## 4.2 Symbolic dynamics

The purpose of this section is to describe precisely the dynamics of  $\phi$  and  $F$  using “symbolic dynamics”. This is very standard for expanding maps [11]. This description refines the structure of the sets  $\mathcal{K}_{a,b}$  introduced before. We would like to emphasize that the use of symbolic dynamics in this paper is due to related to the fact that the initial I.F.S. model in Definition 2.1 is a multivalued map  $\phi$  defined on a union of intervals  $(I_i)_{i=1 \dots N}$ . This is not a “discontinuous Markov partition of a continuous dynamics” [11, p.134].

### 4.2.1 Symbolic dynamics on the trapped set $K \subset I$

Let

$$\mathcal{W}_- := \left\{ (\dots, w_{-2}, w_{-1}, w_0) \in \{1, \dots, N\}^{-\mathbb{N}}, w_{l-1} \rightsquigarrow w_l, \forall l \leq 0 \right\} \quad (4.18)$$

be the set of **admissible left semi-infinite sequences**. For  $w \in \mathcal{W}_-$  and  $i < j$  we write  $w_{i,j} := (w_i, w_{i+1}, \dots, w_j)$  for an extracted sequence. For simplicity we will use the notation

$$\phi_{w_{i,j}} := \phi_{w_{j-1}, w_j} \circ \dots \circ \phi_{w_i, w_{i+1}} : I_{w_i} \rightarrow I_{w_j} \quad (4.19)$$

for the composition of maps. For  $n \geq 0$ , let

$$I_{w_{-n,0}} := \phi_{w_{-n,0}}(I_{w_{-n}}) \subset I_{w_0}. \quad (4.20)$$

For any  $0 < m < n$  we have the strict inclusions

$$I_{w_{-n,0}} \subset I_{w_{-m,0}} \subset I_{w_0}.$$

From (2.1), the size of  $I_{w_{-n,0}}$  is bounded by

$$|I_{w_{-n,0}}| \leq \theta^n |I_{w_0}|,$$

hence the sequence of sets  $(I_{w_{-n,0}})_{n \geq 1}$  is a sequence of non empty and decreasing closed intervals and  $\bigcap_{n=1}^{\infty} I_{w_{-n,0}}$  is a point in  $K$ . We define

**Definition 4.11.** The “symbolic coding map” is

$$S : \begin{cases} \mathcal{W}_- & \rightarrow K \\ w & \mapsto S(w) := \bigcap_{n=1}^{\infty} I_{w_{-n,0}}. \end{cases} \quad (4.21)$$

In some sense we have decomposed the sets  $K_n$ , Eq.(2.6), into individual components:

$$K_n = \bigcup_{w_{-n,0} \in \mathcal{W}_-} I_{w_{-n,0}} \quad (4.22)$$

$$K = \bigcup_{w \in \mathcal{W}_-} S(w).$$

Let us introduce the **left shift**, a multivalued map, defined by

$$L : \begin{cases} \mathcal{W}_- & \rightarrow \mathcal{W}_- \\ (\dots, w_{-2}, w_{-1}, w_0) & \mapsto (\dots, w_{-2}, w_{-1}, w_0, w_1) \end{cases}$$

with  $w_1 \in \{1, \dots, N\}$  such that  $w_0 \rightsquigarrow w_1$ . Let the **right shift** be the univalued map defined by

$$R : \begin{cases} \mathcal{W}_- & \rightarrow \mathcal{W}_- \\ (\dots, w_{-2}, w_{-1}, w_0) & \mapsto (\dots, w_{-2}, w_{-1}) \end{cases}.$$

**Proposition 4.12.** *The following diagram is commutative*

$$\begin{array}{ccc} \mathcal{W}_- & \xrightarrow{S} & K \\ R \uparrow \downarrow L & & \phi^{-1} \uparrow \downarrow \phi \\ \mathcal{W}_- & \xrightarrow{S} & K \end{array} \quad (4.23)$$

and the map  $S : \mathcal{W}_- \rightarrow K$  is one to one. This means that the dynamics of points on the trapped set  $K$  under the maps  $\phi^{-1}, \phi$  is equivalent to the symbolic dynamics of the shift maps  $R, L$  on the set of admissible words  $\mathcal{W}_-$ . Notice that the maps  $R$  and  $\phi^{-1}$  are univalued, whereas the maps  $L$  and  $\phi$  are (in general) multivalued.

*Proof.* From the definition of  $S$  we have

$$\phi_{w_0 w_1} (S(\dots, w_{-2}, w_{-1}, w_0)) = S(\dots, w_{-2}, w_{-1}, w_0, w_1) \quad (4.24)$$

and

$$\phi_{w_{-1} w_0}^{-1} (S(\dots, w_{-2}, w_{-1}, w_0)) = S(\dots, w_{-2}, w_{-1}) \quad (4.25)$$

which gives the diagram (4.23). The map  $S : \mathcal{W}_- \rightarrow K$  is surjective by construction. Let us show that the hypothesis (2.2) implies that it is also injective. Let  $w, w' \in \mathcal{W}_-$  and suppose that  $w \neq w'$ , i.e. there exists  $k \geq 0$  such that  $w_{-k} \neq w'_{-k}$ . From (2.2) we have  $\phi_{w_{-k}, w_{-k+1}} (I_{w_{-k}}) \cap \phi_{w'_{-k}, w'_{-k+1}} (I_{w'_{-k}}) = \emptyset$ . We deduce recursively that  $\phi_{w_{-k}, 0} (I_{w_{-k}}) \cap \phi_{w'_{-k}, 0} (I_{w'_{-k}}) = \emptyset$ . Since  $S(w) \in \phi_{w_{-k}, 0} (I_{w_{-k}})$  and  $S(w') \in \phi_{w'_{-k}, 0} (I_{w'_{-k}})$  we deduce that  $S(w) \neq S(w')$ . Hence  $S$  is one to one.  $\square$

#### 4.2.2 The “future trapped set” $\tilde{K}$ in phase space $T^*I$

Let

$$\mathcal{W}_+ := \left\{ (w_0, w_1, w_2 \dots) \in \{1, \dots, N\}^{\mathbb{N}}, \quad w_l \rightsquigarrow w_{l+1}, \forall l \geq 0 \right\}$$

be the set of admissible right semi-infinite sequences. We still use the notation  $w_{i,j} := (w_i, w_{i+1}, \dots, w_j)$  for an extracted sequence. For any  $n \geq 0$  let

$$\tilde{I}_{w_0, n} := F^{-n} (I_{w_0, n} \times [-R, R]) \quad (4.26)$$

be the image of the rectangle under the univalued map  $F^{-n}$ . Notice that  $\pi(\tilde{I}_{w_0, n}) = I_{w_0}$  where  $\pi(x, \xi) = x$  is the canonical projection map. The map  $F^{-1}$  contracts strictly in the  $\xi$ -variable by the factor  $\theta < 1$  thus  $(\tilde{I}_{w_0, n})_{n \in \mathbb{N}}$  is a sequence of decreasing sets:  $\tilde{I}_{w_0, n+1} \subset \tilde{I}_{w_0, n}$  and we can define the limit

$$\tilde{S} : w \in \mathcal{W}_+ \rightarrow \tilde{S}(w) := \bigcap_{n \geq 0} \tilde{I}_{w_0, n} \subset \tilde{K}. \quad (4.27)$$

**Proposition 4.13.** *For every  $w \in \mathcal{W}_+$ , the set  $\tilde{S}(w)$  is a smooth curve given by*

$$\tilde{S}(w) = \{(x, \zeta_w(x)), \quad x \in I_{w_0}, w \in \mathcal{W}_+\}$$

with

$$\zeta_w(x) = - \sum_{k \geq 1} \phi'_{w_0, k}(x) \cdot \tau'(\phi_{w_0, k}(x)). \quad (4.28)$$

We have an estimate of regularity, uniform in  $w$ :  $\forall \alpha \in \mathbb{N}$ ,  $\exists C_\alpha > 0$ , such that  $\forall w \in \mathcal{W}_+$ ,  $\forall x \in I_{w_0}$ ,

$$|(\partial_x^\alpha \zeta_w)(x)| \leq C_\alpha. \quad (4.29)$$

Moreover, with the Assumption 4.7 of minimal captivity there exists  $a \geq 1$  such that these branches do not intersect on  $\pi^{-1}(K_a)$ ,

$$\forall w, w' \in \mathcal{W}_+, \quad w \neq w' \Rightarrow \pi^{-1}(K_a) \cap \tilde{S}(w) \cap \tilde{S}(w') = \emptyset. \quad (4.30)$$

The set (4.10) can be expressed as

$$\tilde{K} = \bigcup_{w \in \mathcal{W}_+} \tilde{S}(w).$$

*Proof.* From (4.4) we get

$$F^{-1}(\phi_{i,j}(x), \xi) = (x, \phi'_{i,j}(x) (\xi - \tau'(\phi_{i,j}(x))))). \quad (4.31)$$

Iterating this equation we get, that

$$\zeta_{w,n}(x) := - \sum_{k=1}^n \phi'_{w_0, k}(x) \cdot \tau'(\phi_{w_0, k}(x)),$$

fulfills

$$(x, \zeta_{w,n}(x)) = F^{-n}(\phi_{w_0, n}(x), 0),$$

thus  $(x, \zeta_{w,n}(x)) \in \tilde{I}_{w_0, n}$  for all  $n \in \mathbb{N}$  and we get (4.28).

In order to prove (4.29) we can check, that the series of  $\zeta_{w,n}(x)$  and  $\partial_x^\alpha \zeta_{w,n}(x)$  converge with uniform bounds in  $w$  which follows after some calculations from (2.1) and the fact that  $|\phi'_{w_0, k}(x)| \leq \theta^k$  independent of  $w$ .

In order to see (4.30) let  $w, w' \in \mathcal{W}_+$  with  $w \neq w'$  and  $n \in \mathbb{N}$  such that  $w_0, n \neq w'_0, n$  and suppose that there is  $x \in K_a$  and  $\xi \in \mathbb{R}$  such that  $(x, \xi) \in \tilde{S}(w) \cap \tilde{S}(w')$ . Then by



the definition of  $\tilde{S}$ ,  $(x, \xi) \in \tilde{I}_{w_{0,n+a}} \cap \tilde{I}'_{w'_{0,n+a}} \subset \tilde{K}_{n+a}$ . Consequently there are  $(x_1, \xi_1) \in I_{w_{0,n+a}} \times [-R, R]$  and  $(x_2, \xi_2) \in I'_{w'_{0,n+a}} \times [-R, R]$  with  $(x, \xi) = F^{-n-a}(x_1, \xi_1) = F^{-n-a}(x_2, \xi_2)$  and we have

$$F^{-a}(x_1, \xi_1), F^{-a}(x_2, \xi_2)F^n \in (x, \xi).$$

But as  $F^{-a}(x_1, \xi_1) \in \pi^{-1}(I_{w_{0,n}})$  and  $F^{-a}(x_2, \xi_2) \in \pi^{-1}(I'_{w'_{0,n}})$  we clearly have  $F^{-a}(x_1, \xi_1) \neq F^{-a}(x_2, \xi_2)$  because  $w_{0,n} \neq w'_{0,n}$ . And additionally we have chosen  $(x, \xi) \in \mathcal{K}_{a,n+a}$  and from the definition of  $(x_1, \xi_1)$  and  $(x_2, \xi_2)$  we get  $F^{-a}(x_1, \xi_1), F^{-a}(x_2, \xi_2) \in \mathcal{K}_{n+a,a}$ . Thus we have found  $(x, \xi) \in \mathcal{K}_{a,a}$  with  $\#\{F^n(x, \xi) \cap \mathcal{K}_{a,a}\} \geq 2$  which contradicts Assumption 4.7.  $\square$

### 4.2.3 Symbolic dynamics on the trapped set $\mathcal{K}$ in phase space $T^*I$

Recall from (4.14) that  $\mathcal{K} = \pi^{-1}(K) \cap \tilde{K}$ . Let

$$\mathcal{W} := \left\{ (\dots w_{-2}, w_{-1}, w_0, w_1, \dots) \in \{1, \dots, N\}^{\mathbb{Z}}, \quad w_l \rightsquigarrow w_{l+1}, \forall l \in \mathbb{Z} \right\}$$

be the set of bi-infinite admissible sequences. For a given  $w \in \mathcal{W}$  and  $a, b \in \mathbb{N}$ , let

$$\mathcal{I}_{w_{-a,0}, w_{0,b}} := \left( \pi^{-1}(I_{w_{-a,0}}) \cap \tilde{I}_{w_{0,b}} \right) \subset \mathcal{K}_{a,b}$$

where  $\mathcal{K}_{a,b}$  has been defined in (4.11).

**Definition 4.14.** The symbolic coding map is

$$\mathcal{S} : \begin{cases} \mathcal{W} & \rightarrow \mathcal{K} \\ w & \mapsto \mathcal{S}(w) := \bigcap_{n=1}^{\infty} \mathcal{I}_{w_{-n,0}, w_{0,n}} = \left( \pi^{-1}(S(w_-)) \cap \tilde{S}(w_+) \right) \end{cases} \quad (4.32)$$

with  $w_- = (\dots w_{-1}, w_0) \in \mathcal{W}_-$ ,  $w_+ = (w_0, w_1, \dots) \in \mathcal{W}_+$ .

More precisely we can express the point  $\mathcal{S}(w) \in \mathcal{K}$  as

$$\mathcal{S}(w) = (x_{w_-}, \xi_w), \quad x_{w_-} = S(w_-), \quad \xi_w = \zeta_{w_+}(S(w_-)), \quad (4.33)$$

with  $\zeta_{w_+}$  given in (4.28). We also have

$$\mathcal{K}_{a,b} = \bigcup_{w \in \mathcal{W}} \mathcal{I}_{w_{-a,0}, w_{0,b}}.$$

**Proposition 4.15.** *The following diagram is commutative*

$$\begin{array}{ccc} \mathcal{W} & \xrightarrow{\mathcal{S}} & \mathcal{K} \\ R \uparrow \downarrow L & & F^{-1} \uparrow \downarrow F \\ \mathcal{W} & \xrightarrow{\mathcal{S}} & \mathcal{K}. \end{array} \quad (4.34)$$

If Assumption 4.7 of minimal captivity holds true then the map  $\mathcal{S} : \mathcal{W} \rightarrow \mathcal{K}$  is one to one. This means that the univalued dynamics of points on the trapped set  $\mathcal{K}$  under the maps  $F^{-1}, F$  is equivalent to the symbolic dynamics of the full shift maps  $R, L$  on the set of words  $\mathcal{W}$ .

*Proof.* Commutativity of the diagram comes from the construction of  $\mathcal{S}$ . Also  $\mathcal{S}$  is surjective. Let us show that  $\mathcal{S}$  is injective. Let  $w, w' \in \mathcal{W}$ , with  $w \neq w'$ . There exists  $n \geq 0$  such that  $(L^n(w))_- \neq (L^n(w'))_-$ . So  $\mathcal{S}((L^n(w))_-) \neq \mathcal{S}((L^n(w'))_-)$  because  $\mathcal{S} : \mathcal{W}_- \rightarrow \mathcal{K}$  is one to one from Lemma 4.12. Hence  $\mathcal{S}(L^n(w)) \neq \mathcal{S}(L^n(w'))$  and  $F^n(\mathcal{S}(w)) \neq F^n(\mathcal{S}(w'))$  from commutativity of the diagram. We apply  $F^{-n}$  and deduce that  $\mathcal{S}(w) \neq \mathcal{S}(w')$  because  $F^{-1}$  and  $F^{-n}$  are injective on  $\mathcal{K}$  from Assumption 4.7.  $\square$

### 4.3 Relation to the non-local integrability condition of Dolgopyat

We can now discuss the relation of the minimal captive assumption and the non-local integrability (NLI) condition used by Naud and Dolgopyat [40, 16] in order to obtain exponential decay of correlation. For the discussion we use the version of the NLI-condition introduced in [40] where Naud first introduces for a symbolic sequence  $w \in \mathcal{W}_+$  and  $u, v \in I_{w_0}$  the quantity

$$\Delta_w(u, v) := \sum_{k=1}^{\infty} \tau(\phi_{w_0, k}(u)) - \tau(\phi_{w_0, k}(v)) \quad (4.35)$$

as well as the temporal distance function for  $w, w' \in \mathcal{W}_+$  with  $w_0 = w'_0$

$$\varphi_{w, w'}(u, v) := \Delta_w(u, v) - \Delta_{w'}(u, v).$$

According to [40, Definition 2.1] the roof function  $\tau$  fulfills the NLI-condition if there exist  $w, w'$  with  $w_0 = w'_0$  and  $u_0, v_0 \in I_{w_0} \cap K$  such that

$$\frac{\partial \varphi_{w, w'}}{\partial u}(u_0, v_0) \neq 0.$$

Note that (4.35) implies that

$$\frac{\partial \Delta_w}{\partial u}(u_0, v_0) = -\zeta_w(u_0)$$

where  $\zeta_w$  are exactly the functions defined in (4.28). Thus translated to the language of our article, the NLI-condition is the existence of two words  $w, w' \in \mathcal{W}_+$  with  $w_0 = w'_0$  and a point in the trapped set  $u_0 \in I_{w_0} \cap K$  such that  $\zeta_w(u_0) \neq \zeta_{w'}(u_0)$  (i.e. the two branches  $\zeta_w(u_0), \zeta_{w'}(u_0)$  above the point  $u_0 \in K$  are disjoint). In Proposition 4.15 we have however shown, that under the condition of minimal captivity this is true for all  $u_0 \in K$  and for all  $w \neq w'$  we have  $\zeta_w(u_0) \neq \zeta_{w'}(u_0)$ . The minimal captivity assumption thus implies NLI and is much stronger. It has however also stronger implications, for example on the fractal structure of the trapped set  $\mathcal{K}$  as shown in the following section.

#### 4.4 Dimension of the trapped set $\mathcal{K}$

We will now show that the assumption of minimal captivity allows to characterize the fractal structure of the trapped set  $\mathcal{K}$ .

**Proposition 4.16.** *If Assumption 4.7 holds true and if the adjacency matrix  $A$  is symmetric then*

$$\dim_M \mathcal{K} = 2 \dim_M K \quad (4.36)$$

where  $\dim_M B$  stands for the Minkowski dimension of a set  $B$  as defined in Eq.(2.30).

Recall from (2.33) that  $\dim_H K = \dim_M K$ .

For  $w = (w_k)_{k \in \mathbb{Z}} \in \mathcal{W}$ , we note  $w_- = (\dots, w_{-2}, w_{-1}, w_0) \in \mathcal{W}_-$  and  $w_+ = (w_0, w_1, \dots) \in \mathcal{W}_+$ . Let

$$\text{Inv}(w_+) := (\dots w_2, w_1, w_0)$$

be the reversed word. Since the adjacency matrix  $A$  is supposed to be symmetric we have that  $\text{Inv}(w_+) \in \mathcal{W}_-$ . Then, let us consider the following one to one map

$$D : \begin{cases} \mathcal{W} & \rightarrow (\mathcal{W}_- \times \mathcal{W}_-)_l \\ w & \rightarrow (w_-, \text{Inv}(w_+)) \end{cases}$$

where

$$(\mathcal{W}_- \times \mathcal{W}_-)_l := \{(w, w') \in \mathcal{W}_- \times \mathcal{W}_-, \quad w_0 = w'_0\} \quad (4.37)$$

is a subset of  $\mathcal{W}_- \times \mathcal{W}_-$ . The index  $l$  stands for “linked”. Let

$$\Phi := (S \otimes S) \circ D \circ \mathcal{S}^{-1} \quad : \mathcal{K} \rightarrow K \times K$$

where  $\mathcal{S} : \mathcal{W} \rightarrow \mathcal{K}$  has been defined in (4.32) and is shown in Proposition 4.15 to be one to one under Assumption 4.7. The map  $S : \mathcal{W}_+ \rightarrow K$  has been defined in (4.21) and is also one to one. Consider

$$(K \times K)_l := (S \otimes S)((\mathcal{W}_- \times \mathcal{W}_-)_l) \subset K \times K \quad (4.38)$$

the image of (4.37) under the map  $S \otimes S$ . From the previous remarks, the map  $\Phi : \mathcal{K} \rightarrow (K \times K)_l$  is one to one.

**Lemma 4.17.** *The map  $\Phi : \mathcal{K} \rightarrow (K \times K)_l$  is bi-Lipschitz.*

This Lemma is illustrated in Figure 7.1 which shows clearly that the trapped set  $\mathcal{K}$  has a product structure. Before proving Lemma 4.17, let us show how to deduce Proposition 4.16 from it. Since the Hausdorff and Minkowski dimension are invariant under bi-Lipschitz maps [19, p.24], we deduce from this Lemma that

$$\dim_M(\mathcal{K}) = \dim_M(K \times K)_l \quad (4.39)$$

Let us temporarily write  $K_i := K \cap I_i$ . From (4.38) we have that

$$(K \times K)_l = \bigcup_i K_i \times K_i$$

hence

$$\dim_M(K \times K)_l = \max_{1 \leq i \leq N} (2 \dim_M K_i) = 2 \dim_M K \quad (4.40)$$

Eq.(4.39) and (4.40) give Proposition 4.16.

*Proof.* of Lemma 4.17. Let  $w \in \mathcal{W}$ . We write  $w = (w_-, w_+)$  as before and  $x_{w_-} := S(w_-) \in K$ ,  $\rho = (x_{w_-}, \xi_w) = \mathcal{S}(w) \in \mathcal{K}$ . Similarly for another  $w' \in \mathcal{W}$  we get another point  $\rho' = (x_{w'_-}, \xi_{w'}) \in \mathcal{K}$ . We have that

$$\Phi(\rho) = (S(w_-), S(\text{Inv}(w_+))) = (x_{w_-}, x_{\text{Inv}(w_+)}) \in K \times K.$$

That the map  $\Phi$  is bi-Lipschitz means that

$$|\Phi(\rho) - \Phi(\rho')| \asymp |\rho - \rho'|$$

uniformly<sup>7</sup> over  $\rho, \rho'$ . Equivalently this is

$$\left| x_{w_-} - x_{w'_-} \right| + \left| x_{\text{Inv}(w_+)} - x_{\text{Inv}(w'_+)} \right| \asymp \left| x_{w_-} - x_{w'_-} \right| + |\xi_w - \xi_{w'}| \quad (4.41)$$

uniformly over  $w, w' \in \mathcal{W}$ . Let us show (4.41). Let  $w, w' \in \mathcal{W}$  and let  $n \geq 0$  be the integer such that  $(w_+)_j = (w'_+)_j$  for  $0 \leq j \leq n$  but  $(w_+)_{n+1} \neq (w'_+)_{n+1}$ . From the definition (4.20) of the intervals  $I_{w_-,0}$ , we see that the two points  $x_{\text{Inv}(w_+)}, x_{\text{Inv}(w'_+)}$  belong both to the interval  $I_{(\text{Inv}(w_+))_{-n},0}$  but inside it, they belong to the disjoint sub-intervals  $I_{(\text{Inv}(w_+))_{-n-1},0}$  and  $I_{(\text{Inv}(w'_+))_{-n-1},0}$  respectively. Hence

$$\left| x_{\text{Inv}(w_+)} - x_{\text{Inv}(w'_+)} \right| \asymp \left| I_{(\text{Inv}(w_+))_{-n},0} \right|$$

<sup>7</sup>The notation  $|\Phi(\rho) - \Phi(\rho')| \asymp |\rho - \rho'|$  means precisely that there exist  $C > 0$  such that for every  $\rho, \rho'$ ,  $C^{-1} |\rho - \rho'| \leq |\Phi(\rho) - \Phi(\rho')| \leq C |\rho - \rho'|$ .

uniformly over  $w, w' \in \mathcal{W}$ , where  $|I|$  is the length of the interval  $I$ . From the definition (4.26) of the sets  $\tilde{I}_{w_0, n}$  we observe that the points  $\rho = (x_{w_-}, \xi_w)$  and  $\rho' = (x_{w'_-}, \xi_{w'})$  belong respectively to the sets  $\tilde{I}_{w_0, n}$  and  $\tilde{I}_{w'_0, n}$ . Let  $\tilde{w}' := (w'_-, w_+)$ . We have

$$|\rho - \rho'| = \left| (x_{w_-}, \xi_w) - (x_{w'_-}, \xi_{w'}) \right| \quad (4.42)$$

$$\asymp \left| (x_{w_-}, \xi_w) - (x_{w_-}, \xi_{\tilde{w}'}) \right| + \left| (x_{w_-}, \xi_{\tilde{w}'}) - (x_{w'_-}, \xi_{w'}) \right| \quad (4.43)$$

$$\asymp \left| x_{w_-} - x_{w'_-} \right| + |\xi_w - \xi_{\tilde{w}'}|.$$

The points  $\xi_w, \xi_{\tilde{w}'}$  belong to the same set  $\tilde{I}_{w_0, n}$ . However if assumption of “minimal captivity” holds, they belong to disjoint sub-sets  $\tilde{I}_{w_0, n+1}$  and  $\tilde{I}_{w'_0, n+1}$  respectively. Hence

$$|\xi_w - \xi_{\tilde{w}'}| \asymp |J_{w, n}| \quad (4.44)$$

with the interval  $J_{w, n} := \tilde{I}_{w_0, n} \cap \pi^{-1}(x_{w_-})$ . From the bounded distortion principle [19] we have that

$$\forall x, y \in I_{w_{-n, 0}}, \quad |(D\phi_{w_{-n, 0}})(x)| \asymp |(D\phi_{w_{-n, 0}})(y)| \asymp |I_{w_{-n, 0}}|$$

uniformly with respect to  $w, n, x, y$ . From the expression of the canonical map  $F$  in (4.4) and the bounded distortion principle, we have that

$$|J_{w, n}| \asymp |(D\phi_{w_{-n, 0}})(x)|, \quad \forall x \in I_{w_0},$$

uniformly with respect to  $w, n, x$ . Using the previous results we get

$$\begin{aligned} \left| x_{w_-} - x_{w'_-} \right| + |\xi_w - \xi_{w'}| &\asymp \left| x_{w_-} - x_{w'_-} \right| + |\xi_w - \xi_{\tilde{w}'}| \\ &\asymp \left| x_{w_-} - x_{w'_-} \right| + |J_{w, n}| \\ &\asymp \left| x_{w_-} - x_{w'_-} \right| + (D\phi_{w_0, n})(x), \quad \forall x \in I_{w_0}, \\ &\asymp \left| x_{w_-} - x_{w'_-} \right| + |I_{\text{Inv}(w_0, n)}| \\ &\asymp \left| x_{w_-} - x_{w'_-} \right| + \left| x_{\text{Inv}(w_+)} - x_{\text{Inv}(w'_+)} \right|. \end{aligned}$$

We have obtained (4.41) and finished the proof of Lemma 4.17 and Proposition 4.16.  $\square$

## 5 Proof of Theorem 2.9 for the spectral gap in the semi-classical limit

For the proof of Theorem 2.9, we will follow step by step the same analysis as in Section 3 (and also follow closely the proof of Theorem 2 in [21]). The main difference now is that  $\hbar \ll 1$  is a semi-classical parameter (not fixed anymore). In other words, we just

perform a linear rescaling in cotangent space:  $\xi_h := \hbar\xi$ . Our quantization rule for a symbol  $A(x, \xi_h) \in S^{-m}(\mathbb{R})$ , Eq.(3.14) writes now (see [37] p.22), for  $\varphi \in \mathcal{S}(\mathbb{R})$ :

$$\left(\hat{A}\varphi\right)(x) := \frac{1}{2\pi\hbar} \int A(x, \xi_h) e^{i(x-y)\xi_h/\hbar} \varphi(y) dy d\xi_h. \quad (5.1)$$

For simplicity we will still write  $\xi$  instead of  $\xi_h$  below.

## 5.1 The escape function

Let  $1 < \kappa < 1/\theta$  and  $R > 0$  given in Lemma 4.4. Let  $m > 0$ ,  $\eta > 0$  (small) and consider a  $C^\infty$  function  $A_m(x, \xi)$  on  $T^*\mathbb{R}$  so that:

$$\begin{aligned} A_m(x, \xi) &:= \langle \xi \rangle^{-m} && \text{for } |\xi| > R + \eta \\ &:= 1 && \text{for } \xi \leq R \end{aligned}$$

where  $\langle \xi \rangle := (1 + \xi^2)^{1/2}$ .  $A_m$  belongs to the symbol class  $S^{-m}(\mathbb{R})$  defined in (3.12).

From Eq. (4.6) we can deduce, similarly to Eq.(3.13) and if  $\eta$  is small enough, that:

$$\forall x \in I, \forall |\xi| > R, \forall i \rightsquigarrow j \quad \frac{A_m(F_{i,j}(x, \xi))}{A_m(x, \xi)} \leq C^m < 1, \quad \text{with } C = \sqrt{\frac{R^2 + 1}{\kappa^2 R^2 + 1}} < 1. \quad (5.2)$$

This means that the function  $A_m$  is an **escape function** since it decreases strictly along the trajectories of  $F$  outside the zone  $\mathcal{Z}_0 := I \times [-R, R]$ . For any point  $(x, \xi) \in T^*I$  we have the more general bound:

$$\forall x \in I, \forall \xi \in \mathbb{R}, \forall i \rightsquigarrow j \quad \frac{A_m(F_{i,j}(x, \xi))}{A_m(x, \xi)} \leq 1. \quad (5.3)$$

Let  $\hbar > 0$ . Using the quantization rule (5.1), the symbol  $A_m$  can be quantized giving a  $\hbar$ -pseudodifferential operator  $\hat{A}_m$  which is self-adjoint and invertible on  $C^\infty(I)$ . In our case  $\hat{A}_m$  is simply a multiplication operator by  $A_m(\xi)$  in  $\hbar$ -Fourier space.

## 5.2 Using the Egorov Theorem

Let us consider the Sobolev space

$$H^{-m}(\mathbb{R}) := \hat{A}_m^{-1}(L^2(\mathbb{R}))$$

which is the usual Sobolev space as a linear space, except for the norm which depends on  $\hbar$ . Then  $\hat{F} : H^{-m}(\mathbb{R}) \rightarrow H^{-m}(\mathbb{R})$  is unitary equivalent to

$$\hat{Q} := \hat{A}_m \hat{F}_{\hbar, \chi} \hat{A}_m^{-1} : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}).$$

Let  $n \in \mathbb{N}^*$  be a fixed time which will be made large at the end of the proof, and define

$$\hat{P}^{(n)} := \hat{Q}^{*n} \hat{Q}^n = \hat{A}_m^{-1} \hat{F}_{h,\chi}^{*n} \hat{A}_m^2 \hat{F}_{h,\chi}^n \hat{A}_m^{-1}. \quad (5.4)$$

From Egorov Theorem, as in Lemma 3.7, we have that  $\hat{B} := \hat{F}_{h,\chi}^{*n} \hat{A}_m^2 \hat{F}_{h,\chi}^n$  is a PDO with principal symbol

$$\begin{aligned} B(x, \xi) &= \chi^2(x) \sum_{j \text{ s.t. } i \rightsquigarrow j} |\phi'_{i,j}(x)| e^{2\text{Re}(V(\phi_{i,j}(x)))} A_m^2(F_{i,j}(x, \xi)), \quad (x, \xi) \in T^*I \\ &= \chi^2(x) \sum_{j \text{ s.t. } i \rightsquigarrow j} e^{2D((\phi_{i,j}(x)))} A_m^2(F_{i,j}(x, \xi)) \end{aligned}$$

where we have used the “damping function”  $D(x) := \text{Re}(V(x)) - \frac{1}{2} \log(|(\phi^{-1})'(x)|)$  already defined in (2.23). Iteratively for every  $n \geq 1$ , Egorov’s Theorem gives that  $(\hat{F}_{h,\chi}^*)^n \hat{A}_m^2 \hat{F}_{h,\chi}^n$  is a PDO with principal symbol

$$B_n(x, \xi) = \chi^2(x) \sum_{w_{-n,0} \in \mathcal{W}_-} e^{2D_{w_{-n,0}}(x)} A_m^2(F_{w_{-n,0}}(x, \xi))$$

where  $\mathcal{W}_+$  is the set of admissible sequences, defined in (4.18), with the Birkhoff sum  $D_{w_{-n,0}}(x) := \sum_{k=1}^n D(\phi_{w_{-n,-k}}(x))$  and

$$F_{w_{-n,0}} := F_{w_{-1}, w_0} \circ \dots \circ F_{w_{-n}, w_{-n+1}}.$$

With the theorem of composition of PDO [57, chap.4] we obtain that  $\hat{P}^{(n)}$  is a PDO of order 0 with principal symbol given by

$$P^{(n)}(x, \xi) = \left( \chi^2(x) \sum_{w_{-n,0} \in \mathcal{W}_-} e^{2D_{w_{-n,0}}(x)} \frac{A_m^2(F_{w_{-n,0}}(x, \xi))}{A_m^2(x, \xi)} \right). \quad (5.5)$$

We define

$$\gamma(n) := \sup_{x \in I, w_{-n,0} \in \mathcal{W}_-} \frac{1}{n} D_{w_{-n,0}}(x)$$

hence  $e^{2D_{w_{-n,0}}(x)} \leq e^{2n\gamma(n)}$ .

From Theorem 2.6, the spectrum of  $\hat{F}_{h,\chi}$  does not depend on the choice of  $\chi$ . Here we take  $a \geq 0$  as given in Assumption 4.7 and we choose  $\chi$  such that  $\chi \equiv 1$  on  $K_{a+1}$ ,  $\chi \equiv 0$  on  $\mathbb{R} \setminus K_a$ . We have  $P(x, \xi) = 0$  if  $x \in \mathbb{R} \setminus K_a$ .

Now we will bound the positive symbol  $P^{(n)}(x, \xi)$  from above, considering  $x \in K_a$  and different possibilities for the trajectory  $F_{w_{-n,0}}(x, \xi)$ :

1. If  $|\xi| > R$ , Eq.(5.2) gives

$$\frac{A_m^2(F_{w_{-n,0}}(x, \xi))}{A_m^2(x, \xi)} = \frac{A_m^2(F_{w_{-n,0}}(x, \xi))}{A_m^2(F_{w_{-n,-1}}(x, \xi))} \frac{A_m^2(F_{w_{-n,-1}}(x, \xi))}{A_m^2(F_{w_{-n,-2}}(x, \xi))} \cdots \frac{A_m^2(F_{w_{-n,-n+1}}(x, \xi))}{A_m^2(x, \xi)} \quad (5.6)$$

$$\leq (C^{2m})^n \quad (5.7)$$

therefore

$$P^{(n)}(x, \xi) \leq (\#\mathcal{W}_n) e^{2n\gamma(n)} (C^{2m})^n \leq (Ne^{2\gamma(n)} C^{2m})^n.$$

We have used that  $\#\mathcal{W}_n \leq N^n$ . Notice that  $C^{2m}$  can be made arbitrarily small if  $m$  is large.

2. If  $|\xi| \leq R$ , we have from the hypothesis of minimal captivity (Assumption 4.7) and Proposition 4.10 that at time  $(n-1)$  every point  $(x', \xi')$  of the set  $F^{n-1}(x, \xi)$  except finitely many points, satisfy  $|\xi'| > R$ . Using (5.3) and (5.2), for all these points one has  $\frac{A_m^2(F_{w_{-n,0}}(x, \xi))}{A_m^2(x, \xi)} \leq C^{2m}$  and for the exceptional point one can only write  $\frac{A_m^2(F_{w_{-n,0}}(x, \xi))}{A_m^2(x, \xi)} \leq 1$ . This gives

$$P^{(n)}(x, \xi) \leq e^{2n\gamma(n)} ((\#\mathcal{W}_n - 1) C^{2m} + C') \leq \mathcal{B}$$

with the bound

$$\mathcal{B} := e^{2n\gamma(n)} (N^n C^{2m} + C'). \quad (5.8)$$

With the  $L^2$  **continuity theorem** for pseudodifferential operators [37, 15] this implies that in the limit  $\hbar \rightarrow 0$

$$\left\| \hat{P}^{(n)} \right\|_{L^2} \leq \mathcal{B} + \mathcal{O}_n(\hbar). \quad (5.9)$$

Polar decomposition of  $\hat{Q}^n$  gives

$$\left\| \hat{Q}^n \right\|_{L^2} \leq \left\| \hat{Q}^n \right\|_{L^2} = \sqrt{\left\| \hat{P}^{(n)} \right\|_{L^2}} \leq (\mathcal{B} + \mathcal{O}_n(\hbar))^{1/2}. \quad (5.10)$$

Let  $\gamma_+ = \limsup_{n \rightarrow \infty} \gamma(n)$ . If we let  $\hbar \rightarrow 0$  first, and  $m \rightarrow +\infty$  giving  $C^{2m} \rightarrow 0$ , and  $n \rightarrow \infty$ , we obtain  $(\mathcal{B} + \mathcal{O}_n(\hbar))^{1/(2n)} \rightarrow e^{\gamma_+}$ . Therefore for any  $\rho > e^{\gamma_+}$ , there exists  $n_0 \in \mathbb{N}$ ,  $\hbar_0 > 0$ ,  $m_0 > 0$  such that for any  $\hbar \leq \hbar_0$ ,  $m > m_0$ ,

$$\left\| \hat{F}_{\hbar, \chi}^{n_0} \right\|_{H^{-m}} = \left\| \hat{Q}^{n_0} \right\|_{L^2} \leq \rho^{n_0}. \quad (5.11)$$

Also, there exists  $c > 0$  independent of  $\hbar \leq \hbar_0$ , such that for any  $r$  such that  $0 \leq r < n_0$  we have  $\left\| \hat{Q}^r \right\|_{L^2} < c\rho^r$ . As a consequence for any  $n \in \mathbb{N}$  we write  $n = kn_0 + r$  with  $0 \leq r < n_0$  and

$$\left\| \hat{F}_{\hbar, \chi}^n \right\|_{H^{-m}} = \left\| \hat{Q}^n \right\|_{L^2} \leq \left\| \hat{Q}^{n_0} \right\|_{L^2}^k \left\| \hat{Q}^r \right\|_{L^2} \leq \rho^n \frac{\left\| \hat{Q}^r \right\|_{L^2}}{\rho^r} \leq c\rho^n.$$



This estimate implies as well the bound on the spectral radius (2.24) as the bound on the resolvent (2.26):

For any  $n$  the spectral radius of  $\hat{Q}$  satisfies [47, p.192]

$$r_s(\hat{Q}) \leq \left\| \hat{Q}^n \right\|^{1/n} \leq c^{1/n} \rho.$$

So we get that for  $\hbar \rightarrow 0$ ,

$$r_s(\hat{F}_{\hbar, \chi}) = r_s(\hat{Q}) \leq e^{\gamma^+} + o(1). \quad (5.12)$$

In order to obtain the bound (2.26) on the resolvent let  $|z| > \rho_2 > \rho$ . The relation  $(z - \hat{F}_{\hbar, \chi})^{-1} = z^{-1} \sum_{n \geq 0} \left( \frac{\hat{F}_{\hbar, \chi}}{z} \right)^n$  gives that

$$\left\| (z - \hat{F}_{\hbar, \chi})^{-1} \right\|_{H^{-m}} \leq |z|^{-1} \sum_{n \geq 0} \frac{\left\| \hat{F}_{\hbar, \chi}^n \right\|_{H^{-m}}}{|z|^n} \leq |z|^{-1} c_{\rho_1} \sum_{n \geq 0} \frac{\rho_1^n}{|z|^n} = \frac{c_{\rho_1}}{|z| - \rho_1} \leq \frac{c_{\rho_1}}{\rho_2 - \rho_1} =: C_{\rho_2}$$

which finishes the proof of Theorem 2.9.

## 6 Proof of Theorem 2.16 about the fractal Weyl law

We will prove this result once more by conjugating the transfer operator by an escape function as in previous Section 5. However we need first to improve the properties of the escape function. The fractal Weyl estimate will then follow from general trace estimates of PDOs and general lemmas on singular values of compact operators which we recall in the Appendices.

### 6.1 A refined escape function

#### 6.1.1 Distance function

The escape function  $A$  will be constructed from a distance function  $\delta$ . For  $x \in I$ , let

$$\tilde{K}(x) := \tilde{K} \cap (\{x\} \times \mathbb{R}) \quad (6.1)$$

where  $\tilde{K}$  has been defined in (4.10). With this notation we can define the following distance function.

**Definition 6.1.** Let  $x \in I_{w_0}$  and  $\xi \in \mathbb{R}$ , we define the distance of  $(x, \xi)$  to the set  $\tilde{K}$  given in (4.10) by

$$\delta(x, \xi) := \text{dist}(\xi, \tilde{K}(x)) = \min_{w \in \mathcal{W}_+} |\xi - \zeta_w(x)|. \quad (6.2)$$

We will show that the distance function  $\delta(x, \xi)$  decreases along the trajectories of  $F$ . First, the next Lemma shows how the branches  $\zeta_w$  are transformed under the canonical map  $F$ . This formula follows from straightforward calculations.

**Lemma 6.2.** *For every  $w_+ = (w_0, w_1, \dots) \in \mathcal{W}_+$ ,  $x \in I_{w_0}$  we have*

$$F_{w_0, w_1}(x, \zeta_{w_+}(x)) = (x', \zeta_{L(w_+)}(x')) \quad (6.3)$$

with  $L(w_+) := (w_1, w_2, \dots)$  and  $x' = \phi_{w_0, w_1}(x)$ .

**Lemma 6.3.**  $\forall i, j, i \rightsquigarrow j, \forall x \in I_i, \forall \xi \in \mathbb{R}$ ,

$$\delta(F_{i,j}(x, \xi)) \geq \frac{1}{\theta} \delta(x, \xi)$$

where  $\theta < 1$  is given by (2.1).

*Proof.* Let  $i = w_0 \rightsquigarrow j = w_1$ ,  $x \in I_{w_0}$ . Let  $(x', \xi') := F_{w_0, w_1}(x, \xi)$  with  $x' \in I_{w_1}$ . We use (6.3) and also that  $F_{w_0, w_1}$  is expansive in  $\xi$  by a factor larger than  $\theta^{-1} > 1$  (Eq.(4.4)), and get

$$|\xi' - \zeta_{L(w_+)}(x')| = \left| (F_{w_0, w_1}(x, \xi) - F_{w_0, w_1}(x, \zeta_{w_+}(x))) \right|_{\xi} \geq \frac{1}{\theta} |\xi - \zeta_{w_+}(x)|.$$

Thus

$$\begin{aligned} \delta(F_{w_0, w_1}(x, \xi)) &= \min_{w \in \mathcal{W}_+} |\xi' - \zeta_w(x')| = \min_{w_+ \in \mathcal{W}_+} |\xi' - \zeta_{L(w_+)}(x')| \\ &\geq \frac{1}{\theta} \min_{w_+ \in \mathcal{W}_+} |\xi - \zeta_{w_+}(x)| = \frac{1}{\theta} \delta(x, \xi). \end{aligned}$$

□

### 6.1.2 Escape function

The aim of this section is to prove the existence of an escape function with the following properties:

**Proposition 6.4.**  $\forall 1 < \kappa < \theta^{-1}, \exists C_0 > 0$ , s.t.  $\forall \mu, 0 \leq \mu < \frac{1}{2}, \forall m > 0$ , there exists an  $\hbar$ -dependent order function  $A_{m,\mu} \in \mathcal{O}\mathcal{F}^{m\mu}(\langle \xi \rangle^{-m})$  (as defined in Definition B.3) which fulfills the following “decay condition”:

$\forall i, j$ , s.t.  $i \rightsquigarrow j$  and  $\forall (x, \xi) \in I_i \times \mathbb{R}$  s.t.  $\delta(x, \xi) > C_0 \hbar^\mu$  the following estimate holds:

$$\left( \frac{A_{m,\mu} \circ F_{i,j}}{A_{m,\mu}} \right) (x, \xi) \leq \kappa^{-m}. \quad (6.4)$$

In order to prove the above proposition we first remark that the distance function (6.2) is not differentiable, however Lipschitz.

**Lemma 6.5.** Let  $C_1 := \sup_{x \in I, \omega \in \mathcal{W}_+} |(\partial_x \zeta_\omega)(x)|$ . Then  $\delta : T^*I \rightarrow \mathbb{R}^+$  is a Lipschitz function with constant  $C_1 + 1$

*Proof.* Let  $x, y \in I_i$ , then from the fact, that  $|(\partial_x \zeta_\omega)(x)|$  is uniformly bounded by  $C_1$  we have

$$|\delta(x, \xi) - \delta(y, \xi)| \leq C_1 |x - y|.$$

On the other hand clearly

$$|\delta(y, \xi) - \delta(y, \zeta)| \leq |\xi - \zeta|$$

thus

$$|\delta(x, \xi) - \delta(y, \zeta)| \leq C_1 |x - y| + |\xi - \zeta| \leq (C_1 + 1) \text{dist}((x, \xi), (y, \zeta)).$$

□

Next we choose  $0 \leq \mu < 1/2$  and regularize the function  $\delta$  at the scale  $\hbar^\mu$ . For this we choose  $\chi \in C_0^\infty(\mathbb{R}^2)$  with support in the unit ball  $B_1(0)$  of  $\mathbb{R}^2$  and  $\chi(x, \xi) > 0$  for  $\|(x, \xi)\| < 1$ . This function can be rescaled to

$$\chi_{\hbar^\mu}(x, \xi) := \frac{1}{\hbar^{2\mu} \|\chi\|_{L^1}} \chi\left(\frac{x}{\hbar^\mu}, \frac{\xi}{\hbar^\mu}\right)$$

such that  $\text{supp} \chi_{\hbar^\mu} \subset B_{\hbar^\mu}(0)$  and  $\int \chi_{\hbar^\mu}(x) dx = 1$ . Now we can define the regularized distance function by

$$\tilde{\delta}(x, \xi) := \int_{T^*I} \delta(x', \xi') \chi_{\hbar^\mu}(x - x', \xi - \xi') dx' d\xi'.$$

This smoothed distance function  $\tilde{\delta}$  differs only at order  $\hbar^\mu$  from the original one because

$$\begin{aligned} \left| \tilde{\delta}(x, \xi) - \delta(x, \xi) \right| &= \left| \int_{\mathbb{R}^2} (\delta(x, \xi) - \delta(x - x', \xi - \xi')) \chi_{\hbar^\mu}(x', \xi') dx' d\xi' \right| \\ &\leq \sup_{(x', \xi') \in B_{\hbar^\mu}(0)} |(\delta(x, \xi) - \delta(x - x', \xi - \xi'))| \\ &\leq (C_1 + 1) \hbar^\mu. \end{aligned} \quad (6.5)$$

Furthermore we get the following estimates for its derivatives:

**Lemma 6.6.** *For all  $\alpha, \beta \in \mathbb{N}$  the estimate*

$$|\partial_x^\alpha \partial_\xi^\beta \tilde{\delta}(x, \xi)| \leq C_{\alpha, \beta} \hbar^{-\mu(\alpha+\beta)} (\delta(x, \xi) + C \hbar^\mu)$$

*holds*

*Proof.* From the definition of  $\chi_{\hbar^\mu}$  we have  $\|\partial_x^\alpha \partial_\xi^\beta \chi_{\hbar^\mu}\|_\infty \leq C_{\alpha, \beta} \hbar^{-2-(\alpha+\beta)\mu}$  and thus:

$$\begin{aligned} \left| \left( \partial_x^\alpha \partial_\xi^\beta \tilde{\delta}(x, \xi) \right) \right| &= \int_{T^*I} \delta(x', \xi') \partial_x^\alpha \partial_\xi^\beta \chi_{\hbar^\mu}(x - x', \xi - \xi') dx' d\xi' \\ &\leq \pi \hbar^{2\mu} \|\delta\|_{\infty, B_{\hbar^\mu}(x, \xi)} C_{\alpha, \beta} \hbar^{-(2+\alpha+\beta)\mu} \\ &\leq \pi C_{\alpha, \beta} \hbar^{-(\alpha+\beta)\mu} (\delta(x, \xi) + (C_1 + 1) \hbar^\mu) \end{aligned}$$

where we used the Lipschitz property of  $\delta$  (Lemma 6.5) in the last inequality.  $\square$

As  $|\delta(x, \xi)| \leq |\xi| + C$  the above lemma gives us directly that  $\tilde{\delta} \in S_\mu^1(T^*I)$ . Now we define the escape function as:

$$A_{m, \mu}(x, \xi) := \hbar^{m\mu} \left( \hbar^{2\mu} + \left( \tilde{\delta}(x, \xi) \right)^2 \right)^{-\frac{m}{2}}. \quad (6.6)$$

This is obviously a smooth function and it fulfills the conditions of an  $\hbar$ -dependent order function (c.f. Definition B.3)

**Lemma 6.7.** *The function  $A_{m, \mu}$  defined in (6.6) is an  $\hbar$ -dependent order function  $A_{m, \mu} \in \mathcal{O}\mathcal{F}^{m\mu}(\langle \xi \rangle^{-m})$  as defined in Definition B.3.*

*Proof.* As  $\tilde{K} \subset I \times [-R, R]$  we obtain  $\min(0, |\xi| - R) \leq \tilde{\delta}(x, \xi) \leq |\xi| + R$ . This implies that  $A_{m, \mu}(x, \xi) \leq \tilde{C} \langle \xi \rangle^{-m}$  and that  $A_{m, \mu}(x, \xi) \geq C' \hbar^{m\mu} \langle \xi \rangle^{-m}$ . It remains thus to show, that for arbitrary  $\alpha, \beta \in \mathbb{N}$  one has:

$$|\partial_x^\alpha \partial_\xi^\beta A_{m, \mu}(x, \xi)| \leq C_{\alpha, \beta} \hbar^{-\mu(\alpha+\beta)} A_{m, \mu}(x, \xi) \quad (6.7)$$

where  $C_{\alpha,\beta}$  depends only on  $\alpha$  and  $\beta$ . First consider the case  $\alpha = 1, \beta = 0$ :

$$|\partial_x A_{m,\mu}(x, \xi)| = \left| \hbar^{m\mu} m \frac{(\partial_x \tilde{\delta}(x, \xi)) \tilde{\delta}(x, \xi)}{\left( \hbar^{2\mu} + \left( \tilde{\delta}(x, \xi) \right)^2 \right)^{\frac{m+2}{2}}} \right| \leq C \hbar^{-\mu} A_{m,\mu}(x, \xi)$$

where we used  $\tilde{\delta} \leq \sqrt{\hbar^{2\mu} + \tilde{\delta}^2}$  and  $|\partial_x \tilde{\delta}| \leq C \hbar^{-\mu} \sqrt{\hbar^{2\mu} + \tilde{\delta}^2}$  which follows from Lemma 6.6 together with (6.5). Inductively one obtains the estimate for arbitrary  $\alpha, \beta \in \mathbb{N}$  by repeated use of Lemma 6.6 and (6.5).  $\square$

Finally it remains to show the decay estimates for  $\left( \frac{A_{m,\mu} \circ F_{i,j}}{A_{m,\mu}} \right)(x, \xi)$ . Combining (6.5) with Lemma 6.3 we then get

$$\begin{aligned} \tilde{\delta}(F_{i,j}(x, \xi)) &\geq \delta(F_{i,j}(x, \xi)) - (C_1 + 1) \hbar^\mu \geq \frac{1}{\theta} \delta(x, \xi) - (C_1 + 1) \hbar^\mu \\ &\geq \frac{1}{\theta} \tilde{\delta}(x, \xi) - \left( \frac{1}{\theta} + 1 \right) (C_1 + 1) \hbar^\mu \end{aligned}$$

and thus

$$\frac{A_{m,\mu}(F_{i,j}(x, \xi))}{A_{m,\mu}(x, \xi)} \leq \left( \frac{1 + \left( \frac{1}{\theta} \cdot \frac{\tilde{\delta}(x, \xi)}{\hbar^\mu} - \tilde{C} \right)^2}{1 + \left( \frac{\tilde{\delta}(x, \xi)}{\hbar^\mu} \right)^2} \right)^{\frac{m}{2}} \quad (6.8)$$

where  $\tilde{C} = \left( \frac{1}{\theta} + 1 \right) (C_1 + 1)$ . Clearly the right side of (6.8) converges to  $\left( \frac{1}{\theta} \right)^{-m}$  for  $\frac{\tilde{\delta}(x, \xi)}{\hbar^\mu} \rightarrow \infty$  which proves the existence of a desired  $C_0$  and finishes the proof of Proposition 6.4.

### 6.1.3 Truncation in $x$

Here we choose a similar truncation operator  $\hat{\chi}$  as in Eq.(2.14) but in a finer vicinity of the trapped set  $K$ . First notice that  $K_{\hbar^\mu} \Subset \phi^{-1}(K_{\hbar^\mu})$  where  $K_{\hbar^\mu}$  has been defined in Definition 2.13. For  $\hbar$  small enough we have  $\phi^{-1}(K_{\hbar^\mu}) \Subset I$ . Let  $\chi \in C_{\phi^{-1}(K_{\hbar^\mu})}^\infty$  such that  $\chi(x) = 1$  for  $x \in K_{\hbar^\mu}$ .  $\chi$  can be considered as a function  $\chi(x, \xi) := \chi(x)$  (independent of  $\xi$ ) and we have that  $\chi_\mu \in S_\mu^0(T^*\mathbb{R})$ . As in Eq.(2.14) we define  $\hat{\chi} := \text{Op}_\hbar^w(\chi)$  which is the multiplication operator by  $\chi$  and

$$F_{\hbar,\chi} := \hat{F}_\hbar \hat{\chi}.$$

## 6.2 Weyl law

The Weyl law will give an upper bound on the number of eigenvalues of  $\hat{F}_{\hbar,\chi}$  in the Sobolev spaces  $H^m$ . These estimates will be obtained by conjugating  $\hat{F}_{\hbar,\chi}$  with  $\text{Op}_\hbar^w(A_{m,\mu})$  in the same way as for the discrete spectrum or the spectral gap. Note that we use the Weyl quantization (see Definition B.2) in this section, because we want to obtain self adjoint operators. In order to be able to conjugate we have to show, that  $\text{Op}_\hbar^w(A_{m,\mu}) : H^{-m} \rightarrow L^2$

is an isomorphism. We already know that  $Op_h^w(\langle \xi \rangle^m) : L^2 \rightarrow H^{-m}$  is an isomorphism, thus it suffices to show, that  $\hat{B} := Op_h^w(A_{m,\mu})Op_h^w(\langle \xi \rangle^m) : L^2 \rightarrow L^2$  is invertible. From the  $\hbar$ -local symbol calculus (Theorem B.7) it follows that  $\hat{B}$  is an elliptic operator in the  $\hbar$ -local symbol class  $S_\mu(A_{m,\mu}\langle \xi \rangle^m)$  and thus the invertibility follows from Proposition B.10. Note that it is necessary to work in the  $\hbar$ -local symbol classes as  $\hat{B}$  would not be an elliptic operator in  $S_\mu(1)$ . Proposition B.10 also gives us the leading order of our inverse  $\hat{B}^{-1}$  which is  $A_{m,\mu}^{-1}\langle \xi \rangle^{-m}$ . So the inverse of  $Op_h^w(A_{m,\mu})$  is again a PDO with leading symbol  $A_{m,\mu}^{-1}$ .

With the isomorphism  $Op_h^w(A_{m,\mu}) : H^{-m} \rightarrow L^2$  we can thus define a different scalar product on the Sobolev spaces which turns  $Op_h^w(A_{m,\mu})$  into a unitary operator. The Sobolev space equipped with this scalar product will be denoted by  $\mathcal{H}_{\hbar,\mu}^{-m}$  and the study of  $\hat{F}_\hbar$  is thus unitary equivalent to the study of  $\hat{Q}_m$  defined by the following commutative diagram (where we noted  $\hat{A}_{m,\mu} := Op_h^w(A_{m,\mu})$ ):

$$\begin{array}{ccc} L^2(\mathbb{R}) & \xrightarrow{\hat{Q}_m} & L^2(\mathbb{R}) \\ \downarrow \hat{A}_{m,\mu}^{-1} & & \downarrow \hat{A}_{m,\mu}^{-1} \\ \mathcal{H}_{\hbar,\mu}^{-m} & \xrightarrow{\hat{F}_{\hbar,\chi}} & \mathcal{H}_{\hbar,\mu}^{-m} \end{array} \quad (6.9)$$

In the next Lemma,  $C_0$  and  $\kappa$  are as in Lemma 6.4.

**Lemma 6.8.** *Let  $C_0$  be as in Lemma 6.4. Then for every  $\epsilon > 0$  and  $0 \leq \mu < \frac{1}{2}$  there is  $m_0 > 0$  and  $\tilde{C}_1, \tilde{C}_2 > 0$  such that for all  $m > m_0$  and in the limit  $\hbar \rightarrow 0$  we have:*

$$\#\left\{ \lambda_i^\hbar \in \sigma\left(\hat{F}_{\hbar,\chi}|_{\mathcal{H}_{\hbar,\mu}^m}\right) \mid |\lambda_i^\hbar| \geq \epsilon \right\} \leq \frac{1}{2\pi\hbar} \left( \tilde{C}_1 \text{Leb}\{\mathcal{K}_{C_0\hbar^\mu}\} + \tilde{C}_2\hbar \right). \quad (6.10)$$

Before proving Lemma 6.8, let us show that it implies Theorem 2.16. From Theorem 2.6, the discrete spectrum of  $\hat{F}_{\hbar,\chi}|_{\mathcal{H}_{\hbar,\mu}^m}$  is the Ruelle spectrum of resonances  $\text{Res}\left(\hat{F}_\hbar\right)$ , independent of  $\mu$  and  $m$ . With Assumption 4.7 we can use Eq.(4.36) and that  $\mathcal{K}$  has pure dimension, thus equation (2.32) gives  $\text{Leb}\{\mathcal{K}_{C_1\hbar^\mu}\} = \mathcal{O}\left((\hbar^\mu)^{\text{codim}_M(\mathcal{K})}\right)$ . As  $\text{codim}_M(\mathcal{K}) < 2$  and  $\mu < \frac{1}{2}$ , equation(6.10) gives

$$\begin{aligned} \#\left\{ \lambda_i^\hbar \in \text{Res}\left(\hat{F}_\hbar\right) \mid |\lambda_i^\hbar| \geq \epsilon \right\} &= \mathcal{O}\left(\hbar^{-1}(\hbar^\mu)^{\text{codim}_M(\mathcal{K})}\right) \\ &= \mathcal{O}\left(\hbar^{-1}(\hbar^\mu)^{2-2\dim_H(K)}\right) = \mathcal{O}\left(\hbar^{2\mu-1-2\mu\dim_H(K)}\right) \end{aligned}$$

for any fixed  $0 \leq \mu < 1/2$ . This gives Theorem 2.16 with  $\eta = (1 - 2\mu)(1 - \dim_H(K))$ .

*Proof. of Lemma 6.8.* From (6.9),  $\hat{F}_{\hbar,\chi} : \mathcal{H}_{\hbar,\mu}^{-m} \rightarrow \mathcal{H}_{\hbar,\mu}^{-m}$  is unitary equivalent to

$$\hat{Q}_{m,\mu} := Op_h^w(A_{m,\mu})\hat{F}_{\hbar,\chi}Op_h^w(A_{m,\mu})^{-1} : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}).$$

Consider

$$\hat{P}_\mu := \hat{Q}_{m,\mu}^* \hat{Q}_{m,\mu} = \text{Op}_\hbar^w(A_{m,\mu})^{-1} \hat{\chi} \hat{F}_\hbar^* \text{Op}_\hbar^w(A_{m,\mu})^2 \hat{F}_\hbar \hat{\chi} \text{Op}_\hbar^w(A_{m,\mu})^{-1}.$$

By the composition Theorem B.7 and the Egorov theorem B.11 for  $\hbar$ -local symbols,  $\hat{P}_\mu$  is a PDO with leading symbol  $P_\mu(x, \xi) \in S_\mu^0$ . For  $x \in I_i$ ,  $\xi \in \mathbb{R}$  the leading symbol is given by the same expression as in (3.19):<sup>8</sup>

$$P_\mu(x, \xi) = \chi^2(x) \sum_{j \text{ s.t. } i \rightsquigarrow j} |\phi'_{i,j}(x)| e^{2\text{Re}(V(\phi_{i,j}(x)))} \frac{A_{m,\mu}^2(F_{i,j}(x, \xi))}{A_{m,\mu}^2(x, \xi)} \text{ mod } \hbar^{1-2\mu} S_\mu^{-1}(T^*\mathbb{R}). \quad (6.11)$$

Now using the definition of  $\chi$ , Eq.(6.4) and Lemma 6.4, the operator  $\hat{P}_\mu$  can be decomposed into self-adjoint operators

$$\hat{P}_\mu = \hat{k}_\mu + \hat{r}_\mu$$

where  $\hat{k}_\mu$  is a PDO with symbol  $k_\mu \in S_\mu^{-\infty}$  supported on  $\mathcal{K}_{C_0\hbar^\mu}$  for  $C_0$  being the constant from Lemma 6.4. Hence  $\hat{k}_\mu$  is a trace-class operator. The operator  $\hat{r}_\mu$  is a PDO with symbol  $r_\mu \in S_\mu^0$  such that

$$\|r_\mu\|_\infty \leq \theta e^{2\|\text{Re}(V)\|_\infty} \kappa^{-2m} + \mathcal{O}(\hbar^{1-2\mu}),$$

hence  $\|\hat{r}_\mu\| \leq C\kappa^{-2m} + \mathcal{O}(\hbar^{1-2\mu})$ . Here  $\kappa < 1$  is the constant from Lemma 6.4.

Using Lemma C.1 in Appendix C we have that for every  $\epsilon > 0$ , in the limit  $\hbar \rightarrow 0$ ,

$$\#\left\{ \mu_i^\hbar \in \sigma(\hat{k}_\mu) \mid |\mu_i^\hbar| \geq \epsilon \right\} \leq (2\pi\hbar)^{-1} \left( \tilde{C}_1 \text{Leb}\{\mathcal{K}_{C_0\hbar^\mu}\} + \tilde{C}_2\hbar \right). \quad (6.12)$$

By a standard perturbation argument the same estimates holds for the operator  $\hat{P}_\mu$  (for  $m$  sufficiently large): for every  $\epsilon > 0$ , in the limit  $\hbar \rightarrow 0$ ,

$$\#\left\{ \mu_i^\hbar \in \sigma(\hat{P}_\mu) \mid |\mu_i^\hbar| \geq \epsilon + \|\hat{r}_\mu\| \right\} \leq (2\pi\hbar)^{-1} \left( \tilde{C}_1 \text{Leb}\{\mathcal{K}_{C_0\hbar^\mu}\} + \tilde{C}_2\hbar \right). \quad (6.13)$$

From the definition  $\hat{P}_\mu := \hat{Q}_{m,\mu}^* \hat{Q}_{m,\mu}$ , the  $\sqrt{\mu_i^\hbar}$  are singular values of  $\hat{Q}_{m,\mu}$ . Then corollary A.2 from Appendix A shows that the same estimate holds true for the eigenvalues of  $\hat{Q}_{m,\mu}$ , hence of  $\hat{F}_{\hbar,\chi}$ , yielding the result (6.10).  $\square$

## 7 Numerical results for the truncated Gauss map and Bowen-Series maps

In this section we will present numerical results for two important classes of I.F.S.: the truncated Gauss map and the Bowen-Series maps for convex co-compact hyperbolic surfaces. We will show that both examples satisfy the partially captive property. We will

---

<sup>8</sup>Also for this calculation it is crucial to work with the  $\hbar$ -local calculus in order to obtain sufficient remainder estimates.




Figure 7.1: The trapped set  $\mathcal{K}_N := \mathcal{K}$  for the truncated Gauss map with functions (7.1), for the cases of  $N = 3$  and  $N = 10$  branches. This corresponds to the Gauss-Kuzmin-Wirsing transfer operator (7.2). We have  $\mathcal{K}_N \subset \mathcal{K}_{N+1}$  and for  $N \rightarrow \infty$ , the limit trapped set  $\mathcal{K}_\infty = \bigcup_{N \geq 0} \mathcal{K}_N = \{(x, \xi), x \in ]0, 1[, -\frac{2}{1+x} < \xi < 0\}$  is the band between the marked black lines. (More precisely, we have represented the periodic points with period  $n = 6$ . That explains the sparse aspect of the trapped set).

then give some numerical illustrations of the main theorems presented in this paper and finally discuss the connection between the spectrum of these transfer operators with the resonance spectrum of the Laplacian on hyperbolic surfaces.

## 7.1 The truncated Gauss map

In this section we consider the I.F.S. defined from the truncated Gauss map with  $N$  intervals presented in Example 2.2. We choose the roof function  $\tau$  and the potential function  $V$  which enter in the definition of the transfer operator (2.12) to be:

$$\tau(x) = -J(x), \quad V(x) = (1-a)J(x), \quad a \in \mathbb{R}. \quad (7.1)$$

where  $J(x) = \log(|(\phi^{-1})'(x)|) = \log(|G'(x)|) = -2\log(x)$  has been defined in (2.20). Let us write

$$s = a + ib \in \mathbb{C}, \quad b = \frac{1}{\hbar} > 0.$$

Then for every  $s \in \mathbb{C}$ , the transfer operator  $\hat{F}$  given in (2.12) will be written  $\hat{L}_s = \hat{F}$  and is given by:

$$\hat{L}_s \varphi = \hat{F} \varphi = e^{V(x)} e^{i\frac{1}{\hbar}\tau(x)} \varphi \circ \phi^{-1} = e^{(1-s)J} \varphi \circ \phi^{-1} \quad (7.2)$$

As explained in Section 7.1.1 below, this choice is interesting due its relation with the dynamics on the modular surface. The (adjoint of the) transfer operator  $\hat{F}$  constructed in this way is usually called the **Gauss-Kuzmin-Wirsing transfer operator** or “Dieter-Mayer transfer operator” for the truncated Gauss map [39, 55].

**Proposition 7.1.** *For every  $N \geq 1$ , the minimal captivity assumption 4.7 holds true for the truncated Gauss transfer operator defined by (7.2).*






Figure 7.2: The discrete spectrum of Ruelle resonances  $\lambda_j$  (in log scale writing:  $\log \lambda = x + iy$ ) for the truncated Gauss-Kuzmin-Wirsing transfer operator (7.2) associated to the Gauss map, for  $N = 3$  branches and parameters  $a = 1$ ,  $b = 1/\hbar = 0, 100, 1000$ . The full vertical line is at  $x = \text{Pr}(-J) \simeq -0.4$ . For  $b = 0$  there is the eigenvalue  $\lambda = e^{\text{Pr}(-J)}$  plotted at  $(x, y) = (\text{Pr}(-J), -\pi)$ , corresponding to the “equilibrium measure”. The dashed vertical line is at  $x = \gamma_+$  which is shown in (2.24) to be an asymptotic upper bound for  $b = 1/\hbar \rightarrow \infty$ . In this example it seems to be not optimal.

The proof is given in Section 7.3 below. In this proof we explain the structure of the trapped set  $\mathcal{K}$  with more details.

Consequently, we can apply Theorem 2.9 and deduce that there is an asymptotic spectral gap. In Figure 7.2 we present the numerical Ruelle resonances of the truncated Gauss map with 3 branches for different values of  $\hbar$  and compare them with the prediction of the spectral gap. For the numerical calculation we directly use the conjugated transfer operator  $\hat{Q}_m$  that appears in the proofs of the main theorems and develop it in a Fourier basis (see [23, Section 7] for more details on the numerical calculation of Ruelle resonances via the semiclassical approach). One observes on Figure 7.2 that the asymptotic spectral gap given by  $\gamma_+$  is smaller than the general topological pressure bound  $\text{Pr}(-J)$ , Eq.(2.22). The numerical results indicate however, that this gap  $\gamma_+$  is still not optimal.

We can also apply Theorem 2.16 and deduce a fractal Weyl upper bound for the density of resonances. In Figure 7.3 we determine the behavior of the counting function in dependence of the semiclassical parameter in a double logarithmic plot. In these numerical results we observe indeed an algebraic dependence and the exponent agrees very well with the upper bound of the Hausdorff dimension from Theorem 2.16. Note that this is particularly interesting because it is an important open conjecture that the fractal Weyl upper bounds are sharp (see e.g. [41, Section 6] for a review and further references). This conjecture has been supported by numerical experiments in different contexts, like quantum  $n$ -disk systems [36] or convex co-compact hyperbolic surfaces [8]. The data presented in Figure 7.3 is another support for the general validity of the fractal Weyl law.




Figure 7.3: This is the Weyl law for the model of Gauss map with  $N = 3$  branches. The points represent the number of resonances  $\mathcal{N}(b) = \#\left\{\lambda_j \in \text{Res}\left(\hat{L}_s\right), \log|\lambda_j| > -3.5\right\}$  computed numerically, as a function of the semiclassical parameter  $b = 1/\hbar$  in log scale. The linear fit gives  $\log \mathcal{N}(b) = -0.70 \cdot \log b - 0.96$  which has to be compared to the fractal Weyl law (2.34) giving  $\log \mathcal{N}(b) \leq -\dim_H(K) \cdot \log b + \text{cste}$ . From (2.3) we have  $\dim_H K_3 = 0.705$  giving an excellent agreement with the numerical results and suggesting that the upper bound is in fact optimal.

### 7.1.1 Relation with the zeroes of the Selberg zeta function

For the geodesic flow on the modular surface  $\text{SL}_2\mathbb{Z}\backslash\text{SL}_2\mathbb{R}$  it is possible to define the Selberg zeta function (see Section 7.2.1 below for more comments and references):

$$\zeta_{\text{Selberg}}(s) = \prod_{\gamma} \prod_{m \geq 0} (1 - e^{-(s+m)|\gamma|}), \quad s \in \mathbb{C},$$

where the product is over the primitives periodic orbits  $\gamma$  of the geodesic flow and  $|\gamma|$  denotes the length of the orbit. This zeta function is absolutely convergent for  $\text{Re}(s) > 1$ . Using the Gauss map and continued fractions, R. Bowen and C. Series [10]<sup>9</sup> have shown that a periodic orbit  $\gamma$  is in one to one correspondence with a periodic sequence  $(w_j)_{j \in \mathbb{Z}} \in (\mathbb{N} \setminus \{0\})^{\mathbb{Z}}$  where  $w_j \in \mathbb{N} \setminus \{0\}$  is the index of the branch of the Gauss map  $G_{w_j}^{-1}$  in (2.4). Given  $N \geq 1$ , we can restrict the product  $\prod_{\gamma}$  over periodic orbits above to orbits for which  $w_j \leq N, \forall j \in \mathbb{Z}$ , and define a truncated Selberg zeta function as follows:

$$\zeta_{\text{Selberg},N}(s) = \prod_{\gamma, w_j \leq N, \forall j} \prod_{m \geq 0} (1 - e^{-(s+m)|\gamma|}), \quad s \in \mathbb{C}.$$

On the other hand, for fixed  $s \in \mathbb{C}$ , we have from Theorem 2.6 that the operator  $\hat{L}_s$  has discrete spectrum of Ruelle resonances. It is possible to define the dynamical determinant

<sup>9</sup>For the special case of the modular surface and the Gauss map such a correspondence has in deed been known long before see e.g. [1]

of  $\hat{L}_s$  by

$$d(z, s) := \text{Det} \left( 1 - z \hat{L}_s \right) := \exp \left( - \sum_{n \geq 1} \frac{z^n}{n} \text{Tr}^b \left( \hat{L}_s^n \right) \right), \quad z \in \mathbb{C}$$

where  $\text{Tr}^b \left( \hat{L}_s^n \right)$  stands for the flat trace of Atiyah-Bott. The sum is convergent for  $|z|$  small enough. It is known that for fixed  $s$ , the zeroes of  $d(z, s)$  (as a function of  $z$ ) coincide with multiplicities with the Ruelle resonances of  $\hat{L}_s$  [2]. In the case  $z = 1$ , we also have that  $d(1, s)$  coincides with the truncated Selberg zeta function [45][7]:

$$\text{Det} \left( 1 - \hat{L}_s \right) = \zeta_{\text{Selberg}, N}(s) \quad (7.3)$$

which means that the zeroes of  $\zeta_{\text{Selberg}, N}(s)$  are given (with multiplicity) by the event that 1 is a Ruelle resonance of the transfer operator  $\hat{L}_s$ . This also shows that  $\zeta_{\text{Selberg}, N}(s)$  has a holomorphic extension to the complex plane  $s \in \mathbb{C}$ .

*Remark 7.2.* in [45][7, p.306] they consider the adjoint operator  $\hat{L}_s^*$  called the Perron-Frobenius operator.

## 7.2 Bowen-Series maps for Schottky surfaces

The second class of examples that we consider in this section are Bowen-Series maps for Schottky surfaces [10]. We will follow the notation of D. Borthwick's book [7, chap.15] and recall the definition of a Schottky group given there. Recall that an element  $S = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{R})$  acts on  $\mathbb{H}^2 = \text{SL}_2(\mathbb{R})/\text{SO}_2$  and  $\bar{\mathbb{R}} = \partial\mathbb{H}^2$  by  $S(x) := \frac{ax+b}{cx+d}$ .

**Definition 7.3.** Let  $D_1, \dots, D_{2r}$  be disjoint closed half discs in the Poincaré's half plane  $\mathbb{H}^2 = \text{SL}_2(\mathbb{R})/\text{SO}_2$  with center in  $\mathbb{R} = \partial\mathbb{H}^2 \setminus \{\infty\}$ . There exist elements  $S_i \in \text{SL}_2\mathbb{R}$ ,  $i = 1, \dots, r$  such that  $S_i(\partial D_i) = \partial D_{i+r}$  and  $S_i(\text{Int}(D_i)) = \mathbb{C} \setminus D_{i+r}$ . The group generated by the  $S_i$  is called a **Schottky group**  $\Gamma = \langle S_1, \dots, S_r \rangle$ .

*Remark 7.4.* For convenience we will use a cyclic notation for the indices  $i = 1, \dots, 2r$ . Then one can also define  $S_i$  for  $i = r+1, \dots, 2r$  as in the definition above and obtains  $S_{i+r} = S_i^{-1}$ .

Let  $I_i := D_i \cap \partial\mathbb{H}$ . Then  $(I_i)_{i=1, \dots, 2r}$  are  $N = 2r$  disjoint closed intervals. One has  $S_j(\text{Int}(I_j)) = \partial\mathbb{H} \setminus I_{j+r}$  and we assume that  $S_j$  is expanding on  $I_j$  (this can always be obtained by taking iterations if necessary and localizing further to the trapped set, see [7, Prop.15.4]). The maps  $S_j$  are usually called the Bowen-Series maps. Considering the inverse maps one obtains an I.F.S. according to Definition 2.1 associated to this Schottky group in the following way. For any  $j = 1, \dots, N$  and  $i \neq j+r$  let:

$$\phi_{i,j} := S_j^{-1} = S_{j+r} \quad : I_i \rightarrow S_j^{-1}(I_i) \subset \text{Int}(I_j).$$




Figure 7.4: In this (arbitrary) example, we have  $r = 2$  hyperbolic matrices of  $\mathrm{SL}_2\mathbb{R}$ :  $S_1 = \begin{pmatrix} 4 & \sqrt{5} \\ -\sqrt{5} & -1 \end{pmatrix}$  and  $S_2 = \begin{pmatrix} -1 & \sqrt{5} \\ -\sqrt{5} & 4 \end{pmatrix}$  that generate a Schottky group  $\Gamma = \langle S_1, S_2 \rangle$ . Figure (a) shows the Dirichlet fundamental domain  $\mathbb{H}^2 \setminus (D_1 \cup D_2 \cup D_3 \cup D_4)$  with the intervals  $I_i, i = 1, 2, 3, 4$ , on which the I.F.S. is defined. Figure (b) shows the resulting Schottky surface  $\Gamma \setminus \mathbb{H}^2$ . It has three funnels. Figure (c) shows the graph of the generating functions  $\phi_{i,j} = g_j = S_{j+r} : I_i \rightarrow I_j$  of the associated iterated function system.

The adjacency matrix  $A_{i,j}$  has all entries equal to one except  $A_{i,i+r} = 0$  (see Figure 7.4).

As in (7.1) we make the following choice for the potential and the roof function for  $x \in I_j$

$$\tau(x) = -J(x), \quad V(x) = (1-a)J(x), \quad a \in \mathbb{R}. \quad (7.4)$$

where  $J(x) = \log \left( \left| (\phi_{i,j}^{-1})'(x) \right| \right) = \log (|S'_j(x)|)$  has been defined in (2.20). Let us write

$$s = a + ib \in \mathbb{C}, \quad b = \frac{1}{\hbar} > 0.$$

Then for every  $s \in \mathbb{C}$ , the transfer operator  $\hat{F}$  given in (2.12) will be written  $\hat{L}_s = \hat{F}$  and is given by:

$$\hat{L}_s \varphi = \hat{F} \varphi = e^{V(x)} e^{i \frac{1}{\hbar} \tau(x)} \varphi \circ \phi^{-1} = e^{(1-s)J} \varphi \circ \phi^{-1}. \quad (7.5)$$

The adjoint of our transfer operator  $L_s^* = \hat{F}^*$  is exactly the Ruelle transfer operator defined in [7, p.304] and as we will discuss below, its spectrum is in a close connection to the spectrum of the Laplace operator on the Schottky surface.

**Proposition 7.5.** *The minimal captivity assumption 4.7 holds true for the Bowen-Series transfer operator defined by (7.5).*

The proof is given in Section 7.3 below. Consequently, we can apply Theorem 2.9 and deduce that there is an asymptotic spectral gap. We can also apply Theorem 2.16 and deduce an fractal Weyl upper bound for the density of resonances.

*Remark 7.6.* We recall from Section 4.3 that minimal captivity implies NLI condition. F. Naud in [40] has already shown that this weaker NLI condition holds true for Schottky surfaces.

### 7.2.1 Selberg zeta function and resonances of the Laplacian

For the geodesic flow on a hyperbolic surface it is possible to define the Selberg zeta function

$$\zeta_{\text{Selberg}}(s) = \prod_{\gamma} \prod_{m \geq 0} (1 - e^{-(s+m)|\gamma|})$$

where the product is over primitive periodic orbits  $\gamma$  of the geodesic flow and  $|\gamma|$  denotes the length of the orbit. This zeta function is absolutely convergent for  $\text{Re}(s) > 1$  and has a meromorphic continuation to the whole complex plane. This continuation is particularly interesting as its zeros are either “topological zeros” located on the real axis or resonances of the Laplace operator  $\Delta$  on the corresponding hyperbolic surface  $\Gamma \backslash \mathbb{H}^2$ . These resonances  $s \in \text{Res}(\Delta)$  are defined as the poles of the meromorphic extension of the resolvent[7]:

$$R(s) := (\Delta - s(1-s))^{-1}, \quad s \in \mathbb{C}. \quad (7.6)$$

This correspondence follows from the Selberg trace formula for finite-area surfaces, and has been shown by Patterson-Perry [43] for infinite volume surfaces without cusps and Borthwick, Judge and Perry [9] for infinite volume surfaces with cusps (see also [7] for an overview).

For the transfer operators as defined above, one can define a dynamical zeta function by [7, p.305]

$$d(z, s) := \text{Det} \left( 1 - z \hat{L}_s \right).$$

The dynamical and the Selberg zeta function are equal  $\zeta_{\text{Selberg}}(s) = d(1, s)$  (see [7, th.15.8]). This implies immediately that if  $s \in \mathbb{C}$  is a resonance of the Laplacian on the Schottky surface, then 1 has to be an eigenvalue of  $\hat{L}_s$ :

$$s \in \text{Res}(\Delta) \Leftrightarrow 1 \in \text{Spec} \left( \hat{L}_s \right) \quad (7.7)$$

*Remark 7.7.* For the full Gauss map (i.e. with infinitely many branches) the same correspondence between the resonances of the Laplacian on the modular surface  $\text{SL}_2\mathbb{Z}\backslash\mathbb{H}^2$  and the Dieter-Mayer transfer operator  $\hat{L}_s$  is true and has been developed by Dieter Mayer [39]. For the truncated Gauss map considered in Section 7.1, to our knowledge, no such corresponding surfaces are known.

Using the relation (7.7) between the Ruelle spectrum of the transfer operator  $\hat{L}_s$  and the resonances of the Laplacian, it is possible to deduce from Theorem 2.9 some estimate on the ‘‘asymptotic spectral gap’’ for the resonances of the Laplacian as follows.

**Definition 7.8.** The **asymptotic spectral gap** of resonances of the Laplacian  $\Delta$  is defined by

$$a_{\text{asymp}} := \limsup_{b \rightarrow \infty} \{ \text{Re}(s) \text{ s.t. } s \in \text{Res}(\Delta), |\text{Im}(s)| > b \}.$$

The setting (7.4) gives  $D(x) = V - \frac{1}{2}J = \left(\frac{1}{2} - a\right) J(x)$  hence our estimate (2.25) gives that  $a_{\text{asymp}} \leq \frac{1}{2}$ . However this result concerning the resonances of the hyperbolic Laplacian is not new: from the self-adjoint properties of the Laplacian  $\Delta$  in  $L^2$  space we have that  $\text{Im}(s(1-s)) \leq 0$  and this gives that

$$a_{\text{asymp}} \leq \frac{1}{2}. \quad (7.8)$$

*Remark 7.9.* If  $\delta$  denotes the dimension of the limit set (equal to the dimension of the trapped set  $K$ ) a result from F. Naud gives [40]:  $\exists \varepsilon > 0$  s.t.

$$a_{\text{asymp}} \leq (1 - \varepsilon) \delta$$

which improves (7.8) if  $\delta \leq 1/2$ .

### 7.3 Proof of minimal captivity for both models

We give now the proof of Propositions 7.1 and 7.5. Note first that in both models the contracting maps are Möebius maps i.e. of the form  $x'_j = \phi_{i,j}(x) = \frac{a_j x + b_j}{c_j x + d_j} = g_j(x)$  with  $2 \times 2$  matrices  $g_j = \begin{pmatrix} a_j & b_j \\ c_j & d_j \end{pmatrix}$  with  $D_j := \det g_j = \pm 1$ . For the truncated Gauss map these matrices are

$$g_j = \begin{pmatrix} 0 & 1 \\ 1 & j \end{pmatrix} = G_j^{-1} \quad (7.9)$$

with  $j = 1, \dots, N$  and  $D := D_j = -1$ . For the Bowen-Series maps we have

$$g_j = S_j^{-1} \in \mathrm{SL}_2\mathbb{R} \quad (7.10)$$

with  $j = 1, \dots, 2r$  and  $D := D_j = +1$ .

The following proposition shows that there exists coordinates  $(x, \eta)$  on phase space such that the canonical map  $F = (F_j)_{j=1\dots N}$  is decoupled in a product of identical maps.

**Lemma 7.10.** *The canonical map  $F$  defined in (4.3) is the union of the following maps  $F_j$ , with  $j = 1, \dots, N$ :*

$$(x'_j, \xi'_j) = F_j(x, \xi) = \left( g_j(x), (g_j^{-1})'(x'_j) \xi + \tau'(x'_j) \right) \quad (7.11)$$

$$= \left( g_j(x), D_j \cdot (c_j x + d_j)^2 \xi - 2c_j(c_j x + d_j) \right). \quad (7.12)$$

Using the change of variables  $(x, \eta) = \Phi(x, \xi) \in \overline{\mathbb{R}} \times \overline{\mathbb{R}}$  with  $\overline{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$  and

$$\eta := x - \frac{2D}{\xi}, \quad (7.13)$$

the map  $F_j$  gets the simpler “decoupled expression”

$$(x'_j, \eta'_j) = (\Phi \circ F_j \circ \Phi^{-1})(x, \eta) = (g_j(x), g_j(\eta)). \quad (7.14)$$

*Remark 7.11.* Geometrically these new variables  $(x, \eta)$  can be interpreted as the limit points  $(x, \eta) \in \partial\mathbb{H}$  of a geodesic. The map  $(x', \eta') = (\Phi \circ F \circ \Phi^{-1})(x, \eta)$  is simply the Poincaré's map of the geodesic flow [13].

*Proof.* One has

$$g_j^{-1} = D_j \cdot \begin{pmatrix} d_j & -b_j \\ -c_j & a_j \end{pmatrix}, \quad (g_j^{-1})(y) = \frac{d_j y - b_j}{-c_j y + a_j}$$

and

$$(g_j^{-1})'(y) = D_j \cdot (a_j - c_j y)^{-2} = D_j \cdot (c_j x + d_j)^2 \text{ if } y = g_j(x).$$

The roof function is given by (7.1):

$$\begin{aligned}\tau(y) &= -J(y) = -\log \left( \left| (\phi_{i,j}^{-1})'(y) \right| \right) = -\log \left( \left| (g_j^{-1})'(y) \right| \right) \\ &= 2 \log (a_j - c_j y).\end{aligned}$$

So  $\tau'(y) = -2c_j (a_j - c_j y)^{-1} = -2c_j (c_j x + d_j)$  and

$$(x'_j, \xi'_j) = F_j(x, \xi) \stackrel{(7.11)}{=} (g_j(x), D \cdot (c_j x + d_j)^2 \xi - 2c_j (c_j x + d_j)) \quad (7.15)$$

giving (7.12). Now we use the change of variable

$$\xi = \frac{2D}{x - \eta}. \quad (7.16)$$

So

$$\begin{aligned}\xi'_j &= D \cdot (c_j x + d_j)^2 \xi - 2c_j (c_j x + d_j) \\ &= D \cdot (c_j x + d_j)^2 \frac{2D}{(x - \eta)} - 2c_j (c_j x + d_j) \\ &= \frac{2(c_j x + d_j)}{(x - \eta)} (c_j \eta + d_j).\end{aligned}$$

Then

$$\begin{aligned}\eta'_j &= x'_j - \frac{2D}{\xi'_j} = \frac{a_j x + b_j}{c_j x + d_j} - \frac{D(x - \eta)}{(c_j x + d_j)(c_j \eta + d_j)} \\ &= \frac{(a_j x + b_j)(c_j \eta + d_j) - (a_j d_j - b_j c_j)(x - \eta)}{(c_j x + d_j)(c_j \eta + d_j)} = \frac{a_j \eta + b_j}{c_j \eta + d_j} = g_j(\eta).\end{aligned}$$

□

Recall that the multivalued map  $\phi = (\phi_{i,j} = g_j)_j$  has a trapped set  $K$  defined in (2.7) as  $K = \bigcap_{n \geq 1} \phi^n(I)$ . The basin of  $K$  on  $\overline{\mathbb{R}}$  is  $\mathcal{B}(K) := \{x \in \overline{\mathbb{R}}, \exists n \geq 0, \phi^n(x) \in I\} \subset \overline{\mathbb{R}}$ .

**Lemma 7.12.** *The trapped set in phase space  $\mathcal{K}$  defined in (4.7) is contained in the following set:*

$$\mathcal{K} \subset \{(x, \xi), x \in I, \eta \notin \mathcal{B}(K) \text{ with } (x, \eta) = \Phi(x, \xi)\} \quad (7.17)$$



*Proof.* Let  $(x, \xi) \in I \times \mathbb{R}$  which does not belong to the set defined on the right hand side of (7.17). Then  $\eta \in \mathcal{B}(K)$ . Hence for every admissible word  $w \in \mathcal{W}$ , we have that  $|\phi_{w_0,n}(x) - \phi_{w_0,n}(\eta)| \leq C.\theta^n \xrightarrow{n \rightarrow +\infty} 0$ . From the change of variable (7.16) and the expression (7.14) with the new variables, this gives that  $(x_n, \xi_n) := F_{w_0,n}(x, \xi)$  satisfies

$$|\xi_n| = \frac{2}{|\phi_{w_0,n}(x) - \phi_{w_0,n}(\eta)|} \geq C'.\theta^{-n} \rightarrow +\infty$$

hence  $(x, \xi) \notin \mathcal{K}$ . We deduce (7.17).  $\square$

Finally, we show minimal captivity of the canonical map  $F$ . According to (4.8), we have to show that there exists a neighborhood  $B$  of  $\mathcal{K}$  such that  $\forall (x, \xi) \in B, \# \{F(x, \xi) \cap B\} \leq 1$ . This is true if  $B_j := F_j^{-1}(B), j = 1 \dots N$  are disjoint sets. Using the coordinates  $(x, \eta)$  which decouple the map  $F_j$ , in (7.14), it is equivalent to show that there exists a neighborhood  $\mathcal{B}$  of  $K$  in  $\overline{\mathbb{R}}$  such that  $\mathcal{B}_j := g_j^{-1}(\mathcal{B}) \subset \overline{\mathbb{R}}, j = 1, \dots, N$  are disjoint sets. For this we consider both cases:

**Minimal captivity of the truncated Gauss map.** For this map, let  $\mathcal{B} := ] - \infty, -1[$ . Then the sets  $g_j^{-1}(] - \infty, -1[) = ] - j - 1, j[$ , with  $j = 1 \dots N$ , are mutually disjoint. From the argument above this implies that the truncated Gauss map is minimal captive, i.e. Proposition 7.1 holds. Notice that, from (7.16), in variables  $(x, \xi) \in T^*[0, 1]$  we have

$$B = \{x \in [0, 1], \eta \in ] - \infty, -1[\} = \left\{ (x, \xi), x \in [0, 1], \frac{-2}{x+1} < \xi < 0 \right\}$$

This set  $B$  contains the trapped set  $\mathcal{K}_N$  and is depicted in figure (7.1).

**Minimal captivity of the Bowen Series map.** For this case, let  $\mathcal{B} := I = \bigcup_{j=1}^{2r} I_j$ . Then  $\mathcal{B}_j = g_j^{-1}(\mathcal{B}) = g_{j+r}(I) \subset I_{j+r}$ . Since the sets  $I_{j+r}$  are mutually disjoint, the sets  $\mathcal{B}_j$  are also disjoint. From the argument above this implies that the Bowen Series map on phase space is minimal captive, i.e. Proposition 7.5 holds.

Figure (7.5) shows the sets  $B_j = F_j^{-1}(B)$  with  $B := \{x \in I, \eta \in \mathcal{B}\}$  and

$$B_j = \{x \in I, \eta \in \mathcal{B}_j\}$$

that we have used in the proof of minimal captivity.

## A General lemmas on singular values of compact operators

Let  $(P_n)_{n \in \mathbb{N}}$  be a family of compact operators on some Hilbert space. For every  $n \in \mathbb{N}$  let  $(\lambda_{j,n})_{j \in \mathbb{N}} \in \mathbb{C}$  be the sequence of eigenvalues of  $P_n$  counted with multiplicity (i.e. repeated according to the dimension of the eigenspace) and ordered decreasingly:

$$|\lambda_{0,n}| \geq |\lambda_{1,n}| \geq \dots$$




Figure 7.5: This figure illustrates the choice of the bounding functions in the proof of the minimal captive property for the example of a Schottky surface shown in Figure 7.4. The light shaded regions indicate the set  $B_{(j)} := B \cap (I_j \times \mathbb{R})$  while the darker shaded regions indicate the different pre-images  $B_{(i,j)} := F_{ij}^{-1}(B \cap (I_j \times \mathbb{R})) \subset B_{(i)}$ ,  $i \neq j + 2 \pmod{4}$ . For example, that dark orange shaded regions  $B_{(3,3)}, B_{(4,3)}, B_{(2,3)}$  show the three preimages of the light orange region  $B_{(3)}$ . The trapped set  $\mathcal{K}$  is contained in the union of these  $B_{(i,j)}$ .

In the same manner, define  $(\mu_{j,n})_{j \in \mathbb{N}} \in \mathbb{R}^+$ , the decreasing sequence of singular values of  $P_n$ , i.e. the eigenvalues of  $\sqrt{P_n^* P_n}$ .

**Lemma A.1.** *Suppose there exists a map  $N : \mathbb{N} \rightarrow \mathbb{N}$  s.t.  $N(n) \xrightarrow{n \rightarrow \infty} \infty$  and  $\mu_{N(n),n} \xrightarrow{n \rightarrow \infty} 0$ , then  $\forall C > 1$ ,  $|\lambda_{[C \cdot N(n)],n}| \xrightarrow{n \rightarrow \infty} 0$  where  $[\cdot]$  stands for the integer part.*

**Corollary A.2.** *Suppose there exists a map  $N : \mathbb{N} \rightarrow \mathbb{N}$  s.t.  $\forall \varepsilon > 0$ ,  $\exists A_\varepsilon \geq 0$  s.t.  $\forall n \geq A_\varepsilon$ ,*

$$\#\{j \in \mathbb{N} \text{ s.t. } \mu_{j,n} > \varepsilon\} < N(n),$$

*then  $\forall C > 1$ ,  $\forall \varepsilon > 0$ ,  $\exists B_{C,\varepsilon} \geq 0$  s.t.  $\forall n \geq B_{C,\varepsilon}$*

$$\#\{j \in \mathbb{N} \text{ s.t. } |\lambda_{j,n}| > \varepsilon\} \leq C \cdot N(n). \quad (\text{A.1})$$

*Proof. (Of corollary A.2).* Suppose that for any  $\varepsilon > 0$ , there exists  $A_\varepsilon$  s.t. for all  $n \geq A_\varepsilon$ ,  $\#\{j \in \mathbb{N} \text{ s.t. } \mu_{j,n} > \varepsilon\} < N(n)$ . Then  $\mu_{N(n),n} \xrightarrow{n \rightarrow \infty} 0$  and from Lemma A.1,  $\forall C > 1$ ,  $|\lambda_{[C \cdot N(n)],n}| \xrightarrow{n \rightarrow \infty} 0$ , which can be directly restated as (A.1).  $\square$

*Proof.* (Of lemma A.1). Let  $m_{j,n} := -\log \mu_{j,n}$  and  $l_{j,n} := -\log |\lambda_{j,n}|$ ,  $M_{k,n} := \sum_{j=0}^k m_{j,n}$  and  $L_{k,n} := \sum_{j=0}^k l_{j,n}$ . Weyl inequalities relate singular values and eigenvalues by (see [27] p. 50 for a proof) :

$$\prod_{j=1}^k \mu_{j,n} \geq \prod_{j=1}^k |\lambda_{j,n}|, \quad \forall k \geq 1. \quad (\text{A.2})$$

This rewrites:

$$M_{k,n} \leq L_{k,n}, \quad \forall k, n. \quad (\text{A.3})$$

The sequence  $(l_{j,n})_{j \geq 0}$  is increasing in  $j$  so,  $\forall n, \forall k$  we have

$$k \cdot l_{k,n} \geq L_{k,n}. \quad (\text{A.4})$$

Suppose that  $\mu_{N(n),n} \rightarrow 0$  as  $n \rightarrow \infty$  hence

$$m_{N(n),n} \xrightarrow{n \rightarrow \infty} \infty. \quad (\text{A.5})$$

Let  $C > 1$ . The sequence  $(m_{j,n})_{j \geq 0}$  is increasing in  $j$  hence

$$M_{[C \cdot N(n)],n} \geq ([C \cdot N(n)] - N(n)) \cdot m_{N(n),n}, \quad (\text{A.6})$$

hence

$$\begin{aligned} l_{[C \cdot N(n)],n} &\stackrel{(\text{A.4})}{\geq} \frac{1}{[C \cdot N(n)]} \cdot L_{[C \cdot N(n)],n} \stackrel{(\text{A.3})}{\geq} \frac{1}{[C \cdot N(n)]} M_{[C \cdot N(n)],n} \\ &\stackrel{(\text{A.6})}{\geq} \frac{[C \cdot N(n)] - N(n)}{[C \cdot N(n)]} \cdot m_{N(n),n} \xrightarrow{(\text{A.5})} \infty \end{aligned}$$

Thus  $l_{[C \cdot N(n)],n} \xrightarrow{n \rightarrow \infty} \infty$  and  $|\lambda_{[C \cdot N(n)],n}| \xrightarrow{n \rightarrow \infty} 0$ .  $\square$

## B Symbol classes of local $\hbar$ -order

In this Appendix we will first repeat the definitions of the standard symbol classes which are used in this article as well as their well known quantization rules. Then we will introduce a new symbol class which allows  $\hbar$ -dependent order functions and will prove some of the classical results which are known in the usual case for these new symbol classes.

### B.1 Standard semiclassical symbol classes and their quantization

The standard symbol classes (see e.g. [57] chapter 4 or [15] ch 7) of  $\hbar$ PDO's are defined with respect to an order function  $f(x, \xi)$ . This order function is required to be a smooth positive valued function on  $\mathbb{R}^{2n}$  such that there are constants  $C_0$  and  $N_0$  fulfilling

$$f(x, \xi) \leq C_0 \langle (x, \xi) - (x', \xi') \rangle^{N_0} f(x', \xi') \quad (\text{B.1})$$

where we used the notation  $\langle y \rangle := \sqrt{1 + |y|^2}$  for  $y \in \mathbb{R}^k$ . An important example of such an order function is given by  $f(x, \xi) = \langle \xi \rangle^m$  with  $m \in \mathbb{R}$ .

**Definition B.1.** For  $0 \leq \mu \leq \frac{1}{2}$  and  $k \in \mathbb{R}$  the symbol classes  $\hbar^k S_\mu(f)$  contain all families of functions  $a_\hbar(x, \xi) \in C^\infty(\mathbb{R}^{2n})$  parametrized by a parameter  $\hbar \in ]0, \hbar_0]$  such that

$$|\partial_x^\alpha \partial_\xi^\beta a_\hbar(x, \xi)| \leq C \hbar^{k - \mu(|\alpha| + |\beta|)} \langle \xi \rangle^m$$

where  $C$  depends only on  $\alpha, \beta \in \mathbb{N}^n$ .

Unless we want to emphasize the dependence of the symbol  $a_\hbar$  on  $\hbar$  we will drop the index in the following. For the special case of the order function  $f(x, \xi) = \langle \xi \rangle^m$  we also write  $S_\mu^m = S_\mu(\langle \xi \rangle^m)$ , if  $\mu = 0$  we write  $S(f) := S_0(f)$ .

As quantization we use two different quantization rules in this article which are called standard quantization respectively Weyl quantization.

**Definition B.2.** Let  $a_\hbar \in S_\mu(f)$  the Weyl quantization is a family of operators  $\text{Op}_\hbar^w(a) : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ , defined by

$$(\text{Op}_\hbar^w(a_\hbar)\varphi)(x) = (2\pi\hbar)^{-n} \int e^{\frac{i}{\hbar}\xi(x-y)} a_\hbar\left(\frac{x+y}{2}, \xi\right) \varphi(y) dy d\xi, \quad \varphi \in \mathcal{S}(\mathbb{R}^n). \quad (\text{B.2})$$

while the standard quantization  $\text{Op}_\hbar(a) : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$  is given by

$$(\text{Op}_\hbar(a_\hbar)\varphi)(x) = (2\pi\hbar)^{-n} \int e^{\frac{i}{\hbar}\xi(x-y)} a_\hbar(x, \xi) \varphi(y) dy d\xi, \quad \varphi \in \mathcal{S}(\mathbb{R}^n). \quad (\text{B.3})$$

Both quantization extend continuously to operators on  $\mathcal{S}'(\mathbb{R}^n)$ . While the standard quantization is slightly easier to define, the Weyl quantization has the advantage, that real symbols are mapped to formally self adjoint operators.

## B.2 Definition of the symbol classes $S_\mu(A_\hbar)$

In the standard  $\hbar$ -PDO calculus the symbols are ordered by their asymptotic behavior for  $\hbar \rightarrow 0$ . If we take for example a symbol  $a \in \hbar^k S_\mu(f)$  then  $a(x, \xi)$  is of order  $\hbar^k$  for all  $(x, \xi) \in \mathbb{R}^{2n}$ . The symbol classes that we will now introduce will also allow  $\hbar$ -dependent order function which will allow to control the  $\hbar$ -order of a symbol locally, i.e. dependent on  $(x, \xi)$ . First we define these  $\hbar$ -dependent order functions:

**Definition B.3.** Let  $f$  be an order function on  $\mathbb{R}^{2n}$  and  $0 \leq \mu \leq \frac{1}{2}$ . Let  $A_{\hbar} \in S_{\mu}(f)$  be a (possibly  $\hbar$ -dependent) positive symbol such that for some  $c \geq 0$  there is a constant  $C$  that fulfills

$$A_{\hbar}(x, \xi) \geq C\hbar^c f(x, \xi) \quad (\text{B.4})$$

and that for all multi-indices  $\alpha, \beta \in \mathbb{N}^n$ :

$$\left| \partial_x^{\alpha} \partial_{\xi}^{\beta} A_{\hbar}(x, \xi) \right| \leq C_{\alpha, \beta} \hbar^{-\mu(|\alpha|+|\beta|)} A_{\hbar}(x, \xi) \quad (\text{B.5})$$

holds. Then we call  $A_{\hbar}$  an  $\hbar$ -dependent order function and say  $A_{\hbar} \in \mathcal{OF}^c(f)$ .

**Definition B.4.** Let  $0 \leq \mu \leq \frac{1}{2}$  and  $A_{\hbar}$  be a  $\hbar$  dependent order function. The symbol class  $S_{\mu}(A_{\hbar})$  is then defined to be the space of smooth functions  $a_{\hbar}(x, \xi)$  defined on  $\mathbb{R}^{2n}$  and parametrized by a parameter  $\hbar \in ]0, \hbar_0]$  such that

$$\left| \partial_x^{\alpha} \partial_{\xi}^{\beta} a_{\hbar}(x, \xi) \right| \leq C_{\alpha, \beta} \hbar^{-\mu(|\alpha|+|\beta|)} A_{\hbar}(x, \xi) \quad (\text{B.6})$$

By  $\hbar^k S_{\mu}(A_{\hbar})$  we will as usual denote the symbols  $a_{\hbar}$  for which  $\hbar^{-k} a_{\hbar} \in S_{\mu}(A_{\hbar})$ .

The set  $S_{\mu}(A_{\hbar})$  does only depend on the  $\hbar$ -dependent order function  $A_{\hbar}$  and the real parameter  $0 \leq \mu \leq \frac{1}{2}$ . From the Definition B.3 of  $\hbar$ -dependent order function we conclude however that there is an order function  $f$  such that  $A_{\hbar} \in \mathcal{OF}^c(f)$ .

As  $A_{\hbar}(x, \xi) \leq C_0 f(x, \xi)$  and from (B.5) it is obvious, that

$$S_{\mu}(A_{\hbar}) \subset S_{\mu}(f) \quad (\text{B.7})$$

and via this inclusion for  $a_{\hbar} \in S_{\mu}(A_{\hbar})$  the standard quantization  $Op_{\hbar}(a)$  and the Weyl quantization  $Op_{\hbar}^w(a_{\hbar})$  are well defined and give continuous operators on  $\mathcal{S}(\mathbb{R}^n)$  respectively on  $\mathcal{S}'(\mathbb{R}^n)$  (see e.g. [57, Theorem 4.16]). Furthermore equation (B.4) gives us a second inclusion

$$S_{\mu}(f) \subset \hbar^{-c} S_{\mu}(A_{\hbar}) \quad (\text{B.8})$$

thus combining these two inclusions we have:

$$\hbar^c S_{\mu}(f) \subset S_{\mu}(A_{\hbar}) \subset S_{\mu}(f).$$

As for standard  $\hbar$ -PDO symbols we can define asymptotic expansions:

**Definition B.5.** Let  $a_j \in S_\mu(A_\hbar)$  for  $j = 0, 1, \dots$  then we call  $\sum_j \hbar^j a_j$  an asymptotic expansion of  $a \in S_\mu(A_\hbar)$  (writing  $a \sim \sum_j \hbar^j a_j$ ) if and only if for all  $k \in \mathbb{N}$

$$a - \sum_{j < k} \hbar^j a_j \in \hbar^k S_\mu(A_\hbar).$$

As in for the standard  $\hbar$ -PDOs we have some sort of Borel's theorem also for symbols in  $S_\mu(A_\hbar)$ .

**Proposition B.6.** Let  $a_j \in S_\mu(A_\hbar)$  for  $j = 0, 1, \dots$  then there is a symbol  $a \in S_\mu(A_\hbar)$  such that for all  $k \in \mathbb{N}$ :

$$a - \sum_{j < k} \hbar^j a_j \in \hbar^k S_\mu(A_\hbar). \quad (\text{B.9})$$

*Proof.* Once more we can use the inclusion (B.7) into the standard  $\hbar$ -PDO classes and obtain the existence of a symbol  $a \in S_\mu(f)$  such that (see [57, Theorem 4.15])

$$a - \sum_{j < k} \hbar^j a_j \in \hbar^k S_\mu(f) \quad (\text{B.10})$$

and we will show that this symbol belongs to  $S_\mu(A_\hbar)$  and that (B.9) holds: For the first statement we write

$$a = \underbrace{a - \sum_{j < c} \hbar^j a_j}_{\in \hbar^c S_\mu(f)} + \underbrace{\sum_{j < c} \hbar^j a_j}_{\in S_\mu(A_\hbar)}$$

and use the inverse inclusion (B.8).

In order to prove (B.9) we write

$$a - \sum_{j < k} \hbar^j a_j = a - \underbrace{\sum_{j < k+c} \hbar^j a_j}_{\in \hbar^{c+k} S_\mu(f)} + \underbrace{\sum_{j=k}^{k+c-1} \hbar^j a_j}_{\in \hbar^k S_\mu(A_\hbar)}$$

and use once more (B.8). □

The advantage of this new symbol class is, that the order function  $A_\hbar(x, \xi)$  itself can depend on  $\hbar$  and thus the control in  $\hbar$  can be localized. A simple example for such an order function would be  $A_\hbar = \hbar^{m\mu} \langle \frac{\xi}{\hbar^\mu} \rangle^m \in \mathcal{OF}^{m\mu}(\langle \xi \rangle^m)$ . For  $\xi \neq 0$  this function is of order  $\hbar^0$  whereas for  $\xi = 0$  it is of order  $\hbar^{m\mu}$ . Thus also all symbols in  $S_\mu(A_\hbar)$  have to show this behavior.

### B.3 Composition of symbols

By using the inclusion (B.7) we will show a result for the composition of symbols absolutely analogous to the one in the standard case Theorem 4.18 in [57]. We first note that for  $A_{\hbar} \in \mathcal{O}\mathcal{F}^{c_A}(f_A)$  and  $B_{\hbar} \in \mathcal{O}\mathcal{F}^{c_B}(f_B)$  the product formula for derivative yields that  $A_{\hbar}B_{\hbar} \in \mathcal{O}\mathcal{F}^{c_A+c_B}(f_A f_B)$  and can now formulate the following theorem:

**Theorem B.7.** *Let  $A_{\hbar} \in \mathcal{O}\mathcal{F}^{c_A}(f_A)$  and  $B_{\hbar} \in \mathcal{O}\mathcal{F}^{c_B}(f_B)$  be two  $\hbar$ -dependent order functions and  $a \in S_{\mu}(A_{\hbar})$  and  $b \in S_{\mu}(B_{\hbar})$  two  $\hbar$ -local symbols. Then there is a symbol*

$$a\#b \in S_{\mu}(A_{\hbar}B_{\hbar})$$

such that

$$Op_{\hbar}^w(a)Op_{\hbar}^w(b) = Op_{\hbar}^w(a\#b) \quad (\text{B.11})$$

as operators on  $\mathcal{S}$  and the at first order we have

$$a\#b - ab \in \hbar^{1-2\mu}S_{\mu}(A_{\hbar}B_{\hbar}). \quad (\text{B.12})$$

*Proof.* The standard theorem of composition of  $\hbar$ -PDOs (see e.g. Th 4.18 in [57]) together with the inclusion of symbol-classes (B.7) provides us a symbol  $a\#b \in S_{\mu}(f_A \cdot f_B)$  that fulfills equation (B.11). Furthermore it provides us with a complete asymptotic expansion for  $a\#b$ :

$$a\#b - \sum_{k=0}^{N-1} \left( \frac{1}{k!} \left[ \frac{i\hbar(\langle D_x, D_{\eta} \rangle - \langle D_y, D_{\xi} \rangle)}{2} \right]^k a(x, \xi)b(y, \eta) \right)_{|y=x, \eta=\xi} \in \hbar^{N(1-2\mu)}S_{\mu}(f_A \cdot f_B). \quad (\text{B.13})$$

In order to prove our theorem it thus only rests to show, that  $a\#b \in S_{\mu}(A_{\hbar}B_{\hbar})$  and that equation (B.12) holds. We start with the second one. First let  $N \in \mathbb{N}$  be such that  $(N-1)(1-2\mu) \geq c_A + c_B$ , then equation (B.13) and inclusion (B.8) assure that the remainder term in (B.13) is in  $\hbar^{1-2\mu}S_{\mu}(A_{\hbar}B_{\hbar})$ . For  $0 \leq k \leq N-1$  each term in (B.13) can be written as a sum of finitely many terms of the form

$$\frac{(i\hbar)^k}{2^k k!} \left( D_x^{\alpha} D_{\xi}^{\beta} a(x, \xi) \right) \cdot \left( D_x^{\gamma} D_{\xi}^{\delta} b(x, \xi) \right)$$

where  $\alpha, \beta, \gamma, \delta \in \mathbb{N}^n$  are multi-indices fulfilling  $|\alpha| + |\beta| + |\gamma| + |\delta| = 2k$ . Via the product formula one easily checks, that these terms are all in  $\hbar^{k(1-2\mu)}S_{\mu}(A_{\hbar}B_{\hbar})$  which proves that  $a\#b \in S_{\mu}(A_{\hbar}B_{\hbar})$ .  $\square$

### B.4 Ellipticity and inverses

In this section we will define ellipticity for our new symbol classes and will prove a result on  $L^2$ -invertibility.

**Definition B.8.** We call a symbol  $a \in S_\mu(A_{\hbar})$  elliptic if there is a constant  $C$  such that:

$$|a(x, \xi)| \geq CA_{\hbar}(x, \xi). \quad (\text{B.14})$$

For an  $\hbar$ -dependent order function  $A_{\hbar} \in \mathcal{OF}^c(f)$ , from (B.5) and (B.4) it follows, that  $\hbar^c A_{\hbar}^{-1} \in \mathcal{OF}^c(f^{-1})$  is again a  $\hbar$ -dependent order function and we can formulate the following proposition:

**Proposition B.9.** *If  $a \in S_\mu(A_{\hbar})$  is elliptic then  $a^{-1} \in \hbar^{-c} S_\mu(\hbar^c A_{\hbar}^{-1})$ .*

*Proof.* We have to show, that  $|\partial_x^\alpha \partial_\xi^\beta a^{-1}(x, \xi)| \leq C \hbar^{-\mu(|\alpha|+|\beta|)} A_{\hbar}^{-1}(x, \xi)$  uniformly in  $\hbar, x$  and  $\xi$ . For some first derivative (i.e. for  $\alpha \in \mathbb{N}^{2n}, |\alpha| = 1$ ) we have

$$|\partial_{x,\xi}^\alpha a^{-1}| = \frac{|\partial_{x,\xi}^\alpha a|}{|a^2|} \leq C \frac{\hbar^{-\mu} A_{\hbar}}{A_{\hbar}^2} = C \hbar^{-\mu} A_{\hbar}^{-1}$$

where the inequality is obtained by (B.5) and (B.14). The estimates of higher order derivatives can be obtained by induction.  $\square$

As for standard  $\hbar$ -PDOs this notion of ellipticity implies that the corresponding operators are invertible for sufficiently small  $\hbar$ .

**Proposition B.10.** *Let  $A_{\hbar} \in \mathcal{OF}^c(1)$  and  $a \in S_\mu(A_{\hbar})$  be an elliptic symbol, then  $Op_{\hbar}^w(a) : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$  is a bounded operator. Furthermore there exists  $\hbar_0 > 0$  such that  $Op_{\hbar}^w(a)$  is invertible for all  $\hbar \in ]0, \hbar_0]$ . Its inverse is again bounded and a pseudodifferential operator  $Op_{\hbar}^w(b)$  with symbol  $b \in \hbar^{-c} S_\mu(\hbar^c A_{\hbar}^{-1})$ . At leading order its symbol is given by*

$$b - a^{-1} \in \hbar^{1-2\mu-c} S_\mu(\hbar^c A_{\hbar}^{-1})$$

*Proof.* As  $a \in S_\mu(A_{\hbar}) \subset S_\mu(1)$  the boundedness of  $Op_{\hbar}^w(a)$  follows from Theorem 4.23 in [57]. By Theorem B.7 we calculate

$$Op_{\hbar}^w(a) Op_{\hbar}^w(a^{-1}) = Id + R$$

where  $R = Op_{\hbar}^w(r)$  is a PDO with symbol  $r \in \hbar^{1-2\mu} S_\mu(1)$ . Again from Theorem 4.23 in [57] we obtain  $\|R\|_{L^2} \leq C \hbar^{1-2\mu}$  thus there is  $\hbar_0$  such that  $\|R\|_{L^2} < 1$  for  $\hbar \in ]0, \hbar_0]$ . According to Theorem C.3 in [57] we can conclude that  $Op_{\hbar}^w(a)$  is invertible and that the inverse is



given by  $Op_h^w(a^{-1})(Id + R)^{-1}$ . The semiclassical version of Beal's theorem allows us to conclude that  $(Id + R)^{-1} = \sum_{k=0}^{\infty} (-R)^k$  is a PDO with symbol in  $S_{\mu}(1)$  (cf. Theorem 8.3 and the following remarks in [57]). The representation of  $(Id - R)^{-1}$  as a series finally gives us the symbol of the inverse operator at leading order.  $\square$

## B.5 Egorov's theorem for diffeomorphisms

In this section we will study the behavior of symbols  $a \in S_{\mu}(A_h)$  under variable changes. Let  $\gamma : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a diffeomorphism that equals identity outside some bounded set then the pullback with this coordinate change acts as a continuous operator on  $\mathcal{S}(\mathbb{R}^n)$  by:

$$(\gamma^*u)(x) := u(\gamma(x))$$

which can be extended by its adjoint to a continuous operator  $\gamma^* : \mathcal{S}'(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ . By a variable change of an operator we understand its conjugation by  $\gamma$  and we are interested for which  $a \in S_{\mu}(A_h)$  the conjugated operator  $(\gamma^*)^{-1}Op_h(a)\gamma^*$  is again a  $\hbar$ -PDO with symbol  $a_{\gamma}$ . At leading order this symbol will be the composition of the original symbol with the so called canonical transformation

$$T : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}, (x, \xi) \mapsto (\gamma^{-1}(x), (\partial\gamma(\gamma^{-1}(x)))^T\xi)$$

and the symbol class of  $a_{\gamma}$  will be  $S_{\mu}(A_h \circ T)$ . For the  $A_h \in \mathcal{OF}^c(f)$  defined in Definition B.3 the composition  $A_h \circ T$  will in general however not be a  $\hbar$ -dependent order function itself because the derivatives in  $x$  create a supplementary  $\xi$  factor which has to be compensated (cf. discussion in chapter 9.3 in [57]). We therefore demand in this section that our order function  $A_h$  satisfies:

$$\left| \partial_x^{\alpha} \partial_{\xi}^{\beta} A_h(x, \xi) \right| \leq C_{\alpha, \beta} \hbar^{\mu(|\alpha| + |\beta|)} \langle \xi \rangle^{-|\beta|} A_h(x, \xi). \quad (\text{B.15})$$

A straightforward calculation shows then, that  $A_h \circ T \in \mathcal{OF}^c(f \circ T)$  is again a  $\hbar$ -dependent order function. The same condition has to be fulfilled by the symbol of the conjugated operator:

$$\left| \partial_x^{\alpha} \partial_{\xi}^{\beta} a(x, \xi) \right| \leq \hbar^{-\mu(|\alpha| + |\beta|)} \langle \xi \rangle^{-|\beta|} A_h(x, \xi) \quad (\text{B.16})$$

**Theorem B.11.** *Let  $A_{\hbar}$  be an  $\hbar$ -dependent order function that fulfills (B.15). Let  $a \in S_{\mu}(A_{\hbar})$  be an symbol which fulfills (B.16) and has compact support in  $x$  (i.e.  $\overline{\{x \in \mathbb{R}^n \mid \exists \xi \in \mathbb{R}^n : a(x, \xi) \neq 0\}}$  is compact) and let  $\gamma : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a diffeomorphism. Then there is a symbol  $a_{\gamma} \in S_{\mu}(A_{\hbar} \circ T)$  such that*

$$(Op_{\hbar}(a_{\gamma})u)(\gamma(x)) = (Op_{\hbar}(a)(u \circ \gamma))(x) \quad (\text{B.17})$$

for all  $u \in \mathcal{S}'(\mathbb{R}^n)$ . Furthermore  $a_{\gamma}$  has the following asymptotic expansion.

$$a_{\gamma}(\gamma(x), \eta) \sim \sum_{n=0}^{k-n} \frac{1}{\nu!} \langle i \frac{\hbar}{\langle \eta \rangle} D_y, D_{\xi} \rangle^{\nu} e^{i \frac{\hbar}{\langle \eta \rangle} \langle \rho_x(y), \eta \rangle} a(x, \xi) \Big|_{y=0, \xi=(\partial \gamma(x))^T \eta} \quad (\text{B.18})$$

where  $\rho_x(y) = \gamma(y+x) - \gamma(x) - \gamma'(x)y$ . The terms of the series are in  $\hbar^{\frac{\nu(1-2\mu)}{2}} S_{\mu}(\langle \eta \rangle^{\frac{\nu}{2}} A_{\hbar} \circ T(\gamma(x), \eta))$ .

We will prove this theorem similar to Theorem 18.1.17 in [31] by using a parameter dependent stationary phase approximation (Theorem 7.7.7 in [32]) as well as the following proposition which forms the analog to Proposition 18.1.4 of [31] for our symbol classes and which we will prove first.

**Proposition B.12.** *Let  $a(x, \xi; \hbar) \in C^{\infty}(\mathbb{R}^{2n})$  a family of smooth functions that fulfills*

$$|\partial_x^{\alpha} \partial_{\xi}^{\beta} a(x, \xi)| \leq C \hbar^{-l} \langle \xi \rangle^l f(x, \xi) \quad (\text{B.19})$$

where  $C$  and  $l$  may depend on  $\alpha$  and  $\beta$ . Let  $a_j \in S_{\mu}(A_{\hbar})$ ,  $j = 0, 1, \dots$  be a sequence of symbols such that

$$|a(x, \xi) - \sum_{j < k} \hbar^j a_j(x, \xi)| \leq C \hbar^{\tau k} \langle \xi \rangle^{-\tau k} f(x, \xi) \quad (\text{B.20})$$

where  $\tau > 0$ . Then  $a \in S_{\mu}(A_{\hbar})$  and  $a \sim \sum \hbar^j a_j$ .

*Proof.* We have to show that for all  $k \geq 0$  and  $g_k(x, \xi) := a(x, \xi) - \sum_{j < k} \hbar^j a_j(x, \xi)$  we

have  $|\partial_x^{\alpha} \partial_{\xi}^{\beta} g_k| \leq C \hbar^{k-\mu(|\alpha|+|\beta|)} A_{\hbar}$ . This result can be obtained by iterating the following argument for the first derivative in  $x_1$ :

Let  $e_1 \in \mathbb{R}^n$  be the first eigenvector and  $0 < \varepsilon < 1$ . For arbitrary  $j \in \mathbb{N}$  we can write by Taylor's Formula

$$|g_j(x + \varepsilon e_1, \xi) - g_j(x, \xi) - \partial_{x_1} g_j(x, \xi) \varepsilon| \leq C \varepsilon^2 \sup_{t \in [0, \varepsilon]} |\partial_{x_1}^2 g_j(x + t e_1, \xi)|.$$

From (B.19) and the property, that all  $a_j$  are in  $S_\mu(A_\hbar)$  we get

$$\sup_{t \in [0, \varepsilon]} |\partial_{x_1}^2 g_j(x + te_1, \xi)| \leq C \hbar^{-l} \langle \xi \rangle^l f(x, \xi)$$

for some  $l \in \mathbb{R}$  and get

$$|\partial_{x_1} g_j(x, \xi)| \leq C \varepsilon \hbar^{-l} \langle \xi \rangle^l m(x, \xi) + \frac{|g_j(x + \varepsilon e_1, \xi) - g_j(x, \xi)|}{\varepsilon}$$

which turns for  $j > \frac{2k+2c+l}{\tau}$  and  $\varepsilon = \hbar^{k+l+c} \langle \xi \rangle^{-(k+l+c)}$  into:

$$|\partial_{x_1} g_j(x, \xi)| \leq C \hbar^{c+k} \langle \xi \rangle^{-(c+k)} f(x, \xi) \leq C \hbar^k A_\hbar(x, \xi)$$

where we used (B.8) in the second equation. Thus

$$|\partial_{x_1} g_k(x, \xi)| \leq C \hbar^k A_\hbar(x, \xi) + \left| \sum_{i=k}^j \hbar^i \partial_{x_1} a_i(x, \xi) \right| \leq C \hbar^{k-\mu} A_\hbar(x, \xi)$$

which finishes the proof.  $\square$

After having proven this proposition we can start with the proof of Theorem B.11:

*Proof.* If we define

$$a_\gamma(\gamma(x), \eta) := e^{-\frac{i}{\hbar} \gamma(x) \eta} Op_\hbar(a) e^{\frac{i}{\hbar} \gamma(\cdot) \eta} \quad (\text{B.21})$$

then equation (B.17) holds for all  $e^{\frac{i}{\hbar} x \eta}$  which form a dense subset of  $\mathcal{S}'(\mathbb{R}^n)$ . We thus have to show that  $a_\gamma$  defined in (B.21) is in  $S_\mu(A_\hbar)$  and that (B.18) holds.

We will first write  $a_\gamma$  as an oscillating integral in order to apply the stationary phase theorem. By definition of  $Op_\hbar(a)$  one obtains

$$a_\gamma(\gamma(x), \eta) = \frac{1}{(2\pi\hbar)^n} \iint a(x, \tilde{\xi}) e^{\frac{i}{\hbar} ((x-\tilde{y})\tilde{\xi} + (\gamma(\tilde{y}) - \gamma(x))\eta)} d\tilde{y} d\tilde{\xi}$$

which we can transform by a variable transformation  $\tilde{\xi} = \langle \eta \rangle \xi$  and  $\tilde{y} = y + x$  into

$$a_\gamma(\gamma(x), \eta) = \frac{1}{(2\pi\tilde{\hbar})^n} \iint a(x, \langle \eta \rangle \xi) e^{\frac{i}{\tilde{\hbar}} (-y\xi + (\gamma(y+x) - \gamma(x)) \frac{\eta}{\langle \eta \rangle})} dy d\xi$$

where  $\tilde{\hbar} = \frac{\hbar}{\langle \eta \rangle}$ .

The critical points of the phase function are given by

$$y = 0 \text{ and } \xi = (\partial\gamma(x))^T \frac{\eta}{\langle \eta \rangle}.$$

Let  $\chi \in C_c^\infty([-2, 2]^n)$  such that  $\chi = 1$  on  $[-1, 1]^n$  then we can write

$$a_\gamma(\gamma(x), \eta) = I_1(\tilde{\hbar}) + I_2(\tilde{\hbar})$$

with

$$I_1(\tilde{\hbar}) = \frac{1}{(2\pi\tilde{\hbar})^n} \iint \chi(y) \chi\left(\xi - (\partial\gamma(x))^T \frac{\eta}{\langle\eta\rangle}\right) a(x, \langle\eta\rangle\xi) e^{\frac{i}{\tilde{\hbar}}(-y\xi + (\gamma(y+x) - \gamma(x))\frac{\eta}{\langle\eta\rangle})} dy d\xi$$

and

$$I_2(\tilde{\hbar}) = \frac{1}{(2\pi\tilde{\hbar})^n} \iint \left(1 - \chi(y) \chi\left(\xi - (\partial\gamma(x))^T \frac{\eta}{\langle\eta\rangle}\right)\right) a(x, \langle\eta\rangle\xi) e^{\frac{i}{\tilde{\hbar}}(-y\xi + (\gamma(y+x) - \gamma(x))\frac{\eta}{\langle\eta\rangle})} dy d\xi.$$

While  $I_1(\tilde{\hbar})$  still contains critical points, for  $I_2(\tilde{\hbar})$  there are no critical points in the support of the integrand anymore.

$I_1$  is of the form studied in Theorem 7.7.7 in [32]. Here the role of  $x$  and  $y$  is interchanged and there is an additional parameter  $\frac{\eta}{\langle\eta\rangle}$ . We thus get from this stationary phase theorem

$$\begin{aligned} & \left| I_1(\tilde{\hbar}) - \sum_{\nu=0}^{k-n} \frac{1}{\nu!} \langle i\tilde{\hbar}D_y, D_\xi \rangle^\nu e^{\frac{i}{\tilde{\hbar}}\langle\rho_x(y), \frac{\eta}{\langle\eta\rangle}\rangle} u(x, \xi, y, \eta) \Big|_{y=0, \xi=(\partial\gamma(x))^T \frac{\eta}{\langle\eta\rangle}} \right| \\ & \leq C\tilde{\hbar}^{\frac{k+n}{2}} \sum_{|\alpha|\leq 2k} \sup_{y, \xi} |D_{y, \xi}^\alpha u(x, \xi, y, \eta)| \end{aligned} \quad (\text{B.22})$$

where  $u(x, \xi, y, \eta) = \chi(y) \chi\left(\xi - (\partial\gamma(x))^T \frac{\eta}{\langle\eta\rangle}\right) a(x, \langle\eta\rangle\xi)$ . Because of (B.16) and (B.1) we can estimate

$$\sup_{y, \xi} |D_{y, \xi}^\alpha u(x, \xi, y, \eta)| \leq C\tilde{\hbar}^{-\mu|\alpha|} f(x, (\partial\gamma(x))^T \eta) = C\tilde{\hbar}^{-\mu|\alpha|} f \circ T(\gamma(x), \eta).$$

Thus transforming the expansion (B.22) back to an expansion in  $\hbar$  we get

$$\begin{aligned} & \left| I_1(\hbar) - \sum_{\nu=0}^{k-n} \frac{1}{\nu!} \langle i\frac{\hbar}{\langle\eta\rangle} D_y, D_\xi \rangle^\nu e^{\frac{i}{\hbar}\langle\rho_x(y), \eta\rangle} u(x, \xi, y, \eta) \Big|_{y=0, \xi=(\partial\gamma(x))^T \frac{\eta}{\langle\eta\rangle}} \right| \\ & \leq C\hbar^{\frac{k(1-2\mu)+n}{2}} \langle\eta\rangle^{-\frac{k+n}{2}} f \circ T(\gamma(x), \eta). \end{aligned}$$

As the stationary points for  $I_2$  are not contained in the support of the integrand we get by the non stationary phase theorem:

$$|I_2(\hbar)| \leq C \left(\frac{\hbar}{\langle\eta\rangle}\right)^N f \circ T(\gamma(x), \eta)$$

for all  $N \in \mathbb{N}$ . Thus we finally get

$$\begin{aligned} & \left| a_\gamma(\gamma(x), \eta) - \sum_{\nu=0}^{k-n} \frac{1}{\nu!} \langle i\frac{\hbar}{\langle\eta\rangle} D_y, D_\xi \rangle^\nu e^{\frac{i}{\hbar}\langle\rho_x(y), \eta\rangle} u(x, \xi, y, \eta) \Big|_{y=0, \xi=(\partial\gamma(x))^T \frac{\eta}{\langle\eta\rangle}} \right| \\ & \leq C\hbar^{\frac{k(1-2\mu)+n}{2}} \langle\eta\rangle^{-\frac{k+n}{2}} f \circ T(\gamma(x), \eta). \end{aligned} \quad (\text{B.23})$$

If we show that the elements of the series are in  $\hbar^{\frac{\nu(1-2\mu)}{2}} S_\mu(\langle\eta\rangle^{\frac{\nu}{2}} A_\hbar \circ T(\gamma(x), \eta))$  then this equation is of the form (B.20). The terms of order  $\nu$  in the series are of the form

$$\left(\frac{i\hbar}{\langle\eta\rangle}\right)^\nu \partial_y^\alpha e^{\frac{i}{\hbar}\langle\rho_x(y), \eta\rangle} (\partial_\xi^\alpha a)(x, (\partial\gamma(x))^T \eta) \langle\eta\rangle^\nu \Big|_{y=0}$$

Where  $\alpha \in \mathbb{N}^n$  with  $|\alpha| = \nu$ . The second factor  $(\partial_\xi^\alpha a)(x, (\partial\gamma(x))^T \eta) \langle \eta \rangle^\nu$  is in  $\hbar^{-\mu\nu} S_\mu(A_\hbar \circ T(\gamma(x), \eta))$  as we demanded the condition (B.16) on our symbol  $a$ . Thus it remains to show that the other factor is of order  $\left(\frac{\hbar}{\langle \eta \rangle}\right)^{\frac{\nu}{2}}$  on the support of  $a$ . This is the case because  $\rho_x(y)$  vanishes at second order in  $y = 0$ . Each derivative of  $e^{\frac{i}{\hbar} \langle \rho_x(y), \eta \rangle}$  produces a factor  $\frac{i}{\hbar} \langle \partial_{y_i} \rho_x(0), \eta \rangle$ . But as  $\partial_{y_i} \rho_x(0)$  vanishes we need a second derivative, now acting on  $\partial_{y_i} \rho_x(y)$ , in order to get a contribution. Thus in the worst case  $\partial_y^\alpha e^{\frac{i}{\hbar} \langle \rho_x(y), \eta \rangle}$  is of order  $\left(\frac{\hbar}{\langle \eta \rangle}\right)^{-\frac{\nu}{2}}$ . Thus we have shown that (B.23) is of the form (B.20).

The last thing that we have to show is thus, that  $a_\gamma$  fulfills (B.19). If we consider the definition (B.21) of  $a_\gamma$  we see that  $\partial_x^\alpha \partial_\xi^\beta a_\gamma(\gamma(x), \eta)$  can be written as a sum of terms of the form  $\frac{P(\eta)}{\hbar^k} e^{-\frac{i}{\hbar} \gamma(x) \eta} O p_\hbar(b) e^{\frac{i}{\hbar} \gamma(\cdot) \eta}$  where  $b \in S_\mu(A_\hbar \langle \xi \rangle^j)$  and  $P(\eta)$  is a polynomial in  $\eta$ . The constants  $j, k$  and the degree of  $P(\eta)$  depend on  $\alpha$  and  $\beta$ . Thus writing these terms as oscillating integrals and applying the same arguments as above one gets (B.19).

We have thus shown that all the conditions for proposition B.12 are fulfilled and can conclude that  $a_\gamma$  belongs to  $S_\mu(A_\hbar)$  and that (B.23) is also an asymptotic expansion w.r.t. the order function  $A_\hbar$ .  $\square$

## C Adapted Weyl type estimates

**Lemma C.1.** *Let  $a_\hbar \in S_\mu(\langle x \rangle^{-2} \langle \xi \rangle^{-2})$  with  $0 \leq \mu < \frac{1}{2}$  be a real compactly supported symbol as Definition B.4.  $\forall \hbar > 0$ ,  $\hat{A} := \text{Op}_\hbar^w(a_\hbar)$  is self-adjoint and trace class on  $L^2(\mathbb{R})$  and for any  $\epsilon > 0$ , as  $\hbar \rightarrow 0$  :*

$$(2\pi\hbar) \# \left\{ \lambda_i^\hbar \in \sigma(\hat{A}) \mid |\lambda_i^\hbar| \geq \epsilon \right\} \leq C_1 \text{Leb} \{(x, \xi) ; |a_\hbar(x, \xi)| > 0\} + C_2 \hbar \quad (\text{C.1})$$

where  $C_1$  and  $C_2$  are independent of  $\hbar$ .

*Proof.* As  $a_\hbar$  is compactly supported  $\hat{A}$  is trace class for every  $\hbar$  (see theorem C.17 [57]). Consequently also  $\frac{1}{\epsilon^2} \hat{A}^2$  is trace class and its trace is given by Lidskii's theorem by  $\text{Tr}(\frac{1}{\epsilon^2} \hat{A}^2) = \sum_i \left(\frac{\lambda_i^\hbar}{\epsilon}\right)^2$ . As  $\hat{A}$  is self adjoint all  $\lambda_i^\hbar$  are real and one clearly has

$$\# \left\{ \lambda_i^\hbar \in \sigma(\hat{A}) \mid |\lambda_i^\hbar| \geq \epsilon \right\} \leq \text{Tr} \left( \frac{1}{\epsilon^2} \hat{A}^2 \right).$$

If we denote by  $b_\hbar(x, \xi)$  the complete symbol of  $\hat{A}^2$  we can calculate the trace by the following exact formula

$$\text{Tr}(\hat{A}^2) = \frac{1}{2\pi\hbar} \int b_\hbar(x, \xi) dx d\xi$$

For any  $\mu < \frac{1}{2}$  let  $N_\mu \in \mathbb{N}$  be such that  $N_\mu(1 - 2\mu) \geq 1$ . Then using the asymptotic expansion (B.13) for composition of PDOs up to order  $N_\mu$ ,  $b_\hbar$  can be written as  $b_\hbar = b_\hbar^{(1)} + \hbar b_\hbar^{(2)}$  where  $\text{supp} b_\hbar^{(1)} = \text{supp} a_\hbar$  and  $b_\hbar^{(2)} \in S_\mu(\langle x \rangle^{-4} \langle \xi \rangle^{-4})$ . Note that this decomposition depends on  $\mu$  via the choice of the order  $N_\mu$ . Thus

$$\begin{aligned} \frac{1}{\epsilon^2} \text{Tr}(\hat{A}^2) &= \frac{1}{2\pi\hbar\epsilon^2} \left( \int b_\hbar^{(1)}(x, \xi) dx d\xi + \hbar \int b_\hbar^{(2)}(x, \xi) dx d\xi \right) \\ &\leq \frac{1}{2\pi\hbar} (C_1 \text{Leb}(\text{supp}(a_\hbar)) + C_2 \hbar) \end{aligned}$$

The estimate on the first term is obtained because  $b_\hbar^{(1)} \in S_\mu(\langle x \rangle^{-4} \langle \xi \rangle^{-4})$  implies that  $b_\hbar^{(1)}$  is bounded uniformly in  $\hbar$ . Furthermore as stated above,  $b_\hbar^{(1)}$  is compactly supported in  $\text{supp}(a_\hbar)$ . The estimate on the second term follows from the integrable upper bound  $|b_\hbar^{(2)}| \leq C \langle x \rangle^{-4} \langle \xi \rangle^{-4}$ . Finally note that the  $\epsilon$  dependence can be absorbed in the constants  $C_1$  and  $C_2$ .  $\square$

## References

- [1] E. Artin. Ein mechanisches system mit quasiergodischen bahnen. In *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, volume 3, pages 170–175. Springer, 1924.
- [2] V. Baladi and M. Tsujii. Dynamical determinants and spectrum for hyperbolic diffeomorphisms. *Probabilistic and Geometric Structures in Dynamics*, K. Burns, D. Dolgopyat and Ya. Pesin (eds), *Contemp. Math. (Amer. Math. Soc.)*, Volume in honour of M. Brin's 60th birthday. link, 2006.
- [3] V. Baladi and M. Tsujii. Anisotropic Hölder and Sobolev spaces for hyperbolic diffeomorphisms. *Ann. Inst. Fourier*, 57:127–154, 2007.
- [4] V. Baladi and M. Tsujii. Dynamical determinants and spectrum for hyperbolic diffeomorphisms. In *Geometric and probabilistic structures in dynamics*, volume 469 of *Contemp. Math.*, pages 29–68. Amer. Math. Soc., Providence, RI, 2008.
- [5] S. Barkhofen, T. Weich, A. Potzuweit, H-J. Stöckmann, U. Kuhl, and M. Zworski. Experimental observation of the spectral gap in microwave n-disk systems. *Physical review letters*, 110(16):164102, 2013.
- [6] M. Blank, G. Keller, and C. Liverani. Ruelle-Perron-Frobenius spectrum for Anosov maps. *Nonlinearity*, 15:1905–1973, 2002.
- [7] D. Borthwick. *Spectral theory of infinite-area hyperbolic surfaces*. Birkhauser, 2007.
- [8] D. Borthwick. Distribution of resonances for hyperbolic surfaces. *Experimental Mathematics*, 23:25–45, 2014.

- [9] D. Borthwick, C. Judge, and P. A. Perry. Selberg’s zeta function and the spectral geometry of geometrically finite hyperbolic surfaces. *Comment. Math. Helv.*, 80(3):483–515, 2005.
- [10] R. Bowen and C. Series. Markov maps associated with fuchsian groups. *Publications Mathématiques de l’IHÉS*, 50(1):153–170, 1979.
- [11] M. Brin and G. Stuck. *Introduction to Dynamical Systems*. Cambridge University Press, 2002.
- [12] A. Córdoba and C. Fefferman. Wave packets and fourier integral operators. *Communications in Partial Differential Equations*, 3(11):979–1005, 1978.
- [13] F. Dal’Bo. *Trajectoires géodésiques et horocycliques*. EDP sciences, 2012.
- [14] K. Datchev and S. Dyatlov. Fractal Weyl laws for asymptotically hyperbolic manifolds. *Geom. Funct. Anal.*, 23(4):1145–1206, 2013.
- [15] M. Dimassi and J. Sjöstrand. *Spectral Asymptotics in the Semi-Classical Limit*. Cambridge University Press. London Mathematical Society Lecture Notes, Vol. 268, 1999.
- [16] D. Dolgopyat. On decay of correlations in Anosov flows. *Ann. of Math. (2)*, 147(2):357–390, 1998.
- [17] D. Dolgopyat. On mixing properties of compact group extensions of hyperbolic systems. *Israel J. Math.*, 130:157–205, 2002.
- [18] L. Ermann and D. Shepelyansky. Ulam method and fractal weyl law for perron-frobenius operators. *The European Physical Journal B*, 75(3):299–304, 2010.
- [19] K. Falconer. *Techniques in fractal geometry*. John Wiley & Sons Ltd., Chichester, 1997.
- [20] K.J. Falconer. *Fractal geometry: mathematical foundations and applications*. John Wiley & Sons Inc, 2003.
- [21] F. Faure. Semiclassical origin of the spectral gap for transfer operators of a partially expanding map. *Nonlinearity link*, 24:1473–1498, 2011.
- [22] F. Faure and N. Roy. Ruelle-pollicott resonances for real analytic hyperbolic map. *Nonlinearity. link*, 19:1233–1252, 2006.
- [23] F. Faure, N. Roy, and J. Sjöstrand. A semiclassical approach for anosov diffeomorphisms and ruelle resonances. *Open Math. Journal. link*, 1:35–81, 2008.
- [24] F. Faure and J. Sjöstrand. Upper bound on the density of ruelle resonances for anosov flows. a semiclassical approach. *Comm. in Math. Physics, Issue 2. link*, 308:325–364, 2011.

- [25] F. Faure and M. Tsujii. Band structure of the ruelle spectrum of contact anosov flows. *Comptes rendus - Mathématique* 351 , 385-391, (2013) [link](#), 2013.
- [26] F. Faure and M. Tsujii. Prequantum transfer operator for symplectic anosov diffeomorphism. *Asterisque* 375 (2015), [link](#), pages ix+222 pages, 2015.
- [27] I. Gohberg, S. Goldberg, and N. Krupnik. *Traces and Determinants of Linear Operators*. Birkhauser, 2000.
- [28] S. Gouëzel and C. Liverani. Banach spaces adapted to anosov systems. *Ergodic Theory and dynamical systems*, 26:189–217, 2005.
- [29] A. Grigis and J. Sjöstrand. *Microlocal analysis for differential operators*, volume 196 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1994. An introduction.
- [30] L. Guillope, K. Lin, and M. Zworski. The Selberg zeta function for convex co-compact Schottky groups. *Comm. Math. Phys.*, 245(1):149–176, 2004.
- [31] L. Hörmander. *The analysis of linear partial differential operators III*, volume 257. Springer, 1983.
- [32] L. Hormander. *The Analysis of the Linear Partial Differential Operators I: Distribution Theory and Fourier Analysis. Classics in Mathematics*. Springer: Berlin, 2003.
- [33] O. Jenkinson and M. Pollicott. Calculating Hausdorff dimension of Julia sets and Kleinian limit sets. *American Journal of Mathematics*, 124(3):495–545, 2002.
- [34] A.Yu. Kitaev. Fredholm determinants for hyperbolic diffeomorphisms of finite smoothness. *Nonlinearity*, 12(1):141–179, 1999.
- [35] M. Körber, M. Michler, A. Bäcker, and R. Ketzmerick. Hierarchical fractal weyl laws for chaotic resonance states in open mixed systems. *Physical review letters*, 111(11):114102, 2013.
- [36] W. Lu, S. Sridhar, and M. Zworski. Fractal Weyl laws for chaotic open systems. *Physical review letters*, 91(15):154101, 2003.
- [37] A. Martinez. *An Introduction to Semiclassical and Microlocal Analysis*. Universitext. New York, NY: Springer, 2002.
- [38] P. Mattila. *Geometry of sets and measures in Euclidean spaces: fractals and rectifiability*, volume 44. Cambridge University Press, 1999.
- [39] D. H. Mayer. On the thermodynamic formalism for the gauss map. *Communications in mathematical physics*, 130(2):311–333, 1990.



- [40] F. Naud. Expanding maps on cantor sets and analytic continuation of zeta functions. In *Annales Scientifiques de Ecole Normale Supérieure*, volume 38, pages 116–153. Elsevier, 2005.
- [41] S. Nonnenmacher. Spectral problems in open quantum chaos. *Nonlinearity*, 24(12):R123, 2011.
- [42] S. Nonnenmacher, J. Sjöstrand, and M. Zworski. Fractal Weyl law for open quantum chaotic maps. *Ann. Math. (2)*, 179(1):179–251, 2014.
- [43] S. J. Patterson and P. A. Perry. The divisor of selberg’s zeta function for kleinian groups. *Duke Mathematical Journal*, 106(2):321–390, 2001.
- [44] Y. Pesin. *Lectures on Partial Hyperbolicity and Stable Ergodicity*. European Mathematical Society, 2004.
- [45] M. Pollicott and A.C. Rocha. A remarkable formula for the determinant of the laplacian. *Inventiones Mathematicae*, 130(2):399–414, 1997.
- [46] A. Potzuweit, T. Weich, S. Barkhofen, U. Kuhl, H.-J. Stöckmann, and M. Zworski. Weyl asymptotics: From closed to open systems. *Physical Review E*, 86(6):066205, 2012.
- [47] M. Reed and B. Simon. *Mathematical methods in physics, vol I : Functional Analysis*. Academic press, New York, 1972.
- [48] D. Ruelle. Zeta-functions for expanding maps and Anosov flows. *Inventiones mathematicae*, 34(3):231–242, 1976.
- [49] David Ruelle. The thermodynamic formalism for expanding maps. *Communications in Mathematical Physics*, 125(2):239–262, 1989.
- [50] H. Schomerus and J. Tworzydło. Quantum-to-classical crossover of quasibound states in open quantum systems. *Physical review letters*, 93(15):154102, 2004.
- [51] J. Sjöstrand. Geometric bounds on the density of resonances for semiclassical problems. *Duke Math. J.*, 60(1):1–57, 1990.
- [52] M. Taylor. *Partial differential equations, Vol I*. Springer, 1996.
- [53] M. Taylor. *Partial differential equations, Vol II*. Springer, 1996.
- [54] M. Tsujii. Decay of correlations in suspension semi-flows of angle-multiplying maps. *Ergodic Theory and Dynamical Systems*, 28:291–317, 2008.
- [55] F. Wolfgang. Über die Existenz des zweiten Eigenwertes der  $G_N$ -Operatoren bei grossem  $N$ . *Anz. Österr. Akad. Wiss., Math.-Naturwiss. Kl.*, 1991(7):97–101, 1991.

- [56] M. Zworski. Dimension of the limit set and the density of resonances for convex co-compact hyperbolic surfaces. *Invent. Math.*, 136(2):353–409, 1999.
- [57] M. Zworski. *Semiclassical Analysis*. Graduate Studies in Mathematics Series. Amer Mathematical Society, 2012.