

HIDA FAMILIES AND p -ADIC TRIPLE PRODUCT L -FUNCTIONS

MING-LUN HSIEH

ABSTRACT. We construct the three-variable p -adic triple product L -functions attached to Hida families of elliptic newforms and prove the explicit interpolation formulae at all critical specializations by establishing explicit Ichino's formulae for the trilinear period integrals of automorphic forms. Our formulae perfectly fit the conjectural shape of p -adic L -functions predicted by Coates and Perrin-Riou. As an application, we prove the factorization of certain unbalanced p -adic triple product L -functions into a product of anticyclotomic p -adic L -functions for modular forms. By this factorization, we give a new construction of the anticyclotomic p -adic L -functions for elliptic curves in the definite case via the diagonal cycle Euler system à la Darmon and Rotger and obtain a Greenberg-Stevens style proof of anticyclotomic exceptional zero conjecture for elliptic curves due to Bertolini and Darmon.

CONTENTS

1. Introduction	1
2. Classical modular forms and automorphic forms	8
3. The unbalanced p -adic triple product L -functions	16
4. The balanced p -adic triple product L -functions	28
5. The calculation of local zeta integrals (I)	43
6. The calculation of local zeta integrals (II)	52
7. The interpolation formulae	65
8. Applications to anticyclotomic p -adic L -functions	70
References	75

1. INTRODUCTION

The aim of this paper is to construct the three-variable p -adic triple product L -functions attached to Hida families of elliptic newforms in the unbalanced and balanced case with explicit interpolation formulae at all critical specializations. Let $p > 2$ be a prime. Let \mathcal{O} be a valuation ring finite flat over \mathbf{Z}_p . Let \mathbf{I} be a normal domain finite flat over the Iwasawa algebra $\Lambda = \mathcal{O}[[\Gamma]]$ of the topological group $\Gamma = 1 + p\mathbf{Z}_p$. Let

$$\mathbf{F} = (\mathbf{f}, \mathbf{g}, \mathbf{h})$$

be the triple of primitive Hida families of tame conductor (N_1, N_2, N_3) and nebentypus (ψ_1, ψ_2, ψ_3) with coefficients in \mathbf{I} . Roughly speaking, we construct a three-variable Iwasawa function over the weight space of \mathbf{F} interpolating the square root of the algebraic part of central values of the triple product L -function attached to \mathbf{F}_Q and prove explicit interpolation formulae at all critical specializations. We would like to emphasize that our formulae completely comply with the conjectural form described in [CPR89], [Coa89a] and [Coa89b] and is compatible with other known p -adic L -functions. For example, when \mathbf{g} and \mathbf{h} are primitive Hida families of CM forms by some imaginary quadratic field, we show that the unbalanced p -adic L -function is the product of theta elements à la Bertolini-Darmon. In order to state our result precisely, we need to introduce some notation from Hida theory for elliptic modular forms and technical items such as the modified Euler factors at p and the canonical periods of Hida families in the theory of p -adic L -functions.

Date: July 28, 2017.

2010 *Mathematics Subject Classification.* 11F67, 11F33.

This work was partially supported by a MOST grant 103-2115-M-002-012-MY5.

1.1. Galois representations attached to Hida families. If $\mathcal{F} = \sum_{n \geq 1} \mathbf{a}(n, \mathcal{F})q \in \mathbf{I}[[q]]$ is a primitive cuspidal Hida family of tame conductor $N_{\mathcal{F}}$, let $\rho_{\mathcal{F}} : G_{\mathbf{Q}} = \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q}) \rightarrow \text{GL}_2(\text{Frac } \mathbf{I})$ be the associated big Galois representation such that $\text{Tr } \rho_{\mathcal{F}}(\text{Frob}_{\ell}) = \mathbf{a}(\ell, \mathcal{F})$ for primes $\ell \nmid N_{\mathcal{F}}$, where Frob_{ℓ} is the geometric Frobenius at ℓ and let $V_{\mathcal{F}}$ denote the natural realization of $\rho_{\mathcal{F}}$ inside the étale cohomology groups of modular curves. Thus, $V_{\mathcal{F}}$ is a lattice in $(\text{Frac } \mathbf{I})^2$ with the continuous Galois action via $\rho_{\mathcal{F}}$, and the $\text{Gal}(\overline{\mathbf{Q}}_p/\mathbf{Q}_p)$ -invariant subspace $\text{Fil}^0 V_{\mathcal{F}} := V_{\mathcal{F}}^{I_p}$ fixed by the inertia group I_p at p is free of rank one over \mathbf{I} ([Oht00, Corollary, page 558])). We recall the specialization of $V_{\mathcal{F}}$ at arithmetic points. If k is a positive integer and $\epsilon : \Gamma \rightarrow \overline{\mathbf{Q}}_p^{\times}$ is a finite order character, a point $Q \in \text{Spec } \mathbf{I}(\overline{\mathbf{Q}}_p)$ is called an arithmetic point of weight k and finite part ϵ if $Q|_{\Gamma} : \Gamma \rightarrow \Lambda^{\times} \xrightarrow{Q} \overline{\mathbf{Q}}_p^{\times}$ is given by $Q(x) = x^{k_Q} \epsilon_Q(x)$ for some integer $k_Q \geq 2$ and a finite order character $\epsilon_Q : \Gamma \rightarrow \overline{\mathbf{Q}}_p^{\times}$. For an arithmetic point Q , denote by k_Q the weight of Q and ϵ_Q the finite part of Q . Let $\mathfrak{X}_{\mathbf{I}}^+$ be the set of arithmetic points of \mathbf{I} . For each arithmetic point $Q \in \mathfrak{X}_{\mathbf{I}}^+$, the specialization $V_{\mathcal{F}_Q} := V_{\mathcal{F}} \otimes_{\mathbf{I}, Q} \overline{\mathbf{Q}}_p$ is the geometric p -adic Galois representation associated with the eigenform \mathcal{F}_Q of constructed by Shimura and Deligne.

1.2. Triple product L -functions. Let $\mathbf{V} = V_{\mathbf{f}} \widehat{\otimes}_{\mathcal{O}} V_{\mathbf{g}} \widehat{\otimes}_{\mathcal{O}} V_{\mathbf{h}}$ be the triple product Galois representation of rank eight over \mathcal{R} a finite extension of the three-variable Iwasawa algebra given by

$$\mathcal{R} = \mathbf{I} \widehat{\otimes}_{\mathcal{O}} \mathbf{I} \widehat{\otimes}_{\mathcal{O}} \mathbf{I}.$$

Let $\mathfrak{X}_{\mathcal{R}}^+ \subset \text{Spec } \mathcal{R}(\overline{\mathbf{Q}}_p)$ be the weight space of arithmetic points of \mathcal{R} given by

$$\mathfrak{X}_{\mathcal{R}}^+ := \{ \underline{Q} = (Q_1, Q_2, Q_3) \in (\mathfrak{X}_{\mathbf{I}}^+)^3 \mid k_{Q_1} + k_{Q_2} + k_{Q_3} \equiv 0 \pmod{2} \}.$$

For each arithmetic point $\underline{Q} = (Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathcal{R}}^+$, the specialization $\mathbf{V}_{\underline{Q}} = V_{\mathbf{f}_{Q_1}} \otimes V_{\mathbf{g}_{Q_2}} \otimes V_{\mathbf{h}_{Q_3}}$ is a p -adic geometric Galois representation of pure weight $w_{\underline{Q}} := k_{Q_1} + k_{Q_2} + k_{Q_3} - 3$. Let $\omega : (\mathbf{Z}/p\mathbf{Z})^{\times} \rightarrow \mu_{p-1}$ be the Teichmüller character. We assume that

$$(ev) \quad \psi_1 \psi_2 \psi_3 = \omega^{2a} \text{ for some } a \in \mathbf{Z}.$$

Then (ev) implies that the determinant $\det \mathbf{V} = \mathcal{X}^2 \varepsilon_{\text{cyc}}$, where ε_{cyc} is the p -adic cyclotomic character and \mathcal{X} is a \mathcal{R} -adic p -ramified Galois character with $\mathcal{X}(\mathbf{c}) = (-1)^a$ (\mathbf{c} is the complex conjugation). Note that the specialization of \mathcal{X} at \underline{Q} can be written as the product $\mathcal{X}_{\underline{Q}} = \chi_{\underline{Q}} \varepsilon_{\text{cyc}}^{-\frac{w_{\underline{Q}}+1}{2}}$ with a finite order character $\chi_{\underline{Q}}$. We consider the critical twist

$$\mathbf{V}^{\dagger} = \mathbf{V} \otimes \mathcal{X}^{-1}.$$

Then \mathbf{V}^{\dagger} is self-dual in the sense that $(\mathbf{V}^{\dagger})^{\vee}(1) = \mathbf{V}^{\dagger}$. Next we briefly recall the complex L -function associated with the specialization $\mathbf{V}_{\underline{Q}}^{\dagger}$. For each place ℓ , denote by $W_{\mathbf{Q}_{\ell}}$ the Weil-Deligne group of \mathbf{Q}_{ℓ} . To the geometric p -adic Galois representation $\mathbf{V}_{\underline{Q}}^{\dagger}$, we can associate the Weil-Deligne representation $\text{WD}_{\ell}(\mathbf{V}_{\underline{Q}}^{\dagger})$ of $W_{\mathbf{Q}_{\ell}}$ over $\overline{\mathbf{Q}}_p$ (See [Tat79, (4.2.1)] for $\ell \neq p$ and [Fon94, (4.2.3)] for $\ell = p$). Fixing an isomorphism $\iota_p : \overline{\mathbf{Q}}_p \simeq \mathbf{C}$, we define the complex L -function of $\mathbf{V}_{\underline{Q}}^{\dagger}$ by the Euler product

$$L(\mathbf{V}_{\underline{Q}}^{\dagger}, s) = \prod_{\ell < \infty} L_{\ell}(\mathbf{V}_{\underline{Q}}^{\dagger}, s)$$

of the local L -factors $L_{\ell}(\mathbf{V}_{\underline{Q}}^{\dagger}, s)$ attached to $\text{WD}_{\ell}(\mathbf{V}_{\underline{Q}}^{\dagger}) \otimes_{\overline{\mathbf{Q}}_p, \iota_p} \mathbf{C}$ ([Del79, (1.2.2)], [Tay04, page 85]). On the other hand, if we denote by $\pi_{\mathbf{f}_{Q_1}} = \otimes_v \pi_{\mathbf{f}_{Q_1}, v}$ (resp. $\pi_{\mathbf{g}_{Q_2}}, \pi_{\mathbf{h}_{Q_3}}$) the irreducible unitary cuspidal automorphic representation of $\text{GL}_2(\mathbf{A})$ associated with \mathbf{f}_{Q_1} (resp. $\mathbf{g}_{Q_2}, \mathbf{h}_{Q_3}$) and let

$$\Pi_{\underline{Q}} = \pi_{\mathbf{f}_{Q_1}} \times \pi_{\mathbf{g}_{Q_2}} \times \pi_{\mathbf{h}_{Q_3}} \otimes \chi_{\underline{Q}}^{-1}$$

be the irreducible unitary automorphic representation of $\text{GL}_2(\mathbf{A}) \times \text{GL}_2(\mathbf{A}) \times \text{GL}_2(\mathbf{A})$. Denote by $L(s, \Pi_{\underline{Q}})$ the automorphic L -function defined by Garrent, Piatetski-Shapiro and Rallis attached to the triple product $\Pi_{\underline{Q}}$. The analytic theory of $L(s, \Pi_{\underline{Q}})$ such as functional equations and analytic continuation has been explored extensively in the literature (cf. [PSR87]), and thanks to [Ram00, Theorem 4.4.1], we have

$$L\left(s + \frac{1}{2}, \Pi_{\underline{Q}}\right) = \Lambda(\mathbf{V}_{\underline{Q}}^{\dagger}, s) := \Gamma_{\mathbf{V}_{\underline{Q}}^{\dagger}}(s) \cdot L(\mathbf{V}_{\underline{Q}}^{\dagger}, s).$$

Here $\Gamma_{\mathbf{V}_Q^\dagger}(s)$ is the archimedean L -factor of \mathbf{V}_Q^\dagger and is a finite product of four classical Γ -functions (see (1.4)). Moreover, there is a positive integer $N(\mathbf{V}_Q^\dagger)$ and the root number $\varepsilon(\mathbf{V}_Q^\dagger) \in \{\pm 1\}$ such that the complete L -function $\Lambda(\mathbf{V}_Q^\dagger, s)$ satisfies the functional equation

$$\Lambda(\mathbf{V}_Q^\dagger, s) = \varepsilon(\mathbf{V}_Q^\dagger) \cdot N(\mathbf{V}_Q^\dagger)^{-s} \cdot \Lambda(\mathbf{V}_Q^\dagger, -s).$$

We thus have a good understanding of the complex analytic behavior of $L(\mathbf{V}_Q^\dagger, s)$. In this article, we shall investigate the arithmetic of the critical central value $L(\mathbf{V}_Q^\dagger, 0)$ and study the p -adic analytic behavior of the algebraic part of $L(\mathbf{V}_Q^\dagger, 0)$ viewed as a function on the weight space $\mathfrak{X}_{\mathcal{R}}^+$. Note that the rationality of the central L -values of triple product L -functions has been proved in [HK91]. It is natural to first consider the behavior of the root number $\varepsilon(\mathbf{V}_Q^\dagger)$ of \mathbf{V}_Q^\dagger (or Π_Q) over the weight space. The global root number

$$\varepsilon(\mathbf{V}_Q^\dagger) = \prod_{\ell \leq \infty} \varepsilon(\text{WD}_\ell(\mathbf{V}_Q^\dagger))$$

is defined as the product of local constants, where $\varepsilon(?)$ is the local epsilon factor attached to a Weil-Deligne representation (*cf.* [Tat79, page 21]) with respect to the standard choice of a non-trivial additive character of \mathbf{Q}_p and measures on \mathbf{Q}_p in [Del79, 5.3]. For each arithmetic point $Q \in \mathfrak{X}_{\mathcal{R}}^+$, we put

$$\Sigma^-(Q) := \left\{ \ell: \text{prime factor of } N_1 N_2 N_3 \mid \varepsilon(\text{WD}_\ell(\mathbf{V}_Q^\dagger)) = -1 \right\}.$$

It is known that there is a subset Σ^- of prime factors of N such that $\Sigma^- = \Sigma^-(Q)$ for all $Q \in \mathfrak{X}_{\mathcal{R}}^+$. For the archimedean root number, we partition the weight space $\mathfrak{X}_{\mathcal{R}}^+$ into $\mathfrak{X}_{\mathcal{R}}^f \sqcup \mathfrak{X}_{\mathcal{R}}^g \sqcup \mathfrak{X}_{\mathcal{R}}^h \sqcup \mathfrak{X}_{\mathcal{R}}^{\text{bal}}$, where $\mathfrak{X}_{\mathcal{R}}^f$ is the unbalanced range dominated by f given by

$$\mathfrak{X}_{\mathcal{R}}^f = \{(Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathcal{R}}^+ \mid k_{Q_1} + k_{Q_2} + k_{Q_3} \leq 2k_{Q_1}\}$$

($\mathfrak{X}_{\mathcal{R}}^g$ and $\mathfrak{X}_{\mathcal{R}}^h$ are defined likewise), and $\mathfrak{X}_{\mathcal{R}}^{\text{bal}}$ is the balanced range

$$\mathfrak{X}_{\mathcal{R}}^{\text{bal}} = \{(Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathcal{R}}^+ \mid k_{Q_1} + k_{Q_2} + k_{Q_3} > 2k_{Q_i} \text{ for all } i = 1, 2, 3\}.$$

The union $\mathfrak{X}_{\mathcal{R}}^{\text{unb}} := \mathfrak{X}_{\mathcal{R}}^f \sqcup \mathfrak{X}_{\mathcal{R}}^g \sqcup \mathfrak{X}_{\mathcal{R}}^h$ is called the unbalanced range. Then we know that

$$\begin{aligned} \varepsilon(\text{WD}_\infty(\mathbf{V}_Q^\dagger)) &= +1 \text{ if } Q \in \mathfrak{X}_{\mathcal{R}}^{\text{unb}}; \\ \varepsilon(\text{WD}_\infty(\mathbf{V}_Q^\dagger)) &= -1 \text{ if } Q \in \mathfrak{X}_{\mathcal{R}}^{\text{bal}}. \end{aligned}$$

1.3. The modified Euler factors at p . Let $G_{\mathbf{Q}_p}$ be the decomposition group at p . We consider the following rank four $G_{\mathbf{Q}_p}$ -invariant subspaces of \mathbf{V}_Q :

$$(1.1) \quad \begin{aligned} \text{Fil}_f \mathbf{V} &:= \text{Fil}^0 V_f \otimes V_g \otimes V_h; \\ \text{Fil}_{\text{bal}} \mathbf{V} &:= \text{Fil}^0 V_f \otimes \text{Fil}^0 V_g \otimes V_h + V_f \otimes \text{Fil}^0 V_g \otimes \text{Fil}^0 V_h + \text{Fil}^0 V_f \otimes V_g \otimes \text{Fil}^0 V_h. \end{aligned}$$

Let $\bullet \in \{f, \text{bal}\}$. Define the filtrations $\text{Fil}_\bullet^+ \mathbf{V}^\dagger := \text{Fil}_\bullet \mathbf{V} \otimes \mathcal{X}^{-1} \subset \mathbf{V}^\dagger$. The pair $(\text{Fil}_\bullet^+ \mathbf{V}^\dagger, \mathfrak{X}_{\mathcal{R}}^\bullet)$ satisfies the *Panchishkin condition* in [Gre94, page 217]) in the sense that for each arithmetic point $Q \in \mathfrak{X}_{\mathcal{R}}^\bullet$, the Hodge-Tate numbers of $\text{Fil}_\bullet^+ \mathbf{V}_Q^\dagger$ are all positive, while the Hodge-Tate numbers of $\mathbf{V}^+ / \text{Fil}_\bullet^+ \mathbf{V}_Q^\dagger$ are all non-positive.¹ Now we can define the modified p -Euler factor by

$$(1.2) \quad \mathcal{E}_p(\text{Fil}_\bullet^+ \mathbf{V}_Q^\dagger) := \frac{L_p(\text{Fil}_\bullet^+ \mathbf{V}_Q^\dagger, 0)}{\varepsilon(\text{WD}_p(\text{Fil}_\bullet^+ \mathbf{V}_Q^\dagger)) \cdot L_p(\mathbf{V}_Q^\dagger / \text{Fil}_\bullet^+ \mathbf{V}_Q^\dagger, 0)} \cdot \frac{1}{L_p(\mathbf{V}_Q^\dagger, 0)}.$$

We note that this modified p -Euler factor is precisely the product of the factor $\mathcal{L}_p^{(\rho)}(\mathbf{V}_Q^\dagger)$ in [Coa89b, page 109, (18)] and the reciprocal of the local L -factor $L_p(0, \mathbf{V}_Q^\dagger)$.

¹The Hodge-Tate number of $\mathbf{Q}_p(1)$ is one in our convention.

1.4. Hida's canonical periods. To make our interpolation formula meaningful, we must give the precise definition of periods for the motive \mathbf{V}_Q^\dagger . We begin by recalling Hida's canonical period of a \mathbf{I} -adic primitive cuspidal Hida family \mathcal{F} of tame conductor $N_{\mathcal{F}}$. Let $\mathfrak{m}_{\mathbf{I}}$ be the maximal ideal of \mathbf{I} . For a subset Σ of the support of $N_{\mathcal{F}}$, we consider the following

Hypothesis (CR, Σ). The residual Galois representation $\bar{\rho}_{\mathcal{F}} := \rho_{\mathcal{F}} \pmod{\mathfrak{m}_{\mathbf{I}}} : G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\bar{\mathbb{F}}_p)$:

- (1) $\bar{\rho}_{\mathcal{F}}$ is absolutely irreducible,
- (2) $\bar{\rho}_{\mathcal{F}}$ is p -distinguished,
- (3) if $\ell \in \Sigma$ and $\ell \equiv 1 \pmod{p}$, then $\bar{\rho}_{\mathcal{F}}$ is ramified at ℓ .

When $\Sigma = \emptyset$ is the empty set, we shall simply write (CR) for (CR, \emptyset). Recall that $\rho_{\mathcal{F}}$ is p -distinguished if the semi-simplification of the restriction of the residual Galois representation $\rho_{\mathcal{F}} \pmod{\mathfrak{m}_{\mathbf{I}}}$ to the decomposition at p is a sum of two characters $\chi_{\mathcal{F}}^+ \oplus \chi_{\mathcal{F}}^-$ with $\chi_{\mathcal{F}}^+ \not\equiv \chi_{\mathcal{F}}^- \pmod{\mathfrak{m}_{\mathbf{I}}}$. Suppose that \mathcal{F} satisfies (CR). The local component of the universal cuspidal ordinary Hecke algebra corresponding to \mathcal{F} is known to be Gorenstein by [MW86, Prop.2, §9] and [Wil95, Corollary 2, page 482], and with this Gorenstein property, Hida proved in [Hid88a, Theorem 0.1] that the congruence module for \mathcal{F} is isomorphic to $\mathbf{I}/(\eta_{\mathcal{F}})$ for some non-zero element $\eta_{\mathcal{F}} \in \mathbf{I}$. Moreover, for any arithmetic point $Q \in \mathfrak{X}_{\mathbf{I}}^+$, the specialization $\eta_{\mathcal{F}_Q} = Q(\eta_{\mathcal{F}})$ generates the congruence ideal of \mathcal{F}_Q . We denote by \mathcal{F}_Q° the normalized newform of weight k_Q , conductor $N_Q = N_{\mathcal{F}} p^{n_Q}$ with nebentypus χ_Q corresponding to \mathcal{F}_Q . There is a unique decomposition $\chi_Q = \chi'_Q \chi_{Q,(p)}$, where χ'_Q and $\chi_{Q,(p)}$ are Dirichlet characters modulo $N_{\mathcal{F}}$ and p^{n_Q} respectively. Let $\alpha_Q = \mathbf{a}(p, \mathcal{F}_Q)$. Define the modified Euler factor $\mathcal{E}_p(\mathcal{F}_Q, \mathrm{Ad})$ for adjoint motive of \mathcal{F}_Q by

$$\mathcal{E}_p(\mathcal{F}_Q, \mathrm{Ad}) = \alpha_Q^{-2n_Q} \begin{cases} (1 - \alpha_Q^{-2} \chi_Q(p) p^{k_Q})(1 - \alpha_Q^{-2} \chi_Q(p) p^{k_Q-1}) & \text{if } n_Q = 0, \\ -1 & \text{if } n_Q = 1, \chi_{Q,(p)} = 1 \text{ (so } k_Q = 2), \\ \mathfrak{g}(\chi_{Q,(p)}) \chi_{Q,(p)}(-1) & \text{if } n_Q > 0, \chi_{Q,(p)} \neq 1. \end{cases}$$

Here $\mathfrak{g}(\chi_{Q,(p)})$ is the standard Gauss sum. Fixing a choice of the generator $\eta_{\mathcal{F}}$ and letting $\|\mathcal{F}_Q^\circ\|_{\Gamma_0(N_Q)}^2$ be the usual Petersson norm of \mathcal{F}_Q° , we define the *canonical period* $\Omega_{\mathcal{F}_Q}$ of \mathcal{F} at Q by

$$(1.3) \quad \Omega_{\mathcal{F}_Q} := 2^{k_Q} \cdot \|\mathcal{F}_Q^\circ\|_{\Gamma_0(N_Q)}^2 \cdot \frac{\mathcal{E}_p(\mathcal{F}_Q, \mathrm{Ad})}{\eta_{\mathcal{F}_Q}} \in \mathbf{C}^\times.$$

By [Hid16, Corollary 6.24, Theorem 6.28], one can show that for each arithmetic point Q , up to a p -adic unit, the period $\Omega_{\mathcal{F}_Q}$ is equal to the product of the plus/minus canonical period $\Omega(+; \mathcal{F}_Q^\circ) \Omega(-; \mathcal{F}_Q^\circ)$ introduced in [Hid94, page 488].

1.5. Definitions of Γ -factors and an exceptional finite set Σ_{exc} . We recall the definition of Γ -factors of \mathbf{V}_Q^\dagger following the recipe in [Del79]:

$$(1.4) \quad \Gamma_{\mathbf{V}_Q^\dagger}(s) := \begin{cases} \Gamma_{\mathbf{C}}(s + \frac{w_Q+1}{2}) \Gamma_{\mathbf{C}}(s+1 - k_{Q_1}^*) \Gamma_{\mathbf{C}}(s + k_{Q_2}^*) \Gamma_{\mathbf{C}}(s + k_{Q_3}^*) & \text{if } Q \in \mathfrak{X}_{\mathcal{R}}^f; \\ \Gamma_{\mathbf{C}}(s + \frac{w_Q+1}{2}) \Gamma_{\mathbf{C}}(s + k_{Q_1}^*) \Gamma_{\mathbf{C}}(s + k_{Q_2}^*) \Gamma_{\mathbf{C}}(s + k_{Q_3}^*) & \text{if } Q \in \mathfrak{X}_{\mathcal{R}}^{\mathrm{bal}}. \end{cases}$$

Here $\Gamma_{\mathbf{C}}(s) = 2(2\pi)^{-s} \Gamma(s)$ and

$$k_{Q_i}^* = \frac{k_{Q_1} + k_{Q_2} + k_{Q_3}}{2} - k_{Q_i}, \quad i = 1, 2, 3.$$

For each prime ℓ , let $\tau_{\mathbf{Q}_{\ell^2}}$ be the unique unramified quadratic character of \mathbf{Q}_{ℓ}^\times . Let $(f, g, h) = (\mathbf{f}_{Q_1}, \mathbf{g}_{Q_2}, \mathbf{h}_{Q_3})$ be the specialization of \mathbf{F} at Q and put

$$\begin{aligned} \Sigma_{\mathbf{fg}}^{\mathrm{sc}} &= \{\ell \mid \pi_{f,\ell} \text{ and } \pi_{g,\ell} \text{ are supercuspidal; } \pi_{h,\ell} \text{ is spherical}\}; \\ \Sigma_{\mathbf{fg}} &= \{\ell \in \Sigma_{\mathbf{fg}}^{\mathrm{sc}} \mid \pi_{f,\ell} \simeq \pi_{f,\ell} \otimes \tau_{\mathbf{Q}_{\ell^2}} \simeq \pi_{g,\ell}^\vee \otimes \sigma \text{ for some } \sigma \text{ unramified character}\}. \end{aligned}$$

Define $\Sigma_{\mathbf{fh}}$ and $\Sigma_{\mathbf{fg}}$ likewise. We introduce the finite set

$$(1.5) \quad \Sigma_{\mathrm{exc}} = \Sigma_{\mathbf{gh}} \sqcup \Sigma_{\mathbf{fh}} \sqcup \Sigma_{\mathbf{fg}}.$$

It is known that this set Σ_{exc} does not depend on any particular choice of the specializations of $(\mathbf{f}, \mathbf{g}, \mathbf{h})$.

1.6. **Statement of the main results.** We impose the following technical assumption:

(sf) $\gcd(N_1, N_2, N_3)$ is square-free.

Our first result is the construction of the unbalanced p -adic triple product L -functions:

Theorem A. *In addition to (ev) and (sf), we further suppose that*

- (1) $\Sigma^- = \emptyset$,
- (2) \mathbf{f} satisfies (CR).

Fix a generator $\eta_{\mathbf{f}}$ of the congruence ideal of \mathbf{f} . There exists a unique element $\mathcal{L}_{\mathbf{F}}^{\mathbf{f}} \in \mathcal{R}$ such that for every $\underline{Q} = (Q_3, Q_2, Q_1) \in \mathfrak{X}_{\mathcal{R}}^{\mathbf{f}}$ in the unbalanced range dominated by \mathbf{f} , we have

$$(\mathcal{L}_{\mathbf{F}}^{\mathbf{f}}(\underline{Q}))^2 = \Gamma_{\mathbf{V}_{\underline{Q}}^+}(0) \cdot \frac{L(\mathbf{V}_{\underline{Q}}^{\dagger}, 0)}{(-1)^{k_{Q_1}} \Omega_{\mathbf{f}_{Q_1}}^2} \cdot \mathcal{E}_p(\text{Fil}_{\mathbf{f}}^+ \mathbf{V}_{\underline{Q}}^{\dagger}) \cdot \prod_{\ell \in \Sigma_{\text{exc}}} (1 + \ell^{-1}).$$

This p -adic L -function $\mathcal{L}_{\mathbf{F}}^{\mathbf{f}}$ is unique up to a choice of generators of the congruence ideal of \mathbf{f} , i.e. it is unique up to a unit in \mathbf{I} , but the ratio $\mathcal{L}_{\mathbf{F}}^{\mathbf{f}}/\eta_{\mathbf{f}}$ is a genuine p -adic L -function. From Theorem A, one also obtains different p -adic L -functions $\mathcal{L}_{\mathbf{F}}^{\mathbf{g}}$ and $\mathcal{L}_{\mathbf{F}}^{\mathbf{h}}$ which interpolate central L -values at $\mathfrak{X}_{\mathcal{R}}^{\mathbf{g}}$ and $\mathfrak{X}_{\mathcal{R}}^{\mathbf{h}}$ respectively. These p -adic L -functions $\mathcal{L}_{\mathbf{F}}^{\mathbf{f}}$, $\mathcal{L}_{\mathbf{F}}^{\mathbf{g}}$ and $\mathcal{L}_{\mathbf{F}}^{\mathbf{h}}$ are called *unbalanced* p -adic triple product L -functions as they interpolate a square root of the critical central L -values of the triple product L -function $L(\mathbf{V}_{\underline{Q}}^{\dagger}, s)$ for $\underline{Q} \in \mathfrak{X}_{\mathcal{R}}^{\text{unb}}$ at the unbalanced range; from the interpolation formula, these p -adic L -functions are distinguished by the choices of the modified Euler factor at p and the complex periods. In the literature, the one-variable unbalanced p -adic triple product L -functions were first constructed by Harris and Tilouine in [HT01b] (when $N_1 = N_2 = N_3 = 1$). Darmon and Rotger in [DR14] extended the method in [HT01b] to construct a three-variable power series interpolating the global trilinear period of a triple of Hida families and proved the interpolation formulae at the *balanced range*, which is in connection with the p -adic Abel-Jacobi image of diagonal cycles in a triple product of modular curves. This is a p -adic analogue of the classical Gross-Zagier formula and has obtained very significant arithmetic application to certain equivariant BSD conjectures in [DR17]. On the other hand, it is well known that the relation of the interpolation at the unbalanced range to central L -values is suggested by the main identity of Harris and Kudla [HK91], or in general, Ichino's formula [Ich08], but the interpolation formulae at the unbalanced range in the literature are not precise enough for more refined arithmetic applications such as the formulation of corresponding Iwasawa-Greenberg main conjecture. Therefore, Theorem A complements the literature by providing a precise relation of the values of p -adic triple product L -functions at all arithmetic points in the unbalanced range to central L -values of the complex triple product L -functions.

Our main motivation is to use Theorem A to prove the factorization of p -adic triple product L -functions into a product of anticyclotomic p -adic L -functions. For example, according to Theorem A, if \mathbf{g} and \mathbf{h} are primitive Hida families of CM forms associated with some imaginary quadratic field, then $\mathcal{L}_{\mathbf{F}}^{\mathbf{f}}$ is a product of two anticyclotomic p -adic L -functions for modular forms constructed in [BD96] and [CH16]; in contrast, if \mathbf{f} and \mathbf{g} are primitive Hida families of CM forms, then $\mathcal{L}_{\mathbf{F}}^{\mathbf{f}}$ is a product of two anticyclotomic p -adic L -functions in [BDP13] divided by some Katz p -adic L -function. The latter gives a strengthening of [DLR15, Theorem 3.8] and [Col16]. With this factorization, we can easily show that the anticyclotomic p -adic L -functions in the definite case can be recovered by the Euler system of generalized Kato classes [DR17] (See Remark 8.2) and obtain a new proof of the anticyclotomic exceptional zero conjecture for elliptic curves. In an ongoing project with F. Castella, we will explore this Euler system construction of the anticyclotomic p -adic L -functions and deduce Rubin's formulas for generalized Kato classes in the rank two case.

Next we state our second result about the balanced p -adic triple product L -functions. Let

$$N = \text{lcm}(N_1, N_2, N_3); \quad N^- = \prod_{\ell \in \Sigma^-} \ell.$$

Theorem B. *In addition to (ev) and (sf), we further suppose that*

- (1) $\#(\Sigma^-)$ is odd,
- (2) \mathbf{f}, \mathbf{g} and \mathbf{h} satisfy (CR, Σ^-),
- (3) $N = N^+ N^-$ with $\gcd(N^+, N^-) = 1$.

Then there exists a unique element $\mathcal{L}_{\mathbf{F}}^{\text{bal}} \in \mathcal{R}$ satisfies the following interpolation property: for any arithmetic point $\underline{Q} \in \mathfrak{X}_{\mathcal{R}}^{\text{bal}}$, we have

$$(\mathcal{L}_{\mathbf{F}}^{\text{bal}}(\underline{Q}))^2 = \Gamma_{\mathbf{V}_{\underline{Q}}^{\dagger}}(0) \cdot \frac{L(\mathbf{V}_{\underline{Q}}^{\dagger}, 0)}{\Omega_{\mathbf{f}_{Q_1}} \Omega_{\mathbf{g}_{Q_2}} \Omega_{\mathbf{h}_{Q_3}}} \cdot \mathcal{E}_p(\text{Fil}_{\text{bal}}^+ \mathbf{V}_{\underline{Q}}^{\dagger}) \cdot \prod_{\ell \in \Sigma_{\text{exc}}} (1 + \ell^{-1}).$$

We must mention that the p -adic interpolation of global trilinear period integrals attached to a triple of p -adic families of modular forms in the balanced range was first investigated by Greenberg and Seveso in a pioneering work [GS16]. Our construction is ostensibly different from theirs for their method heavily relies on the theory of Ash-Stevens while our approach is built on classical Hida theory developed in [Hid88b]. Indeed, their method treats more general setting, namely they do not restrict to the ordinary case, while our approach is more well-suited for the future investigation on the arithmetic of the balanced p -adic L -functions such as the μ -invariants and the Iwasawa-Greenberg main conjecture. The situation is more or less similar to the two different constructions of two-variable p -adic L -functions for Hida families given by Greenberg-Stevens and Mazur-Kitagawa. In any case, it is definitely very interesting to compare these two different approaches in the ordinary case.

Remark 1.1. We discuss the exceptional zero phenomenon for the balanced p -adic L -functions. By the Ramanujan conjecture, the modified p -Euler factor $\mathcal{E}_p(\text{Fil}_{\text{bal}}^+ \mathbf{V}_{\underline{Q}}^{\dagger})$ never vanishes unless either of $\mathbf{f}_{Q_1}, \mathbf{g}_{Q_2}, \mathbf{h}_{Q_3}$ is special at p . For example, suppose that $\mathbf{F} = (\mathbf{f}, \mathbf{g}, \mathbf{h})$ is the triple of primitive Hida families passing through the p -stabilized newforms (f_1, f_2, f_3) attached to elliptic curves (E_1, E_2, E_3) over \mathbf{Q} at the weight two specialization \underline{Q} . Let $\alpha_i = \mathbf{a}(p, f_i)$ be the p -th Fourier coefficient of f_i for $i = 1, 2, 3$. Assume E_1 is semi-stable at p (i.e. $\alpha_1 = \pm 1$). Then the formula of the modified p -Euler factor reads

$$\mathcal{E}_p(\text{Fil}_{\text{bal}}^+ \mathbf{V}_{\underline{Q}}^{\dagger}) = \begin{cases} (1 - \alpha_1 \alpha_2 \alpha_3)^3 & \text{if } E_2 \text{ and } E_3 \text{ are semi-stable at } p, \\ p \cdot \alpha_3^{-2} (1 - \frac{\alpha_3}{\alpha_1 \alpha_2})^2 (1 - \frac{\alpha_1}{\alpha_2 \alpha_3})^2 & \text{otherwise.} \end{cases}$$

We thus conclude that $\mathcal{L}_{\mathbf{F}}^{\text{bal}}$ possesses an exceptional zero at \underline{Q} when either (i) E_2 and E_3 are semi-stable at p and $\alpha_1 \alpha_2 \alpha_3 = 1$ or (ii) E_2 and E_3 has good ordinary reduction at p and $\alpha_2 = \alpha_3 \alpha_1$. In the case (i), we even have the vanishing of the central value $L(\mathbf{V}_{\underline{Q}}^{\dagger}, 0) = L(E_1 \times E_2 \times E_3, 2) = 0$ as the global root number

$$\varepsilon(\mathbf{V}_{\underline{Q}}^{\dagger}) = \varepsilon(\text{WD}_p(\mathbf{V}_{\underline{Q}}^{\dagger})) = -\alpha_1 \alpha_2 \alpha_3 = -1,$$

so one might speculate about a p -adic Gross-Zaiger formula relating certain ‘‘second partial derivatives’’ of $\mathcal{L}_{\mathbf{F}}^{\text{bal}}$ at \underline{Q} to the p -adic Abel-Jacobi image of diagonal cycle in the Shimura curve X_{N^+, pN^-} attached to the quaternion algebra ramified precisely at pN^- as [BD07, Theorem 1]. We hope to come back to this question in the near future.

1.7. An outline of the proof. The construction of the unbalanced p -adic L -function is based on Hida’s p -adic Rankin-Selberg convolution (cf. [Hid93]). Denote by $e\mathbf{S}(N, \chi, \mathbf{I}) \subset \mathbf{I}[[q]]$ the space of ordinary \mathbf{I} -adic cusp forms with tame nebentypus χ and by $\mathbf{T}(N, \chi, \mathbf{I})$ the universal ordinary cuspidal Hecke algebra. Decompose the tame nebentypus ψ_1 of \mathbf{f} into a product of Dirichlet characters $\psi_{1,(p)}$ and $\psi_1^{(p)}$ modulo p and N_1 respectively and let $\chi := \psi_{1,(p)} \psi_1^{(p)}$. Let $\check{\mathbf{f}} \in e\mathbf{S}(N_1, \chi, \mathbf{I})$ be the primitive Hida family of \mathbf{f} twisted by $\psi_1^{(p)}$ and let $1_{\check{\mathbf{f}}} \in \mathbf{T}(N_1, \chi, \mathbf{I}) \otimes_{\mathbf{I}} \text{Frac } \mathbf{I}$ be the idempotent corresponding to $\check{\mathbf{f}}$. By the definition of congruence ideals, one can verify that $\eta_{\mathbf{f}} \cdot 1_{\check{\mathbf{f}}}$ indeed belongs to $\mathbf{T}(N, \chi, \mathbf{I})$. In §3.6 (3.8), we construct an auxiliary \mathcal{R} -adic modular form $\mathbf{H}^{\text{aux}} \in e\mathbf{S}(N, \chi, \mathbf{I}) \otimes_{\mathbf{I}, i_1} \mathcal{R} \subset \mathcal{R}[[q]]$, where $i_1 : \mathbf{I} \rightarrow \mathcal{R}$ is the homomorphism $a \mapsto a \otimes 1 \otimes 1$, and then the unbalanced p -adic L -function is defined to be

$$\mathcal{L}_{\mathbf{F}}^{\mathbf{f}} := \text{the first Fourier coefficient of } \eta_{\mathbf{f}} \cdot 1_{\check{\mathbf{f}}} \text{Tr}_{N/N_1}(\mathbf{H}^{\text{aux}}) \in \mathcal{R},$$

where $\text{Tr}_{N/N_1} : e\mathbf{S}(N, \chi, \mathbf{I}) \rightarrow e\mathbf{S}(N_1, \chi, \mathbf{I})$ is the usual trace map.

In the balanced case, Hida theory for definite quaternion algebras plays an important role. Let D be the definite quaternion algebra over \mathbf{Q} of the absolute discriminant N^- , and for each positive integer m , let \tilde{X}_m be the definite Shimura curve of level $\Gamma_1(p^n N)$ associated with D as described in [LV11, §2.1]. These are curves of genus zero equipped with a natural finite covering map $\tilde{\alpha}_m : \tilde{X}_m \rightarrow \tilde{X}_{m-1}$. We let $J_m = \text{Pic } \tilde{X}_m \otimes_{\mathbf{Z}} \mathbf{Z}_p$ and let $J_{\infty} := \varprojlim_{n \rightarrow \infty} J_m$ be the inverse limit induced by $\tilde{\alpha}_m$. Then J_{∞} is a Λ -module

with Hecke action, and its ordinary part J_∞^{ord} is equipped with the action of the Σ^- -new quotient of the universal ordinary cuspidal Hecke algebra of level $\Gamma_1(Np^\infty)$. The \mathbf{I} -module $e\mathbf{S}^D(N, \mathbf{I}) := \text{Hom}_\Lambda(J_\infty^{\text{ord}}, \mathbf{I})$ is called the space of Hida families of definite quaternionic forms. Due to the lack of q -expansions, we do not have the notion of primitive Hida families on definite quaternion algebras. Nonetheless, using the idea of Pollack and Weston [PW11] and Hida theory, for a primitive Hida family \mathcal{F} satisfying (CR, Σ^-), it can be shown that there exists Hecke eigenform $\mathcal{F}^D \in e\mathbf{S}^D(N, \mathbf{I})$, unique up to a unit in \mathbf{I} , characterized by the following properties (i) \mathcal{F}^D shares the same Hecke eigenvalues with \mathcal{F} ; (ii) \mathcal{F}^D is non-zero modulo $\mathfrak{m}_\mathbf{I}$ (Theorem 4.5). We shall call \mathcal{F}^D the primitive Jacquet-Langlands lift of \mathcal{F} . Let $\mathbf{J}_m^{\text{ord}} := J_m^{\text{ord}} \widehat{\otimes}_{\mathcal{O}} J_m^{\text{ord}} \widehat{\otimes}_{\mathcal{O}} J_m^{\text{ord}}$ and $\mathbf{J}_\infty^{\text{ord}} = \varprojlim_{m \rightarrow \infty} \mathbf{J}_m^{\text{ord}}$. Hence, with the assumption (2) in Theorem B, we obtain the primitive Jacquet-Langlands lift $\mathbf{F}^D = \mathbf{f}^D \boxtimes \mathbf{g}^D \boxtimes \mathbf{h}^D \in \text{Hom}(\mathbf{J}_\infty^{\text{ord}}, \mathcal{R})$. On the other hand, in Definition 4.6, we construct a collection of regularized diagonal cycles Δ_m^\dagger in $\mathbf{J}_m^{\text{ord}}$ which are compatible with respect to $\tilde{\alpha}_m$ and thus get the *big diagonal cycle* $\Delta_\infty^\dagger := \varprojlim_{m \rightarrow \infty} \Delta_m^\dagger \in \mathbf{J}_\infty^{\text{ord}}$. In order to achieve the optimal integrality of p -adic L -functions, we actually take a modification $\mathbf{F}^{D*} \in \text{Hom}(\mathbf{J}_\infty^{\text{ord}}, \mathcal{R})$ of \mathbf{F}^D in Definition 4.8, and then define the balanced p -adic L -function

$$\Theta_{\mathbf{F}^D} := \mathbf{F}^{D*}(\Delta_\infty^\dagger) \in \mathcal{R}$$

to be the value of the modified \mathbf{F}^{D*} at Δ_∞^\dagger . This p -adic L -function $\Theta_{\mathbf{F}^D}$ is an analogue of theta elements à la Bertolini and Darmon ([BD96]) in the triple product setting.

To obtain the interpolation formula in Theorem A and B, we first prove that the interpolation $\mathcal{L}_\mathbf{F}^f(\underline{Q})$ at $\underline{Q} \in \mathfrak{X}_\mathcal{R}^f$ (resp. $\mathcal{L}_{f, \Sigma^-}(\underline{Q})$ at $\underline{Q} \in \mathfrak{X}_\mathcal{R}^{\text{bal}}$) is given by the global trilinear period integral of certain automorphic forms in the cuspidal automorphic representation $\Pi_{\underline{Q}}$ of $\text{GL}_2(\mathbf{A}_E)$ (resp. the automorphic representation $\Pi_{\underline{Q}}^D$ of $(D \otimes \mathbf{A}_E)^\times$ via the Jacquet-Langlands transfer), where $E = \mathbf{Q} \oplus \mathbf{Q} \oplus \mathbf{Q}$ is the split étale cubic \mathbf{Q} -algebra (See Proposition 3.7 and 4.9). Thanks to Ichino's formula in [Ich08], we can show that the square of this global trilinear period integral is a product of the central L -value $L(1/2, \Pi_{\underline{Q}})$ and certain local zeta integrals $I_v(\phi_v^* \otimes \phi_v^*)$ (See §3.8.2 for definitions), which we shall call *local Ichino integrals* in the introduction. Hence, the proof of the interpolation formulae boils down to the determination of the values of these local Ichino integrals. In the literature, local Ichino integrals were only computed for some special cases [III10], [NPS14] and [Hu17]. Local Ichino integrals at the real place are completely determined in a recent work [CC17], but the explicit calculation of local Ichino integrals at non-archimedean places in the generality we need is a highly laborious task and occupies a substantial part of this paper. The key ingredient in our computation is Proposition 5.1, a generalization of [MV10, Lemma 3.4.2] by removing several restrictive conditions therein, which reduces the calculation of local Ichino integrals to that of certain local Rankin-Selberg integrals in [GJ78, (1.1.3)]. With local theory of L -functions on $\text{GL}(2) \times \text{GL}(2)$ developed by Jacquet in [Jac72], we are able to work out the calculation of local Rankin-Selberg integrals under (sf) and certain *minimal hypothesis* (See Hypothesis 6.1). This minimal hypothesis, roughly speaking, requires \mathbf{F} to be minimal in the sense that \mathbf{F} has the minimal conductor among Dirichlet twists. It turns out that the modified p -Euler factor $\mathcal{E}_p(\text{Fil}_\bullet^+ \mathbf{V}_{\underline{Q}}^\dagger)$ is given by the p -adic Ichino integral while local Ichino integrals at ramified places ℓ only contributes p -adic units if $\ell \notin \Sigma_{\text{exc}}$ or $(1 + \ell^{-1})$ if $\ell \in \Sigma_{\text{exc}}$. Therefore, taking a suitable Dirichlet twist $\mathbf{F}' = (\mathbf{f} \otimes \chi_1, \mathbf{g} \otimes \chi_2, \mathbf{h} \otimes \chi_3)$ with $\chi_1 \chi_2 \chi_3 = 1$ which satisfies the minimal hypothesis, we obtain the desired p -adic L -functions

$$\mathcal{L}_\mathbf{F}^f := \mathcal{L}_{\mathbf{F}'}^f \otimes \chi^1; \quad \mathcal{L}_\mathbf{F}^{\text{bal}} := \Theta_{\mathbf{F}'}^D.$$

The interpolation formulae is a direct consequence of the explicit evaluation of local Ichino integrals and the comparison between the canonical periods of \mathbf{F} and its Dirichlet twist \mathbf{F}' . The method of this paper has been extended by Ishikawa in [Ish17] to construct p -adic twisted triple product L -functions attached a Hida family of Hilbert modular form over a real quadratic field and a Hida family of elliptic modular forms.

This paper is organized as follows. In §2, we recall basic definitions and facts about classical elliptic modular forms and automorphic forms on $\text{GL}_2(\mathbf{A})$. In §3, we give the construction of the unbalanced p -adic triple product L -functions $\mathcal{L}_\mathbf{F}^f$. The key items used in the construction of \mathbf{H}^{aux} , the test Λ -adic forms \mathbf{g}^* and \mathbf{h}^* , are introduced in Definition 3.3. The main formula is derived in Corollary 3.13, where we show the interpolation of the square of $\mathcal{L}_\mathbf{F}^f$ at the unbalanced range is the product of the central L -value of the triple product L -function and local Ichino integrals at the prime p and ramified primes. In §4, we consider the balanced case. We review Hida's theory for definite quaternionic forms in §4.4 and §4.5. In particular, we present a slightly explicit version of the control theorem in Theorem 4.2 and explain the notion of primitive

Jacquet-Langlands lifts in Theorem 4.5. The construction of the big diagonal cycle Δ_∞^\dagger and the balanced p -adic L -functions are given in §4.6 and §4.7. The relation between the interpolation of the square of our balanced p -adic L -functions and the product of the central L -value and local Ichino integrals is given in Corollary 4.13. In §5, we prepare the tools for the computation of local Ichino integrals and carry out the calculations at the p -adic place, and in §6, we elaborate the calculation of local Ichino integrals at ramified primes. In particular, we show in §6.6 that the local Ichino integrals at ramified places can be interpolated into a unit in the ring \mathcal{R} of three-variable Iwasawa functions. In §7, we show that the canonical periods of a primitive Hida family and its Dirichlet twists are equal up to a unit in \mathbf{I} by the method of level-raising and prove the main results (Theorem 7.1). Finally, we prove the factorization of anticyclotomic p -adic L -functions and give applications in §8.

Notation. The following notations will be used frequently throughout the paper. Let \mathbf{A} be the ring of adèles of \mathbf{Q} . If v is a place of \mathbf{Q} , let \mathbf{Q}_v be the completion of \mathbf{Q} with respect to v , and for $a \in \mathbf{A}^\times$, let $a_v \in \mathbf{Q}_v^\times$ be the v -component of a . Denote by $|\cdot|_v$ (or simply $|\cdot|$ if there is no fear of confusion) the absolute value on \mathbf{Q}_v normalized so that $|\cdot|$ is the usual absolute value on \mathbf{R} if $v = \infty$ and $|\ell|_\ell = \ell^{-1}$ if $v = \ell$ is finite. Let $|\cdot|_{\mathbf{A}}$ be the absolute value on \mathbf{A}^\times given by $|a|_{\mathbf{A}} = \prod_v |a_v|_v$. Let $\zeta_v(s)$ be the usual local zeta function of \mathbf{Q}_v . Namely,

$$\zeta_\infty(s) = \pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right); \quad \zeta_\ell(s) = (1 - \ell^{-s})^{-1}.$$

Define the global zeta function $\zeta_{\mathbf{Q}}(s)$ of \mathbf{Q} by $\zeta_{\mathbf{Q}}(s) = \prod_v \zeta_v(s)$. In particular, $\zeta_{\mathbf{Q}}(2) = \pi/6$.

For a prime ℓ , let $v_\ell : \mathbf{Q}_\ell^\times \rightarrow \mathbf{C}^\times$ be the valuation normalized so that $v_\ell(\ell) = 1$. We shall regard \mathbf{Q}_ℓ and \mathbf{Q}_ℓ^\times as subgroups of \mathbf{A} and \mathbf{A}^\times in a natural way. To avoid possible confusion, denote $\varpi_\ell = (\varpi_{\ell,v}) \in \mathbf{A}^\times$ by the idele defined by $\varpi_{\ell,\ell} = \ell$ and $\varpi_{\ell,v} = 1$ if $v \neq \ell$.

Let $\psi_{\mathbf{Q}} : \mathbf{A}/\mathbf{Q} \rightarrow \mathbf{C}^\times$ be the additive character with the archimedean component $\psi_{\mathbf{R}}(x) = \exp(2\pi\sqrt{-1}x)$ and let $\psi_{\mathbf{Q}_\ell} : \mathbf{Q}_\ell \rightarrow \mathbf{C}^\times$ be the local component of $\psi_{\mathbf{Q}}$ at ℓ .

If R is a commutative ring and $G = \mathrm{GL}_2(R)$, we denote by ρ the right translation of G on the space of \mathbf{C} -valued functions on G : $\rho(g)f(g') = f(g'g)$ and by $\mathbf{1} : G \rightarrow \mathbf{C}$ the constant function $\mathbf{1}(g) = 1$. For a function $f : G \rightarrow \mathbf{C}$ and a character $\chi : R^\times \rightarrow \mathbf{C}^\times$, let $f \otimes \chi : G \rightarrow \mathbf{C}$ denote the function $f \otimes \chi(g) = f(g)\chi(\det g)$.

Let $G_{\mathbf{Q}} = \mathrm{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ be the absolute Galois group of \mathbf{Q} and if $\chi : (\mathbf{Z}/N\mathbf{Z})^\times \rightarrow \mathbf{C}^\times$ is Dirichlet character modulo N , denote by $c_\ell(\chi) \leq v_\ell(N)$ the ℓ -exponent of the conductor of χ . We shall identify χ with the Galois character $\chi : G_{\mathbf{Q}} \rightarrow \mathbf{C}^\times$ via class field theory.

If $\omega : \mathbf{Q}^\times \backslash \mathbf{A}^\times \rightarrow \overline{\mathbf{Q}}^\times$ is a finite order Hecke character, we denote by $\omega_\ell : \mathbf{Q}_\ell^\times \rightarrow \mathbf{C}^\times$ the local component of ω at ℓ . On the other hand, we write $\omega = \omega_{(\ell)}\omega^{(\ell)}$, where $\omega_{(\ell)}$ and $\omega^{(\ell)}$ are finite order Hecke characters of conductor ℓ -power and of prime-to- ℓ conductor respectively.

2. CLASSICAL MODULAR FORMS AND AUTOMORPHIC FORMS

In this section, we recall basic definitions and facts about classical elliptic modular forms and automorphic forms on $\mathrm{GL}_2(\mathbf{A})$. The main purpose of this section is to set up the notation and introduce some Hecke operators on the space of automorphic forms which will be frequently used in the construction of p -adic L -functions.

2.1. Classical modular forms. Let $C^\infty(\mathfrak{H})$ be the space of \mathbf{C} -valued smooth functions on the upper half complex plane \mathfrak{H} . Let k be any integer. Let $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2^+(\mathbf{R})$ act on $z \in \mathfrak{H}$ by $\gamma(z) = \frac{az+b}{cz+d}$, and for $f = f(z) \in C^\infty(\mathfrak{H})$, define

$$f|_k \gamma(z) := f(\gamma(z))(cz+d)^{-k} (\det \gamma)^{\frac{k}{2}}.$$

Recall that the Maass-Shimura differential operators δ_k and ε on $C^\infty(\mathfrak{H})$ are given by

$$\delta_k = \frac{1}{2\pi\sqrt{-1}} \left(\frac{\partial}{\partial z} + \frac{k}{2\sqrt{-1}y} \right) \text{ and } \varepsilon = -\frac{1}{2\pi\sqrt{-1}} y^2 \frac{\partial}{\partial \bar{z}} \quad (y = \mathrm{Im}(z))$$

(cf. [Hid93, (1a, 1b) page 310]). Let N be a positive integer and $\chi : (\mathbf{Z}/N\mathbf{Z})^\times \rightarrow \mathbf{C}^\times$ be a Dirichlet character modulo N . Let m be a non-negative integer. Denote by $\mathcal{N}_k^{[m]}(N, \chi)$ the space of nearly holomorphic modular

forms of weight k , level N and character χ , consisting of slowly increasing functions $f \in C^\infty(\mathfrak{H})$ such that $\varepsilon^{m+1}f = 0$ and

$$f|_k \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \chi(d)f \quad \text{for } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N)$$

(cf. [Hid93, page 314]). Let $\mathcal{N}_k(N, \chi) = \bigcup_{r=0}^\infty \mathcal{N}_k^{[m]}(N, \chi)$. (cf. [Hid93, (1a), page 310]) By definition, $\mathcal{N}_k^{[0]}(N, \chi) = \mathcal{M}_k(N, \chi)$ is the space of classical holomorphic modular forms of weight k , level N and character χ . Denote by $\mathcal{S}_k(N, \chi)$ the space of holomorphic cusp forms in $\mathcal{M}_k(N, \chi)$. Let $\delta_k^m = \delta_{k+2m-2} \cdots \delta_{k+2} \delta_k$. If $f \in \mathcal{N}_k(N, \chi)$ is a nearly holomorphic modular form of weight k , then $\delta_k^m f \in \mathcal{N}_{k+2m}(N, \chi)$ has weight $k+2m$ ([Hid93, page 312]). For a positive integer d , define

$$V_d f(z) = d \cdot f(dz); \quad \mathbf{U}_d f(z) = \frac{1}{d} \sum_{j=0}^{d-1} f\left(\frac{z+j}{d}\right),$$

and recall that the classical Hecke operators T_ℓ for primes $\ell \nmid N$ are given by

$$T_\ell f = \mathbf{U}_\ell f + \chi(\ell) \ell^{k-2} V_\ell f.$$

We say $f \in \mathcal{N}_k(N, \chi)$ is a *Hecke eigenform* if f is an eigenfunction of the all Hecke operators T_ℓ for $\ell \nmid N$ and the operators \mathbf{U}_ℓ for $\ell \mid N$.

If $f \in \mathcal{M}_k(N, \chi)$, let

$$f(q) = \sum_{n \geq 0} \mathbf{a}(n, f) q^n$$

be the q -expansion (at the infinity cusp). If κ is a Dirichlet character modulo M , define $f|[\kappa] \in \mathcal{M}_k(NM^2, \chi\kappa^2)$ the twist of f by κ to be the unique modular form with the q -expansion

$$f|[\kappa](q) = \sum_{n \geq 0, (n, M)=1} \mathbf{a}(n, f) \kappa(n) q^n.$$

2.2. Automorphic forms on $\mathrm{GL}_2(\mathbf{A})$. Let N be a positive integer. Define open-compact subgroups of $\mathrm{GL}_2(\widehat{\mathbf{Z}})$ by

$$U_0(N) = \left\{ g \in \mathrm{GL}_2(\widehat{\mathbf{Z}}) \mid g \equiv \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N\widehat{\mathbf{Z}}} \right\},$$

$$U_1(N) = \left\{ g \in U_0(N) \mid g \equiv \begin{pmatrix} * & * \\ 0 & 1 \end{pmatrix} \pmod{N\widehat{\mathbf{Z}}} \right\}.$$

Let $\omega : \mathbf{Q}^\times \backslash \mathbf{A}^\times \rightarrow \mathbf{C}^\times$ be a finite order Hecke character of level N . We extend ω to a character of $U_0(N)$ defined by $\omega\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) = \prod_{\ell \mid N} \omega_\ell(d_\ell)$ for $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in U_0(N)$, where $\omega_\ell : \mathbf{Q}_\ell^\times \rightarrow \mathbf{C}^\times$ is the ℓ -component of ω . Denote by $\mathcal{A}(\omega)$ the space of automorphic forms on $\mathrm{GL}_2(\mathbf{A})$ with central character ω . For any integer k , let $\mathcal{A}_k(N, \omega) \subset \mathcal{A}(\omega)$ be the space of automorphic forms on $\mathrm{GL}_2(\mathbf{A})$ of weight k , level N and character ω . Namely, $\mathcal{A}_k(N, \omega)$ consists of automorphic forms $\varphi : \mathrm{GL}_2(\mathbf{A}) \rightarrow \mathbf{C}$ such that

$$\varphi(\alpha g u_\infty u_f) = \varphi(g) e^{\sqrt{-1}k\theta} \omega(u_f)$$

$$(\alpha \in \mathrm{GL}_2(\mathbf{Q}), u_\infty = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}, u_f \in U_0(N)).$$

Let $\mathcal{A}_k^0(N, \omega)$ be the space of cusp forms in $\mathcal{A}_k(N, \omega)$.

Next we introduce important local Hecke operators on automorphic forms. At the archimedean place, let $V_\pm : \mathcal{A}_k(N, \omega) \rightarrow \mathcal{A}_{k \pm 2}(N, \omega)$ be the normalized weight raising/lowering operator in [JL70, page 165] given by

$$(2.1) \quad V_\pm = \frac{1}{(-8\pi)} \left(\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \otimes 1 \pm \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \otimes \sqrt{-1} \right) \in \mathrm{Lie}(\mathrm{GL}_2(\mathbf{R})) \otimes_{\mathbf{R}} \mathbf{C}.$$

The level-raising operator $V_\ell : \mathcal{A}_k(N, \omega) \rightarrow \mathcal{A}_k(N\ell, \omega)$ at a finite prime ℓ by

$$V_\ell \varphi(g) := \rho \left(\begin{pmatrix} \varpi_\ell^{-1} & 0 \\ 0 & 1 \end{pmatrix} \right) \varphi.$$

If $d = \prod_{\ell} \ell^{v_{\ell}(d)}$ is an positive integer, define $V_d : \mathcal{A}_k(N, \omega) \rightarrow \mathcal{A}_k(Nd, \chi)$ by

$$V_d = \prod_{\ell} V_{\ell}^{v_{\ell}(d)}.$$

Define the operator \mathbf{U}_{ℓ} on $\varphi \in \mathcal{A}_k(N, \omega)$ by

$$\mathbf{U}_{\ell}\varphi = \sum_{x \in \mathbf{Z}_{\ell}/\ell\mathbf{Z}_{\ell}} \rho\left(\begin{pmatrix} \varpi_{\ell} & x \\ 0 & 1 \end{pmatrix}\right)\varphi.$$

Note that $\mathbf{U}_{\ell}V_{\ell}\varphi = \ell\varphi$ and that if $\ell \mid N$, then $\mathbf{U}_{\ell} \in \text{End}_{\mathbf{C}} \mathcal{A}_k(N, \omega)$. For each prime $\ell \nmid N$, let $T_{\ell} \in \text{End}_{\mathbf{C}} \mathcal{A}_k(N, \omega)$ be the usual Hecke operator defined by

$$T_{\ell} = \mathbf{U}_{\ell} + \omega(\varpi_{\ell})V_{\ell}.$$

We introduce the twisting operator θ_{ℓ}^{κ} attached to a Dirichlet character κ of modulo ℓ^s for some $s > 0$. Let ℓ^n be the conductor of κ . If $n > 0$, define the Gauss sum $\mathfrak{g}(\kappa)$ by

$$\mathfrak{g}(\kappa) = \sum_{x \in (\mathbf{Z}/\ell^n\mathbf{Z})^{\times}} \kappa^{-1}(x) e^{\frac{-2\pi\sqrt{-1}x}{\ell^n}}.$$

For $\varphi \in \mathcal{A}_k(N, \omega)$, we define $\theta_{\ell}^{\kappa}\varphi : \text{GL}_2(\mathbf{A}) \rightarrow \mathbf{C}$ by

$$(2.2) \quad \theta_{\ell}^{\kappa}\varphi = \begin{cases} \varphi - \ell^{-1}V_{\ell}\mathbf{U}_{\ell}\varphi & \text{if } n = 0, \\ \mathfrak{g}(\kappa)^{-1} \sum_{x \in (\mathbf{Z}/\ell^n\mathbf{Z})^{\times}} \kappa^{-1}(x) \rho\left(\begin{pmatrix} 1 & x/\varpi_{\ell}^n \\ 0 & 1 \end{pmatrix}\right)\varphi & \text{if } n > 0. \end{cases}$$

2.3. We give a dictionary between modular forms and automorphic forms. For every Dirichlet character $\chi : (\mathbf{Z}/N\mathbf{Z})^{\times} \rightarrow \mathbf{C}^{\times}$, we let $\chi_{\mathbf{A}}$ be the *adelization* of χ , the unique finite order Hecke character $\chi_{\mathbf{A}} : \mathbf{Q}^{\times} \backslash \mathbf{A}^{\times} / \mathbf{R}_{+}(1 + N\widehat{\mathbf{Z}})^{\times} \rightarrow \mathbf{C}^{\times}$ of level N such that $\chi_{\mathbf{A}}(\varpi_{\ell}) = \chi(\ell)^{-1}$ for any prime $\ell \nmid N$. Each nearly holomorphic modular form $f \in \mathcal{N}_k(N, \chi)$ defines a unique automorphic form $\Phi(f) \in \mathcal{A}_k(N, \chi_{\mathbf{A}}^{-1})$ by the formula

$$(2.3) \quad \Phi(f)(\alpha g_{\infty} u) := (f|_k g_{\infty})(\sqrt{-1}) \cdot \chi_{\mathbf{A}}^{-1}(u)$$

for $\alpha \in \text{GL}_2(\mathbf{Q})$, $g_{\infty} \in \text{GL}_2^{+}(\mathbf{R})$ and $u \in U_0(N)$ (cf. [Cas73, §3]). Conversely, we can recover the form f from $\Phi(f)$ by

$$(2.4) \quad f(x + \sqrt{-1}y) = y^{-\frac{k}{2}} \Phi(f)\left(\begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix}\right).$$

The weight raising/lowering operators are the adelic avatar of the Maass-Shimura differential operators δ_k^m and ε on the space of automorphic forms. A direct computation shows that

$$(2.5) \quad \Phi(\delta_k^m f) = V_{+}^m \Phi(f) \text{ and } \Phi(\varepsilon f) = V_{-} \Phi(f).$$

In particular, f is holomorphic if and only if $V_{-} \Phi(f) = 0$. For a positive integer d , we have

$$(2.6) \quad \Phi(V_d f) = d^{1-\frac{k}{2}} V_d \Phi(f),$$

and for a finite prime ℓ

$$(2.7) \quad \Phi(T_{\ell} f) = \ell^{\frac{k}{2}-1} T_{\ell} \Phi(f); \quad \Phi(\mathbf{U}_{\ell} f) = \ell^{\frac{k}{2}-1} \mathbf{U}_{\ell} \Phi(f).$$

For $f \in \mathcal{M}_k(N, \chi)$ and κ a Dirichlet character modulo a ℓ -power, one verifies that

$$(2.8) \quad \Phi(f|[\kappa]) = \theta_{\ell}^{\kappa} \Phi(f) \otimes \kappa_{\mathbf{A}}^{-1}.$$

2.4. Preliminaries on irreducible representations of $\text{GL}_2(\mathbf{Q}_v)$.

2.4.1. *Measures.* We shall normalize the Haar measures on \mathbf{Q}_v and \mathbf{Q}_v^\times as follows. If $v = \infty$, dx or dy denotes the usual Lebesgue measure on \mathbf{R} and the measure $d^\times y$ on \mathbf{R}^\times is $|y|^{-1} dy$. If $v = \ell$ is a finite prime, denote by dx the Haar measure on \mathbf{Q}_ℓ with $\text{vol}(\mathbf{Z}_\ell, dx) = 1$ and by $d^\times y$ the Haar measure on \mathbf{Q}_ℓ^\times with $\text{vol}(\mathbf{Z}_\ell^\times, d^\times y) = 1$. Define the compact subgroup \mathbf{K}_v of $\text{GL}_2(\mathbf{Q}_v)$ by $\mathbf{K}_v = \text{O}(2, \mathbf{R})$ if $v = \infty$ and $\mathbf{K}_v = \text{GL}_2(\mathbf{Z}_v)$ if v is finite. Let dk_v be the Haar measure on \mathbf{K}_v so that $\text{vol}(\mathbf{K}_v, dk_v) = 1$. Let dg_v be the Haar measure on $\text{PGL}_2(\mathbf{Q}_v)$ given by $dg_v = |y_v|^{-1} dx_v d^\times y_v dk_v$ for $g_v = \begin{pmatrix} y_v & x_v \\ 0 & 1 \end{pmatrix} k_v$ with $y_v \in \mathbf{Q}_v^\times$, $x_v \in \mathbf{Q}_v$ and $k_v \in \mathbf{K}_v$.

2.4.2. *Representations.* Denote by $\chi \boxplus v$ the irreducible principal series representation of $\text{GL}_2(\mathbf{Q}_v)$ attached to two characters $\chi, v : \mathbf{Q}_v^\times \rightarrow \mathbf{C}^\times$ such that $\chi v^{-1} \neq |\cdot|^\pm$. If $v = \infty$ is the archimedean place and $k \geq 1$ is an integer, denote by $\mathcal{D}_0(k)$ the discrete series of lowest weight k if $k \geq 2$ or the limit of discrete series if $k = 1$ with central character sgn^k (the k -th power of the sign function). If v is finite, denote by St the Steinberg representation and by χSt the special representation $\text{St} \otimes \chi \circ \det$.

2.4.3. *L -functions and ε -factors.* For a character $\chi : \mathbf{Q}_v^\times \rightarrow \mathbf{C}^\times$, let $L(s, \chi)$ be the complex L -function and $\varepsilon(s, \chi) := \varepsilon(s, \chi, \psi_{\mathbf{Q}_v})$ be the ε -factor (cf. [Sch02, Section 1.1]). Define the γ -factor

$$(2.9) \quad \gamma(s, \chi) := \varepsilon(s, \chi) \cdot \frac{L(1-s, \chi^{-1})}{L(s, \chi)}.$$

If π is an irreducible admissible generic representation of $\text{GL}_2(\mathbf{Q}_v)$, denote by $L(s, \pi)$ the L -function and by $\varepsilon(s, \pi) := \varepsilon(s, \pi, \psi_{\mathbf{Q}_v})$ the ε -factor defined in [JL70, Theorem 2.18]. Let $\tilde{\pi}$ denote the contragredient representation of π . Denote by $L(s, \pi, \text{Ad})$ the adjoint L -function of π determined in [GJ78].

2.4.4. *Conductors and new vectors.* Let ℓ be a prime. Let (π, \mathcal{V}_π) be an irreducible admissible infinite dimensional representation of $\text{GL}_2(\mathbf{Q}_\ell)$, where \mathcal{V}_π a realization of π . For a non-negative integer n , let

$$\mathcal{U}_1(\ell^n) = \text{GL}_2(\mathbf{Z}_\ell) \cap \begin{pmatrix} \mathbf{Z}_\ell & \mathbf{Z}_\ell \\ \ell^n \mathbf{Z}_\ell & 1 + \ell^n \mathbf{Z}_\ell \end{pmatrix}.$$

Let $c(\pi)$ be the exponent of the conductor of π . By definition, $c(\pi)$ is the smallest integer such that $\mathcal{V}_\pi^{\mathcal{U}_1(\ell^{c(\pi)})}$ the space of $\mathcal{U}_1(\ell^{c(\pi)})$ -fixed vectors is non-zero. Define the subspace $\mathcal{V}_\pi^{\text{new}}$ by

$$\mathcal{V}_\pi^{\text{new}} = \left\{ \xi \in \mathcal{V}_\pi \mid \pi \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) \xi = \xi \text{ for all } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathcal{U}_1(\ell^{c(\pi)}) \right\}.$$

Proposition 2.1 (Multiplicity one for new vectors). *We have $\dim_{\mathbf{C}} \mathcal{V}_\pi^{\text{new}} = 1$.*

PROOF. This is [Cas73, Theorem 1]. □

We shall call $\mathcal{V}_\pi^{\text{new}}$ the new line of π .

2.4.5. *Whittaker models.* Every admissible irreducible infinite dimensional representation π of $\text{GL}_2(\mathbf{Q}_v)$ admits a realization of the Whittaker model $\mathcal{W}(\pi) = \mathcal{W}(\pi, \psi_{\mathbf{Q}_v})$ associated with the additive character $\psi_{\mathbf{Q}_v}$. Recall that $\mathcal{W}(\pi)$ is a subspace of smooth functions $W : \text{GL}_2(\mathbf{Q}_v) \rightarrow \mathbf{C}$ such that

- $W \left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} g \right) = \psi_{\mathbf{Q}_v}(x) W(g)$ for all $x \in \mathbf{Q}_v$,
- if $v = \infty$ is the archimedean place, there exists an integer M such that

$$W \left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \right) = O(|a|^M) \text{ as } |a| \rightarrow \infty.$$

The group $\text{GL}_2(\mathbf{Q}_v)$ (or the Hecke algebra of $\text{GL}_2(\mathbf{Q}_v)$) acts on $\mathcal{W}(\pi)$ via the right translation ρ . We introduce the (normalized) local Whittaker newform W_π in $\mathcal{W}(\pi)$ in the following cases. If $v = \infty$ and $\pi = \mathcal{D}_0(k)$, then the Whittaker local newform $W_\pi \in \mathcal{W}(\pi)$ is defined by

$$(2.10) \quad W_\pi(z) \begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} = \mathbb{I}_{\mathbf{R}_+}(y) \cdot y^{\frac{k}{2}} e^{-2\pi y} \cdot \text{sgn}(z)^k \psi_{\mathbf{R}}(x) e^{\sqrt{-1}k\theta}$$

$$(y, z \in \mathbf{R}^\times, x, \theta \in \mathbf{R}).$$

Here $\mathbb{I}_{\mathbf{R}_+}(a)$ denotes the characteristic function of the set of positive real numbers. If $v = \ell$ is a finite prime, then the local Whittaker newform W_π is the unique function in $\mathcal{W}(\pi)^{\text{new}}$ such that $W_\pi(1) = 1$.

2.5. Ordinary lines in irreducible representations of $\mathrm{GL}_2(\mathbf{Q}_p)$. Let p be a prime. Let (π, \mathcal{V}_π) be an irreducible admissible generic representation of $\mathrm{GL}_2(\mathbf{Q}_p)$ with central character $\omega : \mathbf{Q}_p^\times \rightarrow \mathbf{C}^\times$. Let $N(\mathbf{Z}_p) = \left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \mid x \in \mathbf{Z}_p \right\}$. Define the local \mathbf{U}_p -operator and the local level-raising operator V_p in $\mathrm{End}_{\mathbf{C}}(\mathcal{V}_\pi^{N(\mathbf{Z}_p)})$ by

$$(2.11) \quad \mathbf{U}_p \xi := \sum_{x \in \mathbf{Z}_p/p\mathbf{Z}_p} \pi \left(\begin{pmatrix} p & x \\ 0 & 1 \end{pmatrix} \right) \xi; \quad V_p \xi = \pi \left(\begin{pmatrix} p^{-1} & 0 \\ 0 & 1 \end{pmatrix} \right) \xi.$$

For a Dirichlet character κ of conductor p^n , we define the local twisting operator $\theta_p^\kappa \in \mathrm{End} \mathcal{V}_\pi$ by

$$(2.12) \quad \theta_p^\kappa \xi = \begin{cases} \xi - p^{-1} V_p \mathbf{U}_p \xi & \text{if } n = 0, \\ \mathfrak{g}(\kappa)^{-1} \sum_{x \in (\mathbf{Z}/p^n \mathbf{Z})^\times} \kappa^{-1}(x) \pi \left(\begin{pmatrix} 1 & x/p^n \\ 0 & 1 \end{pmatrix} \right) \xi & \text{if } n > 0. \end{cases}$$

For a character $\chi : \mathbf{Q}_p^\times \rightarrow \mathbf{C}^\times$, define the subspace $\mathcal{V}_\pi^{\mathrm{ord}}(\chi)$ by

$$\mathcal{V}_\pi^{\mathrm{ord}}(\chi) := \left\{ \xi \in \mathcal{V}_\pi^{N(\mathbf{Z}_p)} \mid \mathbf{U}_p \xi = \chi|\cdot|^{-\frac{1}{2}}(p) \cdot \xi, \pi \left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right) \xi = \chi(t) \xi, t \in \mathbf{Z}_p^\times \right\}.$$

Proposition 2.2 (Multiplicity one for ordinary vectors). *The space $\mathcal{V}_\pi^{\mathrm{ord}}(\chi)$ is non-zero if and only if π is either the principal series $\chi \boxplus \chi^{-1}\omega$ or the special representation $\chi|\cdot|^{-\frac{1}{2}}\mathrm{St}$. In this case,*

$$\dim_{\mathbf{C}} \mathcal{V}_\pi^{\mathrm{ord}}(\chi) = 1.$$

PROOF. Replacing π by $\pi \otimes \chi^{-1}|\cdot|^{\frac{1}{2}}$, we may assume $\chi = |\cdot|^{\frac{1}{2}}$. For each n , let

$$\mathcal{V}_\pi^{[n]}[\mathbf{U}_p - 1] = \{ \xi \in \mathcal{V}_\pi \mid \mathbf{U}_p \xi = \xi; \pi(u)\xi = \xi \text{ for all } u \in \mathcal{U}_1(p^n) \}.$$

Let $\mathcal{V}_\pi^{\mathrm{ord}} = \mathcal{V}_\pi^{\mathrm{ord}}(|\cdot|^{\frac{1}{2}})$. Let $c(\omega)$ be the exponent of the conductor of ω and $c^* := \max\{1, c(\omega)\}$. Then it is easy to see that

$$\mathcal{V}_\pi^{\mathrm{ord}} = \bigcup_{n \geq c^*} \mathcal{V}_\pi^{[n]}[\mathbf{U}_p - 1].$$

Suppose that $\pi = |\cdot|^{\frac{1}{2}} \boxplus \omega|\cdot|^{-\frac{1}{2}}$ or the Steinberg representation St . We claim that $\mathcal{V}_\pi^{[n]}[\mathbf{U}_p - 1]$ is non-zero for some n . If ω is ramified or π is Steinberg, then $c(\pi) \geq c^*$ and the new line $\mathcal{V}_\pi^{\mathrm{new}} = \mathcal{V}^{[c(\pi)]}[\mathbf{U}_p - 1]$ is not zero. If ω is unramified, then π is spherical, and it is well known that $\dim_{\mathbf{C}} \mathcal{V}_\pi^{[1]} = 2$ and the characteristic polynomial of \mathbf{U}_ℓ on $\mathcal{V}_\pi^{[1]}$ is given by $(X-1)(X-\omega(p)p)$, so $\mathcal{V}_\pi^{[1]}[\mathbf{U}_\ell - 1]$ is non-zero.

Now suppose that $\mathcal{V}_\pi^{\mathrm{ord}} \neq 0$. Then π must be a principal series or special representation since \mathbf{U}_p is a unipotent operator on $\mathcal{V}_\pi^{[n]}$ if π is supercuspidal. For any $u \in \mathcal{U}_1(p^m)$ with $m \geq 1$ and $\xi \in \mathcal{V}_\pi$, a straightforward calculation shows that

$$\pi(u)\mathbf{U}_p \xi = \sum_{x \in \mathbf{Z}_p/p\mathbf{Z}_p} \pi \left(\begin{pmatrix} p & x \\ 0 & 1 \end{pmatrix} u'_x z_x \right) \xi \text{ for some } u'_x \in \mathcal{U}_1(p^{m+1}), z_x \in 1 + p^m \mathbf{Z}_p.$$

It follows that if $\xi \in \mathcal{V}_\pi^{[m+1]}[\mathbf{U}_p - 1]$, then $\xi \in \mathcal{V}_\pi^{[m]}[\mathbf{U}_p - 1]$ whenever $m \geq c^*$. This implies that $\mathcal{V}_\pi^{\mathrm{ord}} = \mathcal{V}_\pi^{\mathcal{U}_1(p^{c^*})} \neq 0$, and hence $c^* \geq c(\pi) \geq c(\omega)$. If $c^* = c(\omega) > 0$, then $c(\omega) = c(\pi)$, and it follows that $\mathcal{V}_\pi^{\mathrm{ord}} = \mathcal{V}_\pi^{\mathrm{new}}$ is the new line in \mathcal{V}_π and $\pi = \mu \boxplus \mu^{-1}\omega$ with unramified character μ . Since any new vector in $\mu \boxplus \mu^{-1}\omega$ is an eigenvector of \mathbf{U}_p with the eigenvalue $\mu|\cdot|^{-\frac{1}{2}}$, we thus conclude that $\pi = |\cdot|^{\frac{1}{2}} \boxplus \omega|\cdot|^{-\frac{1}{2}}$. If $c(\omega) = 0$, then $c^* = 1$ and $\mathcal{V}_\pi^{\mathrm{ord}} = \mathcal{V}_\pi^{[1]}[\mathbf{U}_\ell - 1]$. It follows that π is a unramified principal series or the Steinberg representation St . If $\pi = \mathrm{St}$, then $\mathcal{V}_\pi^{\mathrm{ord}}$ is the new line. If π is a unramified principal series, then the two dimensional vector space $\mathcal{V}_\pi^{\mathcal{U}_1(p)}$ has a basis $\xi^0 \in \mathcal{V}_\pi^{\mathrm{new}} = \mathcal{V}_\pi^{\mathrm{GL}_2(\mathbf{Z}_p)}$ and $V_p \xi^0$. Since $\mathbf{U}_p V_p \xi^0 = p \xi^0$, \mathbf{U}_p is not a scalar, and thus $\dim_{\mathbf{C}} \mathcal{V}_\pi^{\mathrm{ord}} = \dim_{\mathbf{C}} \mathcal{V}_\pi^{[1]}[\mathbf{U}_p - 1] = 1$. \square

We shall call $\mathcal{V}_\pi^{\mathrm{ord}}(\chi)$ the ordinary line of π with respect to χ whenever it is non-zero.

Corollary 2.3. *If π is either the irreducible principal series $\chi \boxplus \chi^{-1}\omega$ or the special representation $\chi|\cdot|^{-\frac{1}{2}}\text{St}$, then the ordinary line $\mathcal{W}(\pi)^{\text{ord}}(\chi)$ in the Whittaker model is generated by the normalized ordinary Whittaker function W_{π}^{ord} characterized by*

$$W_{\pi}^{\text{ord}}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) = \chi|\cdot|^{\frac{1}{2}}(y)\mathbb{I}_{\mathbf{Z}_p}(y) \quad (y \in \mathbf{Q}_p^{\times}).$$

Here $\mathbb{I}_{\mathbf{Z}_p}$ is the characteristic function of \mathbf{Z}_p .

PROOF. The proof of Proposition 2.2 actually gives the recipe to construct the ordinary line. Indeed, let $W = W_{\pi \otimes \chi^{-1}}$ be the Whittaker local newform of $\pi \otimes \chi^{-1}$. Define $W^{\dagger} \in \mathcal{W}(\pi \otimes \chi^{-1})$ as follows: $W^{\dagger} = W$ if $\pi \otimes \chi^{-1}$ is not spherical and $W^{\dagger} = W - \chi^{-2}\omega|\cdot|^{\frac{1}{2}}(p)\rho\left(\begin{pmatrix} p^{-1} & 0 \\ 0 & 1 \end{pmatrix}\right)W$ if $\pi \otimes \chi^{-1}$ is spherical. An elementary calculation shows that $W^{\dagger} \otimes \chi$ belongs to $\mathcal{W}_{\pi}^{\text{ord}}(\chi)$. By using the explicit formulas of Whittaker newforms ([Sch02, Section 2.4]), we find that $W^{\dagger} \otimes \chi\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) = \chi|\cdot|^{\frac{1}{2}}(y)\mathbb{I}_{\mathbf{Z}_p}(y)$ as desired. \square

2.6. p -stabilized newforms. Let π be a cuspidal automorphic representation of $\text{GL}_2(\mathbf{A})$ and let $\mathcal{A}(\pi)$ be the π -isotypic part in the space of automorphic forms on $\text{GL}_2(\mathbf{A})$. For $\varphi \in \mathcal{A}(\pi)$, the Whittaker function of φ (with respect to the additive character $\psi_{\mathbf{Q}} : \mathbf{A}/\mathbf{Q} \rightarrow \mathbf{C}^{\times}$) is given by

$$W_{\varphi}(g) = \int_{\mathbf{A}/\mathbf{Q}} \varphi\left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}g\right)\psi_{\mathbf{Q}}(-x)dx \quad (g \in \text{GL}_2(\mathbf{A})),$$

where dx is the Haar measure with $\text{vol}(\mathbf{A}/\mathbf{Q}, dx) = 1$. We have the Fourier expansion:

$$\varphi(g) = \sum_{\alpha \in \mathbf{Q}^{\times}} W_{\varphi}\left(\begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix}g\right)$$

(cf. [Bum97, Theorem 3.5.5]). Let $f(q) = \sum_n a(n, f)q^n \in \mathcal{S}_k(N, \chi)$ be a normalized Hecke eigenform, we shall denote by $\pi_f = \otimes'_v \pi_{f,v}$ the cuspidal automorphic representation of $\text{GL}_2(\mathbf{A})$ generated by the automorphic form $\Phi(f)$ attached to f . Then π_f is irreducible and unitary with the central character χ^{-1} . If f is newform, then the conductor of π_f is N , $\Phi(f)$ is the normalized new vector in $\mathcal{A}_0(\pi_f)$ and the Mellin transform

$$Z(s, \Phi(f)) = \int_{\mathbf{A}^{\times}/\mathbf{Q}^{\times}} \Phi(f)\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right)|y|^{s-\frac{1}{2}}d^{\times}y = L(s, \pi_f)$$

is the automorphic L -function of π_f . Here $d^{\times}y$ is the product measure $\prod_v d^{\times}y_v$.

Definition 2.4 (p -stabilized newform). Let p be a prime and fix an isomorphism $\iota_p : \mathbf{C} \simeq \overline{\mathbf{Q}}_p$. We say that a normalized Hecke eigenform $f \in \mathcal{S}_k(Np, \chi)$ is a (ordinary) p -stabilized newform (with respect to ι_p) if f is a new outside p and the eigenvalue of \mathbf{U}_p , i.e. the p -th Fourier coefficient $\iota_p(a(p, f))$, is a p -adic unit. The prime-to- p part of the conductor of f is called *the tame conductor* of f .

Remark 2.5. Let f be a p -stabilized newform. By the multiplicity one for new and ordinary vectors, the Whittaker function of $\Phi(f)$ is a product of local Whittaker functions in $\mathcal{W}(\pi_{f,v})$. To be precise,

$$W_{\Phi(f)}(g) = W_{\pi_{f,p}}^{\text{ord}}(g_v) \prod_{v \neq p} W_{\pi_{f,v}}(g_v) \quad (g = (g_v) \in \text{GL}_2(\mathbf{A})).$$

Comparing the Fourier expansions of $\Phi(f)$ and f via (2.4), we find that

$$(2.13) \quad W_{\pi_{f,\ell}}\left(\begin{pmatrix} \ell & 0 \\ 0 & 1 \end{pmatrix}\right) = \mathbf{a}(\ell, f)\ell^{-\frac{k}{2}} \text{ if } \ell \neq p; \quad W_{\pi_{f,p}}^{\text{ord}}\left(\begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}\right) = \mathbf{a}(p, f)p^{-\frac{k}{2}}.$$

By Corollary 2.3, $W_{\pi_{f,p}}^{\text{ord}} \in \mathcal{W}(\pi_{f,p})^{\text{ord}}(\alpha_{f,p})$, where $\alpha_{f,p}$ is the unramified character with $\alpha_{f,p}(p) = \mathbf{a}(p, f)p^{\frac{1-k}{2}}$.

2.7. The bilinear form. Let $\mathcal{A}^0(\omega)$ be the space of cusp forms in $\mathcal{A}(\omega)$. Let \langle, \rangle denote the $\mathrm{GL}_2(\mathbf{A})$ -equivariant pairing between $\mathcal{A}^0(\omega)$ and $\mathcal{A}^0(\omega^{-1})$ defined by

$$\langle \varphi, \varphi' \rangle = \int_{\mathbf{A}^\times \mathrm{GL}_2(\mathbf{Q}) \backslash \mathrm{GL}_2(\mathbf{A})} \varphi(g) \varphi'(g) d^\tau g$$

for $\varphi \in \mathcal{A}^0(\omega), \varphi' \in \mathcal{A}^0(\omega^{-1})$, where $d^\tau g$ is the Tamagawa measure of $\mathrm{PGL}_2(\mathbf{A})$. The following lemma is well-known (cf. [Wal85, page 217]), and we omit the proof.

Lemma 2.6. *For cusp forms $\varphi \in \mathcal{A}_k^0(N, \omega)$ and $\varphi' \in \mathcal{A}_{-k}^0(N, \omega^{-1})$, we have*

$$\begin{aligned} \langle X\varphi, \varphi' \rangle &= -\langle \varphi, X\varphi' \rangle \text{ for } X \in \mathrm{Lie}(\mathrm{GL}_2(\mathbf{R})), \\ \langle \varphi, \mathbf{U}_\ell \varphi' \rangle &= \ell \langle V_\ell \varphi, \varphi' \rangle \text{ for } \ell \mid N, \\ \langle T_\ell \varphi, \varphi' \rangle &= \omega(\ell) \langle \varphi, T_\ell \varphi' \rangle \text{ for } \ell \nmid N. \end{aligned}$$

Let $\pi = \otimes'_v \pi_v$ be an irreducible unitary cuspidal automorphic representation on $\mathrm{GL}_2(\mathbf{A})$ with central character ω . Denote by $\tilde{\pi}$ the contragredient representation of π . By the multiplicity one theorem, the pairing \langle, \rangle gives rise to the equality $\mathcal{A}(\tilde{\pi}) = \mathcal{A}(\pi) \otimes \omega^{-1}$. For a place v of \mathbf{Q} , define the non-degenerate $\mathrm{GL}_2(\mathbf{Q}_v)$ -equivariant pairing \langle, \rangle between $\mathcal{W}(\pi_v)$ and $\mathcal{W}(\tilde{\pi}_v)$ by

$$(2.14) \quad \langle W, W' \rangle = \int_{\mathbf{Q}_v^\times} W\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) W'\left(\begin{pmatrix} -y & 0 \\ 0 & 1 \end{pmatrix}\right) d^\times y$$

for $W \in \mathcal{W}(\pi_v)$ and $W' \in \mathcal{W}(\tilde{\pi}_v)$. This integral converges absolutely as π_v is unitarizable.

Proposition 2.7. *Let $\varphi \in \mathcal{A}(\pi)$ and $\varphi' \in \mathcal{A}(\tilde{\pi})$. Suppose that $W_\varphi = \prod_v W_v$ and $W_{\varphi'} = \prod_v W'_v$ such that $W_v(1) = W'_v(1) = 1$ for all but finitely many v . Then we have*

$$\langle \varphi, \varphi' \rangle = \frac{2L(1, \pi, \mathrm{Ad})}{\zeta_{\mathbf{Q}}(2)} \prod_v \frac{\zeta_v(2)}{\zeta_v(1)L(1, \pi_v, \mathrm{Ad})} \langle W_v, W'_v \rangle.$$

PROOF. This is [Wal85, Proposition 6]. Note that $W_v = W_{\pi_v}$ and $W'_v = W_{\tilde{\pi}_v}$ are the normalized local Whittaker newforms for all but finitely many v , and if π_v is spherical, then

$$\langle W_{\pi_v}, W_{\tilde{\pi}_v} \rangle = \frac{\zeta_v(1)L(1, \pi_v, \mathrm{Ad})}{\zeta_v(2)},$$

so the right hand side of the equation in the proposition is indeed a finite product. \square

We give the formula of the local pairing of ordinary Whittaker functions.

Lemma 2.8. *Let p be a prime. Suppose that π_p is a principal series $\chi \boxplus v$ or a special representation $\chi|\cdot|^{-\frac{1}{2}} \mathrm{St}$. Let $W_{\pi_p}^{\mathrm{ord}} \in \mathcal{W}(\pi_p)^{\mathrm{ord}}(\chi)$ be the normalized ordinary Whittaker function in Corollary 2.3. If $n \geq \max\{1, c(\pi_p)\}$, then we have*

$$\langle \rho\left(\begin{pmatrix} 0 & p^{-n} \\ -p^n & 0 \end{pmatrix}\right) W_{\pi_p}^{\mathrm{ord}}, W_{\pi_p}^{\mathrm{ord}} \otimes \omega_p^{-1} \rangle = \chi(-1) \chi v^{-1} |\cdot| (p^n) \cdot \gamma(0, v\chi^{-1}) \zeta_p(1).$$

Here $v = \chi^{-1} \omega_p$ and $\gamma(s, -)$ is the γ -factor defined in (2.9).

PROOF. Let $W = W_{\pi_p}^{\mathrm{ord}}$ and $t_n = \begin{pmatrix} 0 & p^{-n} \\ -p^n & 0 \end{pmatrix}$. We first note that if $n \geq \max\{1, c_p(\pi)\}$, then $W\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} t_n\right) = 0$ if $y \notin \mathbf{Z}_p$. Then we have

$$\begin{aligned} \langle \rho(t_n) W, W \otimes \omega_p^{-1} \rangle &= \int_{\mathbf{Q}_p^\times} W\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} t_n\right) W\left(\begin{pmatrix} -y & 0 \\ 0 & 1 \end{pmatrix}\right) \omega_p^{-1}(-y) d^\times y \\ &= \int_{\mathbf{Q}_p^\times} W\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} t_n\right) \chi \omega_p^{-1}(-y) |\cdot|^{s-\frac{1}{2}}(y) d^\times y|_{s=1}. \end{aligned}$$

By the local functional equation for $\mathrm{GL}(2)$ (cf. [Bum97, Theorem 4.7.5]), the last integral equals

$$\begin{aligned} & \omega_p(-p^{-n})\varepsilon(1-s, \pi \otimes \chi^{-1}) \frac{L(s, \pi \otimes \chi\omega_p^{-1})}{L(1-s, \pi \otimes \chi^{-1})} \chi(-1) \int_{\mathbf{Q}_p^\times} W\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -p^{2n} & 0 \end{pmatrix}\right) \chi^{-1}|\cdot|^{1/2-s}(y) d^\times y|_{s=1} \\ & = \omega_p(p^{-n})\chi(-1)\chi(p^{2n}) \cdot |p^{2n}|^{s-\frac{1}{2}} \varepsilon(1-s, \pi \otimes \chi^{-1}) \frac{L(s, \pi \otimes \chi\omega_p^{-1})}{L(1-s, \pi \otimes \chi^{-1})} \zeta_p(1-s)|_{s=1} \end{aligned}$$

Using the formula

$$\varepsilon(1-s, \pi \otimes \chi^{-1}) \frac{L(s, \pi \otimes \chi\omega_p^{-1})}{L(1-s, \pi \otimes \chi^{-1})} = \begin{cases} \varepsilon(1-s, v\chi^{-1}) \frac{\zeta_p(s)L(s, \chi v^{-1})}{\zeta_p(1-s)L(1-s, v\chi^{-1})} & \text{if } \pi_p = \chi \boxplus v, \\ -|p|^{-s} \frac{\zeta_p(s+1)}{\zeta_p(1-s)} & \text{if } \pi_p = \chi|\cdot|^{-\frac{1}{2}}\mathrm{St}, \end{cases}$$

we see that

$$\langle \rho(t_n)W, W \otimes \omega_p^{-1} \rangle = \chi(-1)\omega_p(p^{-n})\chi|\cdot|^{\frac{1}{2}}(p^n) \begin{cases} \gamma(0, v\chi^{-1})\zeta_p(1) & \text{if } \pi_p = \chi \boxplus v, \\ -|p|^{-1}\zeta_p(2) & \text{if } \pi_p = \chi|\cdot|^{-\frac{1}{2}}\mathrm{St}. \end{cases}$$

Finally, we note that if $\pi = \chi|\cdot|^{-\frac{1}{2}}\mathrm{St}$, then $v = \chi|\cdot|^{-1}$ and $\gamma(0, v\chi^{-1})\zeta_p(1) = -|p|^{-1}\zeta_p(2)$. This finishes the proof. \square

2.8. Root numbers and Petersson norms. Let $f \in \mathcal{S}_k(N, \chi)$ be a normalized cuspidal newform of weight k and conductor N . Put $f_c(z) := f(-\bar{z})$. Then it is a classical result that

$$(2.15) \quad f|_k \begin{pmatrix} 0 & -1 \\ N & 0 \end{pmatrix} = w(f) \cdot f_c$$

for some $w(f) \in \mathbf{C}^\times$ with the modulus $|w(f)| = 1$ (cf. [Miy06, Theorem 4.6.15]). This complex number $w(f)$ is called the *root number* of f . By [Hid88c, page 38], we have

$$w(f) = \prod_{\ell < \infty} \varepsilon(1/2, \pi_{f, \ell}).$$

Recall that the Petersson norm of f is defined by

$$\|f\|_{\Gamma_0(N)}^2 = \int_{\Gamma_0(N) \backslash \mathfrak{H}} |f(x + \sqrt{-1}y)|^2 y^k \frac{dx dy}{y^2}.$$

For each integer M , define the matrix $\tau_M = (\tau_{M, v}) \in \mathrm{GL}_2(\mathbf{A})$ by

$$(2.16) \quad \begin{aligned} \tau_{M, \infty} &= \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \tau_{M, \ell} = 1 \text{ if } \ell \nmid M; \\ \tau_{M, \ell} &= \begin{pmatrix} 0 & 1 \\ -\ell v_\ell(M) & 0 \end{pmatrix} \in \mathrm{GL}_2(\mathbf{Q}_\ell) \text{ if } \ell \mid M. \end{aligned}$$

Let $\pi = \pi_f$ be the cuspidal automorphic representation generated by $\Phi(f)$ with central character $\omega(= \chi_{\mathbf{A}}^{-1})$. Define the local norm of the normalized Whittaker newform W_{π_v} by

$$(2.17) \quad B_{\pi_v} = \frac{\zeta_v(2)}{\zeta_v(1)L(1, \pi_v, \mathrm{Ad})} \langle \rho(\tau_{N, v})W_{\pi_v}, W_{\pi_v} \otimes \omega_v^{-1} \rangle.$$

It is straightforward to verify that

$$B_{\pi_\infty} = 2^{-1-k}, \quad B_{\pi_\ell} = 1 \text{ if } \ell \nmid N.$$

By Proposition 2.7 and (2.15), we have

$$(2.18) \quad \|f\|_{\Gamma_0(N)}^2 = \frac{[\mathrm{SL}_2(\mathbf{Z}) : \Gamma_0(N)]}{2^k \cdot w(f)} \cdot L(1, \pi, \mathrm{Ad}) \cdot \prod_{q|N} B_{\pi_q}.$$

3. THE UNBALANCED p -ADIC TRIPLE PRODUCT L -FUNCTIONS

3.1. Ordinary Λ -adic modular forms. Let $p > 2$ be a prime and let \mathcal{O} be the ring of integers of a finite extension of \mathbf{Q}_p . Let \mathbf{I} be a normal domain finite flat over $\Lambda = \mathcal{O}[[1 + p\mathbf{Z}_p]]$. A point $Q \in \text{Spec } \mathbf{I}(\overline{\mathbf{Q}}_p)$, a ring homomorphism $Q : \mathbf{I} \rightarrow \overline{\mathbf{Q}}_p$ is said to be locally algebraic if $Q|_{1+p\mathbf{Z}_p}$ is a locally algebraic character in the sense that $Q(z) = z^{k_Q} \epsilon_Q(z)$ with k_Q an integer and $\epsilon_Q(z) \in \mu_{p^\infty}$. We shall call k_Q the *weight* of Q and ϵ_Q the *finite part* of Q . Let $\mathfrak{X}_{\mathbf{I}}$ be the set of locally algebraic points $Q \in \text{Spec } \mathbf{I}(\overline{\mathbf{Q}}_p)$ of weight $k_Q \geq 1$. A point $Q \in \mathfrak{X}_{\mathbf{I}}$ is called *arithmetic* if the weight $k_Q \geq 2$ and let $\mathfrak{X}_{\mathbf{I}}^+$ be the set of arithmetic points. Let $\wp_Q = \text{Ker } Q$ be the prime ideal of \mathbf{I} corresponding to Q and $\mathcal{O}(Q)$ be the image of \mathbf{I} under Q .

Fix an isomorphism $\iota_p : \mathbf{C}_p \simeq \mathbf{C}$ once and for all. Denote by $\omega : (\mathbf{Z}/p\mathbf{Z})^\times \rightarrow \mu_{p-1}$ the p -adic Teichmüller character. Let N be a positive integer prime to p and let $\chi : (\mathbf{Z}/Np\mathbf{Z})^\times \rightarrow \mathcal{O}^\times$ be a Dirichlet character modulo Np . Denote by $\mathbf{S}(N, \chi, \mathbf{I})$ the space of \mathbf{I} -adic cusp forms of tame level N and (even) branch character χ , consisting of formal power series $\mathbf{f}(q) = \sum_{n \geq 1} \mathbf{a}(n, \mathbf{f})q^n \in \mathbf{I}[[q]]$ with the following property: there exists an integer $a_{\mathbf{f}}$ such that for arithmetic points $Q \in \mathfrak{X}_{\mathbf{I}}^+$ with $k_Q \geq a_{\mathbf{f}}$, the specialization $\mathbf{f}_Q(q)$ is the q -expansion of a cusp form $\mathbf{f}_Q \in \mathcal{S}_{k_Q}(Np^e, \chi\omega^{2-k_Q}\epsilon_Q)$. We call the character χ is the *branch character* of \mathbf{f} .

The space $\mathbf{S}(N, \chi, \mathbf{I})$ is equipped with the action of the usual Hecke operators T_ℓ for $\ell \nmid Np$ as in [Wil88, page 537] and the operators \mathbf{U}_ℓ for $\ell \mid pN$ given by $\mathbf{U}_\ell(\sum_n \mathbf{a}(n, \mathbf{f})q^n) = \sum_n \mathbf{a}(n\ell, \mathbf{f})q^n$. For a positive integer d prime to p , define $V_d : \mathbf{S}(N, \chi, \mathbf{I}) \rightarrow \mathbf{S}(Nd, \chi, \mathbf{I})$ by $V_d(\sum_n \mathbf{a}(n, \mathbf{f})q^n) = d \sum_n \mathbf{a}(n, \mathbf{f})q^{dn}$. Recall that Hida's ordinary projector e is defined by

$$e := \lim_{n \rightarrow \infty} \mathbf{U}_p^{n!}.$$

This ordinary projector e has a well-defined action on the space of classical modular forms preserving the cuspidal part as well as on the space $\mathbf{S}(N, \chi, \mathbf{I})$ of \mathbf{I} -adic cusp forms (*cf.* [Wil88, page 537 and Prop. 1.2.1]). The space $e\mathbf{S}(N, \chi, \mathbf{I})$ is called the space of ordinary \mathbf{I} -adic forms defined over \mathbf{I} . A key result in Hida's theory of ordinary \mathbf{I} -adic cusp forms is that if $\mathbf{f} \in e\mathbf{S}(N, \chi, \mathbf{I})$, then for *every* arithmetic points $Q \in \mathfrak{X}_{\mathbf{I}}$, we have $\mathbf{f}_Q \in e\mathcal{S}_{k_Q}(Np^e, \chi\omega^{2-k_Q}\epsilon_Q)$. We say $\mathbf{f} \in e\mathbf{S}(N, \chi, \mathbf{I})$ is a *primitive Hida family* if for every arithmetic points $Q \in \mathfrak{X}_{\mathbf{I}}$, \mathbf{f}_Q is a p -stabilized cuspidal newform of tame conductor N . Let $\mathfrak{X}_{\mathbf{I}}^{\text{cls}}$ be the set of classical points (for \mathbf{f}) given by

$$\mathfrak{X}_{\mathbf{I}}^{\text{cls}} := \{Q \in \mathfrak{X}_{\mathbf{I}}^{\text{cls}} \mid \mathbf{f}_Q \text{ is the } q\text{-expansion of a classical modular form}\}.$$

Note that $\mathfrak{X}_{\mathbf{I}}^{\text{cls}}$ contains the set of arithmetic points $\mathfrak{X}_{\mathbf{I}}^+$ but may be strictly larger than $\mathfrak{X}_{\mathbf{I}}^+$ as we allow the possibility of the points of weight one.

3.2. Galois representation attached to Hida families. Let $\langle \cdot \rangle : \mathbf{Z}_p^\times \rightarrow 1 + p\mathbf{Z}_p$ be character defined by $\langle x \rangle = x\omega^{-1}(x)$ and write $z \mapsto [z]_\Lambda$ for the inclusion of group-like elements $1 + p\mathbf{Z}_p \rightarrow \mathcal{O}[[1 + p\mathbf{Z}_p]]^\times = \Lambda^\times$. For $z \in \mathbf{Z}_p^\times$, denote by $\langle z \rangle_{\mathbf{I}} \in \mathbf{I}^\times$ the image of $[\langle z \rangle]_\Lambda$ in \mathbf{I} under the structure morphism $\Lambda \rightarrow \mathbf{I}$. By definition, $Q(\langle z \rangle_{\mathbf{I}}) = Q(\langle z \rangle)$ for $Q \in \mathfrak{X}_{\mathbf{I}}$. Let $\epsilon_{\text{cyc}} : G_{\mathbf{Q}} \rightarrow \mathbf{Z}_p^\times$ be the p -adic cyclotomic character and let $\langle \epsilon_{\text{cyc}} \rangle_{\mathbf{I}} : G_{\mathbf{Q}} \rightarrow \mathbf{I}^\times$ be the character $\langle \epsilon_{\text{cyc}} \rangle_{\mathbf{I}}(\sigma) = \langle \epsilon_{\text{cyc}}(\sigma) \rangle_{\mathbf{I}}$. For each Dirichlet character χ , we define $\chi_{\mathbf{I}} : G_{\mathbf{Q}} \rightarrow \mathbf{I}^\times$ by $\chi_{\mathbf{I}} := \sigma_\chi \langle \epsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{-2}$, where σ_χ is the Galois character which sends the geometric Frobenius element Frob_ℓ at ℓ to $\chi(\ell)^{-1}$.

If $\mathbf{f} \in e\mathbf{S}(N, \chi, \mathbf{I})$ is primitive Hida family of tame conductor N , we let $\rho_{\mathbf{f}} : G_{\mathbf{Q}} \rightarrow \text{GL}_2(\text{Frac } \mathbf{I})$ be the \mathbf{I} -adic Galois representation attached to \mathbf{f} characterized by

$$\text{Tr}(\rho_{\mathbf{f}}(\text{Frob}_\ell)) = \mathbf{a}(\ell, \mathbf{f}); \quad \det \rho_{\mathbf{f}}(\text{Frob}_\ell) = \chi\omega^2(\ell) \langle \ell \rangle_{\mathbf{I}} \ell^{-1} \quad (\ell \nmid N).$$

Note that $\det \rho_{\mathbf{f}} = \chi_{\mathbf{I}}^{-1} \cdot \epsilon_{\text{cyc}}^{-1}$. We have a complete knowledge of the description of the restriction of $\rho_{\mathbf{f}}$ to the local decomposition group $G_{\mathbf{Q}_\ell}$. For $\ell = p$, according to [Wil88, Theorem 2.2.1],

$$\rho_{\mathbf{f}}|_{G_{\mathbf{Q}_p}} \sim \begin{pmatrix} \alpha_p & * \\ 0 & \alpha_p^{-1} \chi_{\mathbf{I}}^{-1} \epsilon_{\text{cyc}}^{-1} \end{pmatrix}$$

where $\alpha_p : G_{\mathbf{Q}_p} \rightarrow \mathbf{I}^\times$ the unramified character with $\alpha_p(\text{Frob}_p) = \mathbf{a}(p, \mathbf{f})$.² For $\ell \neq p$, enlarging \mathbf{I} if necessary, we have the following list of $\rho_{\mathbf{f}}|_{G_{\mathbf{Q}_\ell}}$.

²Our representation $\rho_{\mathbf{f}}$ is the dual of $\rho_{\mathcal{F}}$ considered in [Wil88].

- (1) (Principal series) $\rho_{\mathbf{f}}|_{G_{\mathbf{Q}_\ell}}$ is reducible and isomorphic to

$$\alpha_\ell \xi \varepsilon_{\text{cyc}}^{1/2} \langle \varepsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{-1/2} \oplus \alpha_\ell^{-1} \xi' \varepsilon_{\text{cyc}}^{1/2} \langle \varepsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{-1/2}$$

with a unramified characters $\alpha_\ell : G_{\mathbf{Q}_\ell} \rightarrow \mathbf{I}^\times$ and a finite order characters $\xi, \xi' : G_{\mathbf{Q}_\ell} \rightarrow \overline{\mathbf{Q}}^\times$ with $\xi \xi' = \chi^{-1} \omega^{-2}$.

- (2) (Special) $\rho_{\mathbf{f}}|_{G_{\mathbf{Q}_\ell}}$ is indecomposable and

$$\rho_{\mathbf{f}}|_{G_{\mathbf{Q}_\ell}} \sim \begin{pmatrix} \xi \varepsilon_{\text{cyc}} \langle \varepsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{-1/2} & * \\ 0 & \xi \langle \varepsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{-1/2} \end{pmatrix}$$

with a finite order character $\xi : G_{\mathbf{Q}_\ell} \rightarrow \overline{\mathbf{Q}}^\times$ such that $\xi^2 = \chi^{-1} \omega^{-2}$.

- (3) (Supercuspidal) $\rho_{\mathbf{f}}|_{G_{\mathbf{Q}_\ell}}$ is irreducible and $\rho_{\mathbf{f}} \simeq \rho_0 \otimes \langle \varepsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{-1/2}$ with $\rho_0 : G_{\mathbf{Q}_\ell} \rightarrow \text{GL}_2(\overline{\mathbf{Q}})$ irreducible representation of finite image

(*cf.* [SU06, page 689]).

Remark 3.1 (Rigidity of automorphic types). We recall the *rigidity of automorphic types* for a primitive Hida family \mathbf{f} in [FO12, Lemma 2.14]. Let $\ell \neq p$ be a prime. If for some arithmetic point Q the associated cuspidal automorphic representation $\pi_{\mathbf{f}_{Q,\ell}}$ is principal series (resp. special, supercuspidal) of conductor ℓ^n , then for *any* arithmetic point Q' , $\pi_{\mathbf{f}_{Q',\ell}}$ is also principal series (resp. special, supercuspidal) of the same conductor ℓ^n . This is a consequence of the above description of $\rho_{\mathbf{f}}|_{G_{\mathbf{Q}_\ell}}$, the Langlands correspondence and the Ramanujan conjecture for elliptic modular forms (only needed in the case (Special)).

In addition, if $\pi_{\mathbf{f}_{Q,\ell}}$ is a discrete series at any arithmetic point $Q \in \mathfrak{X}_{\mathbf{I}}^+$, then the Weil-Deligne representaiton associated with the specialization of $\rho_{\mathbf{f}} \otimes \langle \varepsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{1/2} |_{G_{\mathbf{Q}_\ell}}$ at Q is independent of Q .

3.3. Hecke algebras and congruence numbers. If N is a positive integer and χ a Dirichlet character modulo N , we let $\mathbb{T}_k(N, \chi)$ be the \mathcal{O} -subalgebra in $\text{End}_{\mathbf{C}} e\mathcal{S}_k(N, \chi)$ generated over \mathcal{O} by the Hecke operators T_ℓ for $\ell \nmid Np$ and the operators \mathbf{U}_ℓ for $\ell \mid Np$. Suppose that N is *prime to* p . Let $\Delta = (\mathbf{Z}/Np\mathbf{Z})^\times$ and $\widehat{\Delta}$ be the group of Dirichlet characters modulo Np . Enlarging \mathcal{O} if necessary, we assume that every $\chi \in \widehat{\Delta}$ takes value in \mathcal{O}^\times . We are going to consider the Hecke algebra $\mathbf{T}(N, \mathbf{I})$ acting on the space of ordinary Λ -adic cusp forms of tame level $\Gamma_1(N)$ defined by

$$\mathbf{S}(N, \mathbf{I})^{\text{ord}} := \bigoplus_{\chi \in \widehat{\Delta}} e\mathbf{S}(N, \chi, \mathbf{I}).$$

In addition to the action of Hecke operators, denote by σ_d the usual diamond operator for $d \in \Delta$ acting on $\mathbf{S}(N, \mathbf{I})^{\text{ord}}$ by $\sigma_d(\mathbf{f})_{\chi \in \widehat{\Delta}} = (\chi(d)\mathbf{f})_{\chi \in \widehat{\Delta}}$. Then the ordinary \mathbf{I} -adic cuspidal Hecke algebra $\mathbf{T}(N, \mathbf{I})$ is defined to be the \mathbf{I} -subalgebra of $\text{End}_{\mathbf{I}} \mathbf{S}(N, \mathbf{I})^{\text{ord}}$ generated over \mathbf{I} by T_ℓ for $\ell \mid Np$, \mathbf{U}_ℓ for $\ell \mid Np$ and the diamond operators σ_d for $d \in \Delta$. Let $Q \in \mathfrak{X}_{\mathbf{I}}^+$ be an arithmetic point. Every $t \in \mathbf{T}(N, \mathbf{I})$ commutes with the specialization: $(t \cdot \mathbf{f})_Q = t \cdot \mathbf{f}_Q$. For $\chi \in \widehat{\Delta}_{Np}$, let $\wp_{Q,\chi}$ be the ideal of $\mathbf{T}(N, \mathbf{I})$ generated by \wp_Q and $\{\sigma_d - \chi(d)\}_{d \in \Delta}$. A classical result [Hid88b, Theorem 3.4] in Hida theory asserts that

$$\mathbf{T}(N, \mathbf{I})/\wp_{Q,\chi} \simeq \mathbb{T}_{k_Q}(Np^e, \chi \omega^{2-k_Q} \epsilon_Q) \otimes_{\mathcal{O}} \mathcal{O}(Q).$$

Let $\mathbf{f} \in e\mathbf{S}(N, \chi, \mathbf{I})$ be a primitive Hida family of tame level N and character χ and let $\lambda_{\mathbf{f}} : \mathbf{T}(N, \mathbf{I}) \rightarrow \mathbf{I}$ be the corresponding homomorphism defined by $\lambda_{\mathbf{f}}(T_\ell) = \mathbf{a}(\ell, \mathbf{f})$ for $\ell \nmid Np$, $\lambda_{\mathbf{f}}(\mathbf{U}_\ell) = \mathbf{a}(\ell, \mathbf{f})$ for $\ell \mid Np$ and $\lambda_{\mathbf{f}}(\sigma_d) = \chi(d)$ for $d \in \Delta$. Let $\mathbf{m}_{\mathbf{f}}$ be the maximal of $\mathbf{T}(N, \mathbf{I})$ containing $\text{Ker } \lambda_{\mathbf{f}}$ and let $\mathbf{T}_{\mathbf{m}_{\mathbf{f}}}$ be the localization of $\mathbf{T}(N, \mathbf{I})$ at $\mathbf{m}_{\mathbf{f}}$. It is the local ring of $\mathbf{T}(N, \mathbf{I})$ through which $\lambda_{\mathbf{f}}$ factors. Recall that the congruence ideal $C(\mathbf{f})$ of the morphism $\lambda_{\mathbf{f}} : \mathbf{T}_{\mathbf{m}_{\mathbf{f}}} \rightarrow \mathbf{I}$ is defined by

$$C(\mathbf{f}) := \lambda_{\mathbf{f}}(\text{Ann}_{\mathbf{T}_{\mathbf{m}_{\mathbf{f}}}}(\text{Ker } \lambda_{\mathbf{f}})) \subset \mathbf{I}.$$

It is well known that $\mathbf{T}_{\mathbf{m}_{\mathbf{f}}}$ is a reduced local finite flat Λ -alegba (*cf.* [DDT94, Prop. 4.7]), and there is an algebra direct sum decomposition

$$(3.1) \quad \lambda : \mathbf{T}_{\mathbf{m}_{\mathbf{f}}} \otimes_{\mathbf{I}} \text{Frac } \mathbf{I} \simeq \text{Frac } \mathbf{I} \oplus \mathcal{B}, \quad t \mapsto \lambda(t) = (\lambda_{\mathbf{f}}(t), \lambda_{\mathcal{B}}(t)),$$

where \mathcal{B} is some semi-simple $(\text{Frac } \mathbf{I})$ -algebra ([Hid88b, Corollaty 3.7]). Then by definition we have

$$C(\mathbf{f}) = \lambda_{\mathbf{f}}(\mathbf{T}_{\mathbf{m}_{\mathbf{f}}} \cap \lambda^{-1}(\text{Frac } \mathbf{I} \oplus \{0\})).$$

Now we impose the following

Hypothesis (CR). The residual Galois representation $\bar{\rho}_{\mathbf{f}}$ of $\rho_{\mathbf{f}}$ is absolutely irreducible and p -distinguished.

Under the above hypothesis, $\mathbf{T}_{\mathbf{m}_{\mathbf{f}}}$ is Gorenstein by [Wil95, Corollary 2, page 482], and with this Gorenstein property of $\mathbf{T}_{\mathbf{m}_{\mathbf{f}}}$, Hida in [Hid88a] proved that the congruence ideal $C(\mathbf{f})$ is generated by one element $\eta_{\mathbf{f}}$, called the congruence number for \mathbf{f} . Let $1_{\mathbf{f}}^*$ be the unique element in $\mathbf{T}_{\mathbf{m}_{\mathbf{f}}} \cap \lambda^{-1}(\text{Frac } \mathbf{I} \oplus \{0\})$ such that $\lambda_{\mathbf{f}}(1_{\mathbf{f}}^*) = \eta_{\mathbf{f}}$. Then $1_{\mathbf{f}} := \eta_{\mathbf{f}}^{-1} 1_{\mathbf{f}}^*$ is the idempotent in $\mathbf{T}_{\mathbf{m}_{\mathbf{f}}} \otimes_{\mathbf{I}} \text{Frac } \mathbf{I}$ corresponding to the direct summand $\text{Frac } \mathbf{I}$ of (3.1). Moreover, for each arithmetic point Q , it is also shown by Hida that the specialization $\eta_{\mathbf{f}}(Q) \in \mathcal{O}(Q)$ is the congruence number for \mathbf{f}_Q and

$$1_{\mathbf{f}} := \eta_{\mathbf{f}}^{-1} 1_{\mathbf{f}}^* \pmod{\wp_{\chi, Q}} \in \mathbb{T}_{k_Q}^{\text{ord}}(Np^r, \chi\omega^{2-k_Q}\epsilon_Q) \otimes_{\mathcal{O}} \text{Frac } \mathcal{O}(Q)$$

is the idempotent with $\lambda_{\mathbf{f}}(1_{\mathbf{f}}) = 1$.

For each Dirichlet character $\psi : (\mathbf{Z}/Np^r\mathbf{Z})^{\times} \rightarrow \mathbf{C}^{\times}$, we have a unique decomposition $\psi = \psi^{(p)}\psi_{(p)}$, where $\psi^{(p)}$ and $\psi_{(p)}$ are Dirichlet characters modulo N and p^r respectively. Let $\bar{\chi} = \chi^{-1}$ be the complex conjugation of χ . Denote by $\check{\mathbf{f}} \in e\mathbf{S}(N, \chi_{(p)}\bar{\chi}^{(p)}, \mathbf{I})$ the primitive Hida family corresponding to the twist $\mathbf{f}[[\bar{\chi}^{(p)}]](q) = \sum_{(n, N)=1} \bar{\chi}^{(p)}(n)\mathbf{a}(n, \mathbf{f})q^n$ (cf. [Dim14, Lemma 6.1]). To be precise, the Fourier coefficients of $\check{\mathbf{f}}$ are given by

$$\mathbf{a}(\ell, \check{\mathbf{f}}) = \begin{cases} \bar{\chi}^{(p)}(\ell)\mathbf{a}(\ell, \mathbf{f}) & \text{if } \ell \nmid N, \\ \mathbf{a}(\ell, \mathbf{f})^{-1}\chi_{(p)}\omega^{2-k_Q}\ell^{-1}\langle \ell \rangle_{\mathbf{I}} & \text{if } \ell \mid N. \end{cases}$$

by [Miy06, Theorem 4.6.16]. For every arithmetic point $Q \in \mathfrak{X}^+$, $\check{\mathbf{f}}_Q$ is the p -stabilized newform attached to $\mathbf{f}_Q[[\bar{\chi}^{(p)}]]$. Moreover, the Atkin-Lehner involution η'_p introduced in [Miy06, (4.6.21), page 168] induces an isomorphism $\eta'_p : \mathcal{S}_k(Np^r, \chi\omega^{2-k_Q}\epsilon_Q) \simeq \mathcal{S}_k(Np^r, \bar{\chi}^{(p)}\chi_{(p)}\omega^{2-k_Q}\epsilon_Q)$ such that $T_{\ell}\eta'_p = \bar{\chi}^{(p)}(\ell)\eta'_p T_{\ell}$ for $\ell \nmid N$ ([Miy06, (4.6.23)]). We thus obtain a Λ -algebra isomorphism $[\bar{\chi}^{(p)}] : \mathbf{T}_{\mathbf{m}_{\mathbf{f}}} \simeq \mathbf{T}_{\mathbf{m}_{\check{\mathbf{f}}}}$ such that $[\bar{\chi}^{(p)}](T_{\ell}) = T_{\ell} \cdot \bar{\chi}^{(p)}(\ell)$ for $\ell \nmid N$ and $\lambda_{\check{\mathbf{f}}} \circ [\bar{\chi}^{(p)}] = \lambda_{\mathbf{f}}$. It follows that

$$(3.2) \quad 1_{\check{\mathbf{f}}}^* = [\bar{\chi}^{(p)}](1_{\mathbf{f}}^*) \text{ and } \eta_{\check{\mathbf{f}}} = \eta_{\mathbf{f}}.$$

3.4. The adjustment of levels for a triple of modular forms. For any positive integer M , let $\text{supp}(M)$ denote the support of M , i.e. the set of prime factors of M . If f is a p -stabilized newform of tame conductor N_1 , let $c_{\ell}(f) := c(\pi_{f, \ell})$ be the exponent of the ℓ -component of N_1 for each prime $\ell \neq p$ and set

$$\begin{aligned} \Sigma_f^1 &= \{\ell : \text{prime} \mid \pi_{f, \ell} \text{ is a principal series}\}; \\ \Sigma_f^0 &= \{\ell : \text{prime} \mid \pi_{f, \ell} \text{ is a discrete series}\}. \end{aligned}$$

To a triple (f, g, h) of p -stabilized newforms of tame conductors (N_1, N_2, N_3) , we are going to associate a set of auxiliary integers, which we call *the adjustment of levels* for (f, g, h) . This adjustment of levels is crucial for the construction of our test Λ -adic modular forms (Definition 3.3 and Definition 4.8) in order to obtain the optimal value of the local zeta integrals in Ichino's formula, and they are defined according to the choice of good test vectors in the space of product of local representations of $\pi_{f, \ell} \times \pi_{g, \ell} \times \pi_{h, \ell}$ (cf. §6.1). Inevitably, the definition is ad-hoc and may seem to be artificial at the first sight. The readers may skip the details and come back until §6.1. To begin with, let $N_{fgh} = \gcd(N_1, N_2, N_3)$ and $N = \text{lcm}(N_1, N_2, N_3)$. Put

$$c_{\ell}^{\min} := v_{\ell}(N_{fgh})$$

and let $\Sigma_{fgh} = \Sigma_f^0 \cap \Sigma_g^0 \cap \Sigma_h^0$. We introduce several disjoint subsets of $\text{supp}(N)$:

$$\begin{aligned} \Sigma_{fgh}^{(I)} &= \{\ell \in \Sigma_f^1 \cup \Sigma_g^1 \cup \Sigma_{fgh} \mid c_{\ell}(h) < \min\{c_{\ell}(f), c_{\ell}(g)\}\}, \\ \Sigma_f^{(IIa)} &= \{\ell \in \Sigma_g^0 \cap \Sigma_h^0 \mid L(s, \pi_{g, \ell} \otimes \pi_{h, \ell}) \neq 1, c_{\ell}(f) = 0\}, \\ \Sigma_f^{(IIb)} &= \{\ell \in \Sigma_g^0 \cap \Sigma_h^0 \mid L(s, \pi_{g, \ell} \otimes \pi_{h, \ell}) = 1, \ell \in \Sigma_f^1, c_{\ell}(f) < \min\{c_{\ell}(g), c_{\ell}(h)\}\}, \\ \Sigma_f^{\max} &= \{\ell : \text{prime factor of } N_1 \mid c_{\ell}(g) = c_{\ell}(h) = c_{\ell}^{\min} < c_{\ell}(f)\}. \end{aligned}$$

Define $\Sigma_{fh}^{(I)}$, $\Sigma_{gh}^{(I)}$, $\Sigma_g^{(IIa)}$, $\Sigma_g^{(IIb)}$, Σ_g^{\max}, \dots , in the same manner. We set

$$\begin{aligned} \mathbf{d}_f^{(I)} &= \prod_{\ell \in \Sigma_{fg}^{(I)}} \ell^{c_\ell(fg) - c_\ell(f)} \cdot \prod_{\ell \in \Sigma_{fh}^{(I)}} \ell^{c_\ell(fh) - c_\ell(f)}, \\ \mathbf{d}_f^{(II)} &= \prod_{\ell \in \Sigma_f^{(IIa)}} \ell^{\lceil \frac{c_\ell(gh)}{2} \rceil} \cdot \prod_{\ell \in \Sigma_f^{(IIb)}} \ell^{c_\ell(gh) - c_\ell(f)}, \\ \mathbf{d}_f^{\max} &= \prod_{\ell \in \Sigma_f^{\max}} \ell^{c_\ell(f) - c_\ell^{\min}}. \end{aligned}$$

Likewise we define $\mathbf{d}_g^{(I)}$, $\mathbf{d}_g^{(II)}$, \mathbf{d}_g^{\max} , $\mathbf{d}_h^{(I)}$, $\mathbf{d}_h^{(II)}$ and \mathbf{d}_h^{\max} . Finally, put

$$(3.3) \quad \mathbf{d}_f = \mathbf{d}_f^{(I)} \mathbf{d}_f^{(II)}, \quad \mathbf{d}_g = \mathbf{d}_g^{(I)} \mathbf{d}_g^{\max} \mathbf{d}_g^{(II)}, \quad \mathbf{d}_h = \mathbf{d}_h^{(I)} \mathbf{d}_h^{\max} \mathbf{d}_h^{(II)}.$$

By definition, we have

$$(3.4) \quad \mathbf{d}_f \mid N/N_1, \quad \mathbf{d}_g \mid N/N_2, \quad \mathbf{d}_h \mid N/N_3.$$

3.5. Definitions of good test Λ -adic modular forms. Let $\mathcal{O} = \mathcal{O}_F$ for some finite extension F of \mathbf{Q}_p . Fixing a topological generator γ_0 of $1 + p\mathbf{Z}_p$, we let $\Lambda = \mathcal{O}[[1 + p\mathbf{Z}_p]] = \mathcal{O}[[T]]$ with $T = \gamma_0 - 1$. For $i = 1, 2, 3$, let \mathbf{I}_i be a normal domain finite flat over Λ and let $\psi_i : (\mathbf{Z}/pN_i\mathbf{Z})^\times \rightarrow \mathcal{O}^\times$ be Dirichlet characters with $\psi_i(-1) = 1$. Throughout this paper, we fix a triplet of primitive Hida families

$$\mathbf{F} := (\mathbf{f}, \mathbf{g}, \mathbf{h}) \in e\mathbf{S}(N_1, \psi_1, \mathbf{I}_1) \times e\mathbf{S}(N_2, \psi_2, \mathbf{I}_2) \times e\mathbf{S}(N_3, \psi_3, \mathbf{I}_3)$$

of tame conductors $\underline{N} = (N_1, N_2, N_3)$ and branch characters $\underline{\psi} = (\psi_1, \psi_2, \psi_3)$. We shall impose the following running hypotheses

$$(ev) \quad \psi_1 \psi_2 \psi_3 = \omega^{2a} \text{ for some } a \in \mathbf{Z};$$

$$(sf) \quad \gcd(N_1, N_2, N_3) \text{ is square-free.}$$

Lemma 3.2. *Let $(Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathbf{I}_1}^{\text{cls}} \times \mathfrak{X}_{\mathbf{I}_2}^{\text{cls}} \times \mathfrak{X}_{\mathbf{I}_3}^{\text{cls}}$ and $(f, g, h) = \mathbf{F}_{\underline{Q}} = (\mathbf{f}_{Q_1}, \mathbf{g}_{Q_2}, \mathbf{h}_{Q_3})$ be the specialization of \mathbf{F} at \underline{Q} . The adjustment of levels $\mathbf{d}_f^\bullet, \mathbf{d}_g^\bullet$ and \mathbf{d}_h^\bullet for $\bullet \in \{(I), (II), \max\}$ are independent of the choice of any arithmetic point \underline{Q} .*

PROOF. The lemma is clear from the rigidity of automorphic types, the description of the restriction of $\rho_{\mathbf{f}}|_{G_{Q_\ell}}$ given in §3.2 and the Langlands correspondence for $\text{GL}(2)$. \square

Definition 3.3 (Test Λ -adic forms). Let $N = \text{lcm}(N_1, N_2, N_3)$. Put

$$\Sigma_{?,0}^{(IIb)} = \left\{ \ell \in \Sigma_{?}^{(IIb)} \mid c_\ell(?) = 0 \right\} \text{ for } ? \in \{f, g, h\}.$$

For each $\ell \in \Sigma_{f,0}^{(IIb)}$ (resp. $\Sigma_{g,0}^{(IIb)}$, $\Sigma_{h,0}^{(IIb)}$), we fix once and for all a root $\beta_\ell(\mathbf{f}) \in \mathbf{I}_1^\times$ (resp. $\beta_\ell(\mathbf{g}) \in \mathbf{I}_2^\times$, $\beta_\ell(\mathbf{h}) \in \mathbf{I}_3^\times$) of the Hecke polynomial $H_{f,\ell}(X) := X^2 - \mathbf{a}(\ell, \mathbf{f})X + \psi_1 \omega^2(\ell) \ell^{-1} \langle \ell \rangle_{\mathbf{I}_1}$ (resp. $H_{g,\ell}(X)$, $H_{h,\ell}(X)$). With the above notation in the previous subsection, we define the pair $(\mathbf{g}^*, \mathbf{h}^*)$ in $e\mathbf{S}(N, \psi_2, \mathbf{I}_2) \times e\mathbf{S}(N, \psi_3, \mathbf{I}_3)$ of the ordinary Λ -adic cusp forms by

$$\begin{aligned} \mathbf{g}^*(q) &= \sum_{I \subset \Sigma_{g,0}^{(IIb)}} (-1)^{|I|} \beta_I(\mathbf{g})^{-1} V_{\mathbf{d}_g/n_I} \mathbf{g}, \\ \mathbf{h}^*(q) &= \sum_{I \subset \Sigma_{h,0}^{(IIb)}} (-1)^{|I|} \beta_I(\mathbf{h})^{-1} V_{\mathbf{d}_h/n_I} \mathbf{h}, \end{aligned}$$

where $n_I = \prod_{\ell \in I} \ell$, $\beta_I(?) = \prod_{\ell \in I} \beta_\ell(?)$ for $? = \mathbf{f}, \mathbf{g}, \mathbf{h}$.

3.6. **The construction of the p -adic L -function in the unblanced case.** We let

$$\mathcal{R} = \mathbf{I}_1 \widehat{\otimes}_{\mathcal{O}} \mathbf{I}_2 \widehat{\otimes}_{\mathcal{O}} \mathbf{I}_3$$

be a finite extension over the three variable Iwasawa algebra

$$\begin{aligned} \mathcal{R}_0 &:= \Lambda \widehat{\otimes}_{\mathcal{O}} \Lambda \widehat{\otimes}_{\mathcal{O}} \Lambda = \mathcal{O}[[T_1, T_2, T_3]], \\ (T_1 = T \otimes 1 \otimes 1, T_2 = 1 \otimes T \otimes 1, T_3 = 1 \otimes 1 \otimes T). \end{aligned}$$

Define the multiplicative map $\Theta : \mathbf{Z}_{(p)}^\times \rightarrow \mathcal{R}^\times$ by

$$\Theta(n) := \psi_{1,(p)} \omega^{-a-1}(n) \langle n \rangle_{\mathbf{I}_1}^{1/2} \langle n \rangle_{\mathbf{I}_2}^{-1/2} \langle n \rangle_{\mathbf{I}_3}^{-1/2}.$$

Define the \mathcal{R} -adic twisting operator $[[\Theta]] : \mathcal{R}[[q]] \rightarrow \mathcal{R}[[q]]$ by

$$\left(\sum_{n \geq 0} \mathbf{a}(n) q^n \right) [[\Theta]] = \sum_{n \geq 0, p \nmid n} \Theta(n) \cdot \mathbf{a}(n) q^n.$$

Here $\psi_{1,(p)}$ is the restriction of the branch character ψ_1 of \mathbf{f} to $(\mathbf{Z}/p\mathbf{Z})^\times$. Define the power series \mathbf{H} by

$$\mathbf{H} := \mathbf{g}^* \cdot \mathbf{h}^* [[\Theta]] \in \mathcal{R}[[q]].$$

Lemma 3.4. *The power series \mathbf{H} belongs to $\mathbf{S}(N, \psi_{1,(p)} \overline{\psi_1}^{(p)}, \mathbf{I}_1) \widehat{\otimes}_{\mathbf{I}_1} \mathcal{R}$.*

PROOF. The following proof is taken from Hida's blue book [Hid93]. Put

$$\mathfrak{X}_{\mathcal{R}}^0 = \{ \underline{Q} = (Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathbf{I}_1}^+ \times \mathfrak{X}_{\mathbf{I}_2}^+ \times \mathfrak{X}_{\mathbf{I}_3}^+ \mid k_{Q_1} = k_{Q_2} + k_{Q_3}, k_{Q_1} \geq k_{Q_2} + 2 \}.$$

For $\underline{Q} \in \mathfrak{X}_{\mathcal{R}}^0$, put

$$\mathbb{k}_0 = \psi_{1,(p)} \omega^{-a-1} \epsilon_{Q_1}^{1/2} \epsilon_{Q_2}^{-1/2} \epsilon_{Q_3}^{-1/2}.$$

Here $\epsilon_\tau^{1/2}$ is the unique square root of ϵ_τ taking value in $1 + p\mathbf{Z}_p$. We verify that $(\mathbf{h}^* [[\Theta]])_{\underline{Q}} = \mathbf{h}_{Q_3} [[\mathbb{k}_0]] \in \mathcal{S}_{k_{Q_3}}(N, \psi_{1,(p)}^2 \psi_1^{-1} \psi_2^{-1} \epsilon_{Q_1} \epsilon_{Q_2}^{-1})$, and hence we find that for every $\underline{Q} \in \mathfrak{X}_{\mathcal{R}}^0$,

$$(3.5) \quad \mathbf{H}_{\underline{Q}} = \mathbf{g}_{Q_2}^* \cdot \mathbf{h}_{Q_3}^* [[\mathbb{k}_0]] \in \mathcal{S}_{k_{Q_1}}(N, \psi_{1,(p)} \overline{\psi_1}^{(p)} \omega^{2-k_{Q_1}} \epsilon_{Q_1}).$$

We have $\mathcal{R}_0 = \mathcal{O}[[T_1, T_2, Z]]$ with $Z = (1 + T_1)^{-1}(1 + T_2)(1 + T_3) - 1$. Let $L_0 = \text{Frac } \mathcal{R}_0$ and $L = \text{Frac } \mathcal{R}$ be a finite extension of L_0 . Let $\alpha_1, \dots, \alpha_n$ be a basis of \mathcal{R} over \mathcal{R}_0 and write $\mathbf{H} = \sum_{j=1}^n \mathbf{H}^{(j)} \alpha_j$ with $\mathbf{H}^{(j)} \in \mathcal{R}_0[[q]]$. On the other hand, if we let $\{\alpha_j^*\}_{j=1, \dots, n}$ be the dual basis of $\{\alpha_j\}_{j=1, \dots, n}$ with respect to the trace map $\text{Tr} : L \rightarrow L_0$. Then $\mathbf{H}^{(j)} = \text{Tr}(\mathbf{H} \alpha_j^*)$. Let $\mathbf{u} = 1 + p$. By (3.5), we can write $\mathbf{H}^{(j)} = \mathbf{H}^{(j)}(T_1, T_2, Z) \in \mathcal{O}[[T_1, T_2, Z]][[q]]$ so that

$$(3.6) \quad \mathbf{H}^{(j)}(\mathbf{u}^{k_1} \zeta_1 - 1, \mathbf{u}^{k_2} \zeta_2 - 1, \zeta_3 - 1) = \text{Tr}(\mathbf{H}_{\underline{Q}} \alpha_i(\underline{Q})) \in \mathcal{S}_{k_1}(N, \psi_{1,(p)} \overline{\psi_1}^{(p)} \omega^{2-k_1})$$

for all but finite many positive integers k_1, k_2 with $k_1 \geq k_2 + 2$ and $\zeta_i \in \mu_{p^\infty}$ ($i = 1, 2, 3$), where $\underline{Q} = (Q_1, Q_2, Q_3)$ are some arithmetic points of weights $(k_1, k_2, k_1 - k_2)$ and finite parts $(\epsilon_{Q_1}, \epsilon_{Q_2}, \epsilon_{Q_3})$, $\epsilon_{Q_i}(z)$ is the finite order character with $\epsilon_{Q_i}(u) = \zeta_i$.

To prove the lemma, it suffices to show that

$$(3.7) \quad \mathbf{H}^{(j)}(T_1, T_2, Z) \in \mathbf{S} \widehat{\otimes}_{\mathcal{O}} \mathcal{O}[[T_2, Z]], \quad \mathbf{S} := \mathbf{S}(N, \psi_{1,(p)} \overline{\psi_1}^{(p)}, \mathcal{O}[[T_1]]),$$

which in turn, by [Hid93, Lemma 1 in page 328], is equivalent to showing that $\mathbf{H}^{(j)}(T_1, T_2, \zeta) \in \mathbf{S} \widehat{\otimes}_{\mathcal{O}} \mathcal{O}[[\zeta]][[T_2]]$ for every $\zeta \in \mu_{p^\infty}$. Now we repeat the arguments in [Hid93, page 226-227]. Let a be a positive integer such that \mathbf{g}_Q is a classical modular form for all $Q \in \mathfrak{X}_{\mathbf{I}}$ with $k_Q = a$. We define the power series inductively

$$\begin{aligned} H_0(T_1, T_2) &= \mathbf{H}^{(j)}(T_1, T_2, \zeta - 1), \quad Y_m = T_2 - (\mathbf{u}^{m+a-1} - 1) \in p\mathcal{O}[[T_2]], \\ H_m(T_1, T_2) &= \frac{H_{m-1}(T_1, T_2) - H_{m-1}(T_1, \mathbf{u}^{m+a-1} - 1)}{Y_m} \in \mathcal{O}[[T_1, T_2]][[q]] \end{aligned}$$

for $m = 1, 2, \dots$. Then (3.6) implies that $H_0(T_1, \mathbf{u}^a - 1) \in \mathbf{S} \otimes_{\mathcal{O}} \mathcal{O}[\zeta]$ and by induction, we find easily that $H_m(T_1, \mathbf{u}^{m+a} - 1) \in \mathbf{S} \otimes_{\mathcal{O}} \mathcal{O}[\zeta]$ for all $m = 0, 1, \dots$. On other hand, by construction we have

$$\mathbf{H}^{(j)}(T_1, T_2, \zeta - 1) = \sum_{m=0}^{\infty} H_m(T_1, \mathbf{u}^{m+2} - 1) \prod_{i=1}^m Y_i.$$

It is clear that the right hand side is a convergent power series and belongs to $\mathbf{S} \widehat{\otimes}_{\mathcal{O}} \mathcal{O}[\zeta][[T_2]]$. \square

The above lemma allows us to apply the ordinary projector e to \mathbf{H} and obtain $e\mathbf{H}$ an ordinary Λ -adic modular form with coefficients in \mathcal{R} . We define an auxiliary \mathcal{R} -adic form $\mathbf{H}^{\text{aux}} \in e\mathbf{S}(N, \psi_{1,(p)} \overline{\psi_1}^{(p)}, \mathbf{I}_1) \widehat{\otimes}_{\mathbf{I}_1} \mathcal{R}$ by

$$(3.8) \quad \mathbf{H}^{\text{aux}} := \sum_{I \subset \Sigma_{g,h}^{(b)}} (-1)^I \frac{\psi_{1,(p)}(n_I/\mathbf{d}_f) \langle n_I/\mathbf{d}_f \rangle_{\mathbf{I}_1} \mathbf{d}_f}{\beta_I(\mathbf{f}) n_I} \cdot \mathbf{U}_{\mathbf{d}_f/n_I}(e\mathbf{H}).$$

Now we are ready to define the p -adic L -function following the construction in [Hid85, (4.6)]. Denote by $\text{Tr}_{N/N_1} : e\mathbf{S}(N, \psi_{1,(p)} \overline{\psi_1}^{(p)}, \mathbf{I}_1) \rightarrow e\mathbf{S}(N_1, \psi_{1,(p)} \overline{\psi_1}^{(p)}, \mathbf{I}_1)$ the usual trace map (cf. [Hid88c, page 14]).

Definition 3.5. The unbalanced p -adic triple product L -function $\mathcal{L}_{\mathbf{F}}^{\mathbf{f}}$ is defined by

$$\mathcal{L}_{\mathbf{F}}^{\mathbf{f}} := \mathbf{a}(1, 1_{\mathbf{f}}^* \text{Tr}_{N/N_1}(\mathbf{H}^{\text{aux}})) \in \mathcal{R}.$$

3.7. Global trilinear period integrals. Define the weight space for the triple $(\mathbf{f}, \mathbf{g}, \mathbf{h})$ in the \mathbf{f} -dominated unbalanced range by

$$\mathfrak{X}_{\mathcal{R}}^{\mathbf{f}} := \{ \underline{Q} = (Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathbf{I}_1}^+ \times \mathfrak{X}_{\mathbf{I}_2}^{\text{cls}} \times \mathfrak{X}_{\mathbf{I}_3}^{\text{cls}} \mid k_{Q_1} \geq k_{Q_2} + k_{Q_3}, k_{Q_1} \equiv k_{Q_2} + k_{Q_3} \pmod{2} \}.$$

In this subsection, we relate the value of $\mathcal{L}_{\mathbf{F}}^{\mathbf{f}}(\underline{Q})$ at a point $\underline{Q} = (Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathcal{R}}^{\mathbf{f}}$ to a global trilinear period integral of a test triple of modular forms. To this end, it is necessary to work in the framework of automorphic forms. Let $(k_1, k_2, k_3) = (k_{Q_1}, k_{Q_2}, k_{Q_3})$ and let r be an integer greater than $\max\{1, c_p(\epsilon_{Q_1}), c_p(\epsilon_{Q_2}), c_p(\epsilon_{Q_3})\}$. Recall that the specialization

$$(f, g, h) := \mathbf{F}_{\underline{Q}} = (\mathbf{f}_{Q_1}, \mathbf{g}_{Q_2}, \mathbf{h}_{Q_3}) \in \mathcal{S}_{k_1}(N_1 p^r, \chi_f) \times \mathcal{S}_{k_2}(N_2 p^r, \chi_g) \times \mathcal{S}_{k_3}(N_3 p^r, \chi_h)$$

are p -stabilized cuspidal newforms with characters modulo Np^r

$$\chi_f = \psi_1 \epsilon_{Q_1} \omega^{2-k_1}, \chi_g = \psi_2 \epsilon_{Q_2} \omega^{2-k_2} \text{ and } \chi_h = \psi_3 \epsilon_{Q_3} \omega^{2-k_3}.$$

Let $\varphi_f = \Phi(f)$, $\varphi_g = \Phi(g)$ and $\varphi_h = \Phi(h)$ be the associated automorphic cusp forms as in (2.3). Then

$$(\varphi_f, \varphi_g, \varphi_h) \in \mathcal{A}_{k_1}^0(N_1 p^r, \omega_f) \times \mathcal{A}_{k_2}^0(N_2 p^r, \omega_g) \times \mathcal{A}_{k_3}^0(N_3 p^r, \omega_h),$$

and the central characters $\omega_f, \omega_g, \omega_h$ are the adelizations

$$\omega_f = (\chi_f^{-1})_{\mathbf{A}}, \omega_g = (\chi_g^{-1})_{\mathbf{A}}, \omega_h = (\chi_h^{-1})_{\mathbf{A}}.$$

Denote by $(\beta_{\ell}(f), \beta_{\ell}(g), \beta_{\ell}(h))$ the specialization $(\beta_{\ell}(\mathbf{f})(Q_1), \beta_{\ell}(\mathbf{g})(Q_2), \beta_{\ell}(\mathbf{h})(Q_3))$. For each finite prime ℓ , define the polynomial $\mathcal{Q}_{f,\ell} \in \mathcal{O}[X]$ by

$$\mathcal{Q}_{f,\ell}(X) = X^{v_{\ell}(\mathbf{d}_f)} \begin{cases} 1 & \text{if } \ell \notin \Sigma_{gh}^{(b)}, \\ (1 - \beta_{\ell}(f)^{-1} \ell^{\frac{k_1}{2}-1} X^{-1}) & \text{if } \ell \in \Sigma_{gh}^{(b)}. \end{cases}$$

We define $\mathcal{Q}_{g,\ell}(X)$ and $\mathcal{Q}_{h,\ell}(X)$ likewise. Set

$$(3.9) \quad \varphi_f^* = \prod_{\ell} \mathcal{Q}_{f,\ell}(V_{\ell}) \varphi_f, \quad \varphi_g^* = \prod_{\ell} \mathcal{Q}_{g,\ell}(V_{\ell}) \varphi_g \text{ and } \varphi_h^* = \prod_{\ell} \mathcal{Q}_{h,\ell}(V_{\ell}) \varphi_h.$$

By (2.6), we see that

$$\varphi_g^* = \mathbf{d}_g^{\frac{k_{Q_2}}{2}-1} \Phi(\mathbf{g}_{Q_2}^*) \text{ and } \varphi_h^* = \mathbf{d}_h^{\frac{k_{Q_3}}{2}-1} \Phi(\mathbf{h}_{Q_3}^*).$$

Decompose $\omega_f = \omega_{f,(p)}\omega_f^{(p)}$, where $\omega_{f,(p)}$ and $\omega_f^{(p)}$ are finite order Hecke characters of p -power conductor and prime-to- p conductor respectively. By definition, $\omega_{f,(p)}$ is the adelization of $\chi_{f,(p)}^{-1}$, the restriction of χ_f^{-1} to $(\mathbf{Z}/p^r\mathbf{Z})^\times$. Let \check{f} be the primitive Hida family corresponding to the twist $f|[\overline{\psi_1}^{(p)}]$ and put

$$\check{\varphi}_f = \Phi(\check{f}) \in \mathcal{A}_{k_1}^0(N_1 p^r, \omega_f^{-1} \omega_{f,(p)}^2).$$

We introduce the modified p -Euler factor $\mathcal{E}_p(f, \text{Ad})$ for the adjoint motive attached to the p -stabilized newform f . Let $\alpha_{f,p} : \mathbf{Q}_p^\times \rightarrow \mathbf{C}^\times$ be the unramified character as in Remark 2.5. Let $\beta_{f,p} := \alpha_{f,p}^{-1} \omega_{f,p}$. Hence the local component $\pi_{f,p}$ is either the principal series $\alpha_{f,p} \boxplus \beta_{f,p}$ or the special representation $\alpha_{f,p} |\cdot|^{-\frac{1}{2}} \text{St}$. Define the modified p -Euler factor $\mathcal{E}_p(f, \text{Ad})$ by

$$\begin{aligned} \mathcal{E}_p(f, \text{Ad}) &= \varepsilon(1, \beta_{f,p} \alpha_{f,p}^{-1}) L(0, \beta_{f,p} \alpha_{f,p}^{-1}) L(1, \beta_{f,p} \alpha_{f,p}^{-1}) \\ &= \mathbf{a}(p, f)^{-c_p(\pi_f)} \cdot p^{c_p(\pi_f)(\frac{k_1}{2}-1)} \varepsilon(1/2, \pi_{f,p}) \\ &\quad \times \begin{cases} (1 - \alpha_{f,p}^{-2} \omega_{f,p}(p))(1 - \alpha_{f,p}^{-2}(p)p^{-1}) & \text{if } c_p(\pi_f) = 0, \\ 1 & \text{if } c_p(\pi_f) > 0. \end{cases} \end{aligned} \quad (3.10)$$

Define \mathcal{J}_∞ and $t_n \in \text{GL}_2(\mathbf{A})$ by

$$\mathcal{J}_\infty = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \in \text{GL}_2(\mathbf{R}), \quad t_n = \begin{pmatrix} 0 & p^{-n} \\ -p^n & 0 \end{pmatrix} \in \text{GL}_2(\mathbf{Q}_p) \hookrightarrow \text{GL}_2(\mathbf{A}). \quad (3.11)$$

Lemma 3.6. *Let notation be as above. We have*

$$\langle \rho(\mathcal{J}_\infty t_n) \varphi_f, \check{\varphi}_f \otimes \omega_{f,(p)}^{-1} \rangle = \frac{\zeta_{\mathbf{Q}}(2)^{-1}}{[\text{SL}_2(\mathbf{Z}) : \Gamma_0(N_1)]} \cdot \|f^\circ\|_{\Gamma_0(N_f^\circ)}^2 \cdot \mathcal{E}_p(f, \text{Ad}) \cdot \frac{\omega_{f,p}^{-1} \alpha_{f,p}^2 |\cdot| (-p^n) \zeta_p(2)}{\zeta_p(1)}.$$

PROOF. Let $\pi = \otimes_v \pi_v$ be the irreducible automorphic cuspidal representation generated by $\varphi_f = \Phi(f)$ and let $\omega = \omega_f$ be the central character of π . Let $\varphi' = \rho(\mathcal{J}_\infty t_n) \varphi_f \in \mathcal{A}_{-k_{Q_1}}^0(N p^r, \omega)$ and $\varphi'' = \check{\varphi}_f \otimes \omega_{f,(p)}^{-1} \in \mathcal{A}_{k_{Q_1}}^0(N p^r, \omega^{-1})$. Then $\varphi' \in \mathcal{A}(\pi)$ and $\varphi'' \in \mathcal{A}(\pi^\vee)$. Since φ_f and $\check{\varphi}_f$ are automorphic forms attached to p -stabilized cuspidal newforms f and \check{f} , and $\omega_{f,(p)}$ is unramified outside p , according to Remark 2.5, the Whittaker functions $W_{\varphi'}$ and $W_{\varphi''}$ have the factorizations

$$W_{\varphi'} = \rho(t_n) W_{\pi_p}^{\text{ord}} \cdot \rho(\mathcal{J}_\infty) W_{\pi_\infty} \prod_{v \neq p, \infty} W_{\pi_v}, \quad W_{\varphi''} = W_{\pi_p}^{\text{ord}} \otimes \omega_p^{-1} \cdot W_{\pi_\infty}^\vee \prod_{v \neq p, \infty} W_{\pi_v}^\vee,$$

where $W_{\pi_p}^{\text{ord}} \in \mathcal{W}_{\pi_p}^{\text{ord}}(\alpha_p)$ is the ordinary Whittaker functions attached to the character α_p . On the other hand, let $\varphi^\circ = \Phi(f^\circ)$ be the normalized newform in $\mathcal{A}(\pi)$ and let $\overline{\varphi^\circ} \in \mathcal{A}(\pi^\vee)$ be the complex conjugation of φ° . Then $\rho(\mathcal{J}_\infty) \overline{\varphi^\circ}$ is the normalized newform in $\mathcal{A}(\pi^\vee)$.

Let $\alpha = \alpha_{f,p}$, $\beta = \beta_{f,p}$. Combining Proposition 2.7, Lemma 2.8 and the formula

$$\varepsilon(1/2, \pi_p) = \begin{cases} \varepsilon(1/2, \beta) & \text{if } \pi_p = \alpha \boxplus \beta \\ -\alpha |\cdot|_p^{-\frac{1}{2}}(p) & \text{if } \pi_p = \alpha |\cdot|_p^{-\frac{1}{2}} \text{St}, \end{cases}$$

we find that

$$\begin{aligned} \frac{\langle \varphi', \varphi'' \rangle}{\langle \varphi^\circ, \overline{\varphi^\circ} \rangle} &= \frac{\langle \rho(t_n) W_{\pi_p}^{\text{ord}}, W_{\pi_p}^{\text{ord}} \otimes \omega_p^{-1} \rangle}{\langle W_{\pi_p}, W_{\pi_p}^\vee \rangle} \\ &= \omega_p(-1) \alpha \beta^{-1} |\cdot|_p(p^n) \cdot \varepsilon(1/2, \pi_p) \begin{cases} (1 - \beta \alpha^{-1} |\cdot|_p(p))(1 - \beta \alpha^{-1}(p))(1 + p^{-1})^{-1} & \text{if } c(\pi_p) = 0, \\ \alpha^{-1} |\cdot|_p^{\frac{1}{2}}(p^{c(\pi_p)}) & \text{if } c(\pi_p) > 0. \end{cases} \end{aligned}$$

From above equation together with the following equation ([III10, page 1403])

$$\begin{aligned} \langle \varphi^\circ, \overline{\varphi^\circ} \rangle &= \frac{\zeta_{\mathbf{Q}}(2)^{-1}}{[\text{SL}_2(\mathbf{Z}) : \Gamma_0(N_1 p^s)]} \|f^\circ\|_{\Gamma_0(N_{f^\circ})}^2 \\ &= \|f^\circ\|_{\Gamma_0(N_{f^\circ})}^2 \frac{\zeta_{\mathbf{Q}}(2)^{-1}}{[\text{SL}_2(\mathbf{Z}) : \Gamma_0(N_1)]} \begin{cases} 1 & \text{if } c(\pi_p) = 0 \\ p^s (1 + p^{-1})^{-1} & \text{if } c(\pi_p) > 0, \end{cases} \end{aligned}$$

we can directly deduce the lemma. \square

We may regard $F := \mathbf{F}_{\underline{Q}} = (f, g, h)$ as the modular form on \mathfrak{H}^3 of weight (k_1, k_2, k_3) given by $F(z_1, z_2, z_3) = f(z_1)g(z_2)h(z_3)$. Let ω_F be the central character of $F|_{\mathfrak{H}}$ given by

$$\omega_F = \omega_f \omega_g \omega_h.$$

Let \mathbb{k} be the Dirichlet character modulo p^r defined by

$$(3.12) \quad \mathbb{k} = \psi_{1,(p)} \omega^{-a-1+\frac{k_2+k_3-k_1}{2}} \epsilon_{Q_1}^{1/2} \epsilon_{Q_2}^{-1/2} \epsilon_{Q_3}^{-1/2}.$$

By definition, $\mathbb{k}^2 = \chi_{f,(p)}^2 \chi_f^{-1} \chi_g^{-1} \chi_h^{-1}$. Define the character $\omega_F^{1/2}$ by

$$(3.13) \quad \omega_F^{1/2} = \omega_{f,(p)} \mathbb{k}_{\mathbf{A}} = \omega^{-a+\frac{k_1+k_2+k_3}{2}-3} \epsilon_{Q_1}^{-1/2} \epsilon_{Q_2}^{-1/2} \epsilon_{Q_3}^{-1/2}.$$

Then $\omega_F^{1/2}$ is a finite order Hecke character unramified outside p , and

$$(\omega_F^{1/2})^2 = \omega_f \omega_g \omega_h = \omega_F$$

as the notation suggests. Let $E = \mathbf{Q} \oplus \mathbf{Q} \oplus \mathbf{Q}$ be the split cubic étale algebra over \mathbf{Q} . Let

$$m = \frac{k_1 - k_2 - k_3}{2}.$$

Define the automorphic cusp form ϕ_F^* on $\mathrm{GL}_2(\mathbf{A}_E)$ by

$$(3.14) \quad \begin{aligned} \phi_F^* &:= (\rho(\mathcal{J}_{\infty}) \varphi_f^* \otimes \omega_F^{-1/2}) \boxtimes \varphi_g^* \boxtimes V_+^m \theta_p^{\mathbb{k}} \varphi_h^*, \\ \phi_F^*(x_1, x_2, x_3) &= \varphi_f^*(x_1 \mathcal{J}_{\infty}) \cdot \varphi_g^*(x_2) \cdot V_+^m \theta_p^{\mathbb{k}} \varphi_h^*(x_3) \cdot \omega_F^{-1/2}(\det x_1). \end{aligned}$$

Here $\theta_p^{\mathbb{k}}$ is the twisting operator as in (2.2). Put

$$\mathbf{t}_n = (t_n, 1, 1) \in \mathrm{GL}_2(E_p).$$

We shall relate the valuation of our p -adic L -function $\mathcal{L}_p(\underline{Q})$ at \underline{Q} to the global trilinear period $I(\rho(\mathbf{t}_n)\phi_F^*)$ defined by

$$I(\rho(\mathbf{t}_n)\phi_F^*) := \int_{\mathbf{A} \times \mathrm{GL}_2(\mathbf{Q}) \setminus \mathrm{GL}_2(\mathbf{A})} \phi_F^*(xt_n, x, x) d^r x.$$

Put

$$(3.15) \quad \mathbf{d}_F^{\mathbb{k}/2} := \mathbf{d}_f^{\frac{k_1-2}{2}} \mathbf{d}_g^{\frac{k_2-2}{2}} \mathbf{d}_h^{\frac{k_3-2}{2}}.$$

Proposition 3.7. *We have*

$$\mathcal{L}_F^{\mathbf{f}}(\underline{Q}) = \frac{\zeta_{\mathbf{Q}}(2)[\mathrm{SL}_2(\mathbf{Z}) : \Gamma_0(N)]}{\eta_f^{-1} \|f^\circ\|_{\Gamma_0(N, f^\circ)}^2 \mathcal{E}_p(f, \mathrm{Ad})} \cdot I(\rho(\mathbf{t}_n)\phi_F^*) \cdot \frac{\zeta_p(1)}{\omega_{f,p}^{-1} \alpha_{f,p}^2 | \cdot |_p (-p^n) \zeta_p(2)} \cdot \frac{1}{\mathbf{d}_F^{\mathbb{k}/2}}.$$

PROOF. First of all, since $\check{\mathbf{f}}_{Q_1}$ is a p -stabilized newform, by the multiplicity one for new and ordinary vectors, we have

$$1_{\check{\mathbf{f}}_{Q_1}}^* \mathrm{Tr}_{N/N_1}(\mathbf{H}_{\underline{Q}}^{\mathrm{aux}}) = \mathcal{L}_p^{\mathbf{f}}(\underline{Q}) \cdot \check{\mathbf{f}}_{Q_1}.$$

It follows that

$$(3.16) \quad \langle \rho(\mathcal{J}_{\infty} t_n) \varphi_f \otimes \omega_{f,(p)}^{-1}, \check{\varphi}_f \rangle \cdot \mathcal{L}_p^{\mathbf{f}}(\underline{Q}) = \langle \rho(\mathcal{J}_{\infty} t_n) \varphi_f \otimes \omega_{f,(p)}^{-1}, 1_{\check{\mathbf{f}}_{Q_1}}^* \mathrm{Tr}_{N/N_1} \Phi(\mathbf{H}_{\underline{Q}}^{\mathrm{aux}}) \rangle.$$

Since $\Theta(n)(\underline{Q}) = \mathbb{k}(n)n^m$ for $n \in \mathbf{Z}_{(p)}^\times$, from [Hid93, equation (2), page 330], we deduce that

$$(3.17) \quad e\mathbf{H}_{\underline{Q}} = e(\mathbf{g}_{Q_2}^* \mathbf{d}^m(\mathbf{h}_{Q_3}^* | [\mathbb{k}])) = e\mathrm{Hol}(\mathbf{g}_{Q_2}^* \delta_{k_{Q_3}}^m(\mathbf{h}_{Q_3}^* | [\mathbb{k}])),$$

where $\mathbf{d} = q \frac{d}{dq}$ is Serre's p -adic differential operator, Hol is the holomorphic projection as in [Hid93, (8a), page 314] and $\delta_{k_{Q_3}}^m$ is the Maass-Shimura differential operator. Put $H = \mathbf{g}_{Q_2}^* \cdot \delta_{k_{Q_3}}^m \mathbf{h}_{Q_3}^* | [\mathbb{k}]$. Using (2.5), (2.6) and (2.8), we see that

$$\varphi_H := \Phi(H) = \mathbf{d}_g^{1-\frac{k_2}{2}} \mathbf{d}_h^{1-\frac{k_3}{2}} \cdot \varphi_g^* \cdot V_+^m \theta_p^{\mathbb{k}} \varphi_h^* \otimes \mathbb{k}_{\mathbf{A}}^{-1}.$$

Then H is a nearly holomorphic cusp form of weight k_{Q_1} and $\varphi_H \in \mathcal{A}_{k_1}^0(Np^r, \omega_f^{-1} \omega_{f,(p)}^2)$ has a decomposition

$$\varphi_H = \mathrm{Hol}(\varphi_H) + V_+ \varphi'_1 + V_+^2 \varphi'_2 + \cdots + V_+^n \varphi'_n,$$

where $\text{Hol}(\varphi_H)$ and $\{\varphi'_j\}_{j=1,\dots,n}$ are holomorphic automorphic forms. It follows that $\text{Hol}(\varphi_H) = \Phi(\text{Hol}(H))$.

Let $1_f^* \in \mathbb{T}^{\text{ord}}(N_1 p^s, \chi_f)$ and $\eta_f \in \mathcal{O}(Q_1)$ denote the specializations of 1_f^* and η_f at Q_1 respectively. As a consequence of strong multiplicity one theorem for modular forms, the idempotent $1_f = \eta_f^{-1} 1_f^* \in \mathbb{T}^{\text{ord}}(N_1 p^s, \chi_f) \otimes_{\mathcal{O}} \text{Frac } \mathcal{O}(Q_1)$ is generated by the Hecke operators T_ℓ for $\ell \nmid Np$, so by Lemma 2.6 and (3.2), the right hand side of (3.16) equals

$$(3.18) \quad \langle 1_f^* \text{Tr}_{N/N_1} \rho(\mathcal{J}_\infty t_n) \varphi_f \otimes \omega_{f,(p)}^{-1}, \Phi(\mathbf{H}_{\underline{Q}}^{\text{aux}}) \rangle = \eta_f [K_0(N_1) : K_0(N)] \cdot \langle \rho(\mathcal{J}_\infty t_n) \varphi_f \otimes \omega_{f,(p)}^{-1}, \Phi(\mathbf{H}_{\underline{Q}}^{\text{aux}}) \rangle.$$

Note that for any prime $\ell \neq p$, $\omega_{f,(p)}(\varpi_\ell) = \chi_{f,(p)}(\ell)$ is the specialization of $\psi_{1,(p)}(\ell) \langle \ell \rangle_{\mathbf{I}_1}$ at Q_1 . From the definition (3.8), (3.17) and Lemma 2.6, we find that the pairing in the right hand side of (3.18) equals

$$\begin{aligned} & \mathbf{d}_f^{-\frac{k_1}{2}} \sum_{I \subset \Sigma_{gh}^{(b)}} (-1)^I \frac{n_I^{\frac{k_1}{2}}}{\beta_I(f)} \chi_{f,(p)}(n_I / \mathbf{d}_f) \cdot \langle \rho(\mathcal{J}_\infty t_n) \varphi_f \otimes \omega_{f,(p)}^{-1}, \mathbf{U}_{\mathbf{d}_f / n_I} e\text{Hol}(\varphi_H) \rangle \\ &= \mathbf{d}_f^{1-\frac{k_1}{2}} \langle \rho(\mathcal{J}_\infty t_n) \varphi_f^* \otimes \omega_{f,(p)}^{-1}, e\text{Hol}(\varphi_H) \rangle. \end{aligned}$$

On the other hand, it is straightforward to verify by Lemma 2.6 that

$$\begin{aligned} \langle \rho(t_n) \mathbf{U}_p \varphi, \varphi' \rangle &= \langle \varphi, \mathbf{U}_p \varphi' \rangle, \\ \langle \rho(\mathcal{J}_\infty) \varphi, V_+ \varphi' \rangle &= - \langle \rho(\mathcal{J}_\infty) V_- \varphi, \varphi' \rangle \end{aligned}$$

(cf. [Hid85, (5.4)]), and together with (3.13), it follows that

$$\begin{aligned} \langle \rho(\mathcal{J}_\infty t_n) \varphi_f^* \otimes \omega_{f,(p)}^{-1}, e\text{Hol}(\varphi_H) \rangle &= \langle \rho(\mathcal{J}_\infty t_n) \varphi_f^* \otimes \omega_F^{-1/2}, \varphi_g^* \cdot V_+^m \theta_p^k \varphi_h^* \rangle \\ &= I(\rho(\mathbf{t}_n) \phi_F^*). \end{aligned}$$

Combining the above equation with (3.16) and (3.18), we find that

$$\langle \rho(\mathcal{J}_\infty t_n) \varphi_f, \check{\varphi}_f \otimes \omega_{f,(p)}^{-1} \rangle \cdot \mathcal{L}_p^f(\underline{Q}) = \eta_f [\Gamma_0(N_1) : \Gamma_0(N)] \mathbf{d}_F^{-\kappa/2} \cdot I(\rho(\mathbf{t}_n) \phi_F^*).$$

From Lemma 3.6 the formula of the pairing in the left hand side, we obtain the lemma. \square

3.8. Ichino's period integral formula for triple products.

3.8.1. *The setting.* In this subsection, we apply Ichino's formula to express $I(\rho(\mathbf{t}_n) \phi_F^*)$ as a product of the central value of the triple product L -function attached to F and normalized local trilinear integrals. We retain the notation in the previous subsection. Let

$$\pi_1 = \pi_f \otimes \omega_F^{-1/2}, \quad \pi_2 = \pi_g \text{ and } \pi_3 = \pi_h$$

with central characters $\omega_1 = \omega_g^{-1} \omega_h^{-1}$, $\omega_2 = \omega_g$ and $\omega_3 = \omega_h$ respectively. Let

$$\Pi_{\underline{Q}} = \pi_1 \boxtimes \pi_2 \boxtimes \pi_3$$

be an irreducible unitary cuspidal automorphic representation of $\text{GL}_2(\mathbf{A}_E)$ and let $\mathcal{A}(\Pi_{\underline{Q}}) = \mathcal{A}(\pi_1) \otimes \mathcal{A}(\pi_2) \otimes \mathcal{A}(\pi_3)$ be the unique automorphic realization of $\Pi_{\underline{Q}}$. For brevity, we simply write Π for $\Pi_{\underline{Q}}$. For each place v , let $\mathcal{V}_{\Pi_v} = \mathcal{V}_{\pi_{1,v}} \otimes \mathcal{V}_{\pi_{2,v}} \otimes \mathcal{V}_{\pi_{3,v}}$ denote a realization of Π_v , where $\mathcal{V}_{\pi_{i,v}}$ is a realization of $\pi_{i,v}$ for $i = 1, 2, 3$. Then we have the factorizations

$$\Pi \simeq \bigotimes_v \Pi_v, \quad \mathcal{A}(\Pi) \simeq \bigotimes_v \mathcal{V}_{\Pi_v}.$$

We let $\phi_F = \varphi_1 \boxtimes \varphi_2 \boxtimes \varphi_3 \in \mathcal{A}(\Pi)$, where

$$\varphi_1 = \varphi_f \otimes \omega_F^{-1/2}, \quad \varphi_2 = \varphi_g \text{ and } \varphi_3 = \varphi_h.$$

Then we have a factorization $\phi_F = \bigotimes_v \phi_v$ via the above isomorphism. Since φ_f, φ_g and φ_h are p -stabilized newforms and $\omega_F^{1/2}$ is unramified outside p , we find that $\phi_v = \varphi_{1,v} \otimes \varphi_{2,v} \otimes \varphi_{3,v} \in \mathcal{V}_{\Pi_v}^{\text{new}}$ if $v \neq p$ and $\phi_p = \varphi_{1,p} \otimes \varphi_{2,p} \otimes \varphi_{3,p} \in \mathcal{V}_{\Pi_p}^{\text{ord}}$.

- $\varphi_{i,v} \in \mathcal{V}_{\pi_{i,v}}^{\text{new}}$ is a new vector if $v \neq p$,

- $\varphi_{i,p} \in \mathcal{V}_{\pi_i,p}^{\text{ord}}(\chi_{i,p})$ is an ordinary vector attached to the character $\chi_{i,p} : \mathbf{Q}_p^\times \rightarrow \mathbf{C}^\times$, where

$$(3.19) \quad \chi_{1,p} = \alpha_{f,p} \omega_{F,p}^{-1/2}, \quad \chi_{2,p} = \alpha_{g,p} \quad \text{and} \quad \chi_{3,p} = \alpha_{h,p}$$

($\alpha_{?,p}$ is the character attached to a p -stabilized newform ? defined in Remark 2.5).

For each finite prime ℓ , define the polynomial $\mathcal{Q}_{1,\ell}(X) \in \mathcal{O}[X]$ by

$$(3.20) \quad \mathcal{Q}_{1,\ell}(X) = X^{v_\ell(\mathbf{d}_f)} \begin{cases} 1 & \text{if } \ell \notin \Sigma_{f,0}^{(\text{Ib})}, \\ (1 - \omega_F^{1/2}(\varpi_\ell) \beta_\ell(f)^{-1} \ell^{\frac{k_1}{2}-1} X^{-1}) & \text{if } \ell \in \Sigma_{f,0}^{(\text{Ib})}. \end{cases}$$

Set $\mathcal{Q}_{2,\ell}(X) = \mathcal{Q}_{g,\ell}(X)$ and $\mathcal{Q}_{3,\ell}(X) = \mathcal{Q}_{h,\ell}(X)$. Let $\widehat{\mathbf{d}}_f = \prod_\ell \varpi_\ell^{v_\ell(\mathbf{d}_f)} \in \widehat{\mathbf{Q}}^\times$. We put

$$\begin{aligned} \varphi_1^* &:= \prod_\ell \mathcal{Q}_{1,\ell}(V_\ell) \varphi_1 = \omega_F^{1/2}(\widehat{\mathbf{d}}_f) \cdot \varphi_f^* \otimes \omega_F^{-1/2}, \\ \varphi_2^* &= \varphi_g^*; \quad \varphi_3^* = \varphi_h^*. \end{aligned}$$

We give the factorization of the automorphic form ϕ_F^* defined in (3.14). By definition,

$$\phi_F^* = C_1 \cdot \rho(\mathcal{J}_\infty) \varphi_1^* \boxtimes \varphi_2^* \boxtimes V_+^m \theta_p^k \varphi_3^* \quad (C_1 := \omega_{F,\infty}^{-1/2}(-1) \omega_F^{-1/2}(\widehat{\mathbf{d}}_f)).$$

In view of (3.9), we find that that $\phi_F^* = C_1 \cdot \otimes_v \phi_v^*$, where

$$(3.21) \quad \phi_v^* = \begin{cases} \pi_{1,\infty}(\mathcal{J}_\infty) \varphi_{1,\infty} \otimes \varphi_{2,\infty} \otimes V_+^m \varphi_{3,\infty} & \text{if } v = \infty, \\ \varphi_{1,p} \otimes \varphi_{2,p} \otimes \theta_p^k \varphi_{3,p} & \text{if } v = p, \\ \mathcal{Q}_{1,\ell}(V_\ell) \varphi_{1,\ell} \otimes \mathcal{Q}_{2,\ell}(V_\ell) \varphi_{2,\ell} \otimes \mathcal{Q}_{3,\ell}(V_\ell) \varphi_{3,\ell} & \text{if } v = \ell \nmid p. \end{cases}$$

Here θ_p^k is the local twisting operator attached to \mathbb{k} as in (2.12) and V_ℓ is the level-raising operator as in (2.11). Note that $\phi_\ell^* = \phi_\ell$ is a new vector in \mathcal{V}_{Π_ℓ} for $\ell \nmid pN$.

Next we consider the contragredient representation $\widetilde{\Pi} = \widetilde{\pi}_1 \otimes \widetilde{\pi}_2 \otimes \widetilde{\pi}_3$. We put

$$\widetilde{\varphi}_i = \varphi_i \otimes \omega_i^{-1} \quad \text{and} \quad \widetilde{\varphi}_i^* = \varphi_i^* \otimes \omega_i^{-1}, \quad i = 1, 2, 3.$$

Define $\widetilde{\phi}_F$ and $\widetilde{\phi}_F^* \in \mathcal{A}(\widetilde{\Pi})$ by

$$\begin{aligned} \widetilde{\phi}_F &= \widetilde{\varphi}_1 \boxtimes \widetilde{\varphi}_2 \boxtimes \widetilde{\varphi}_3, \\ \widetilde{\phi}_F^* &= \rho(\mathcal{J}_\infty) \widetilde{\varphi}_1^* \boxtimes \widetilde{\varphi}_2^* \boxtimes V_+^m \theta_p^k \widetilde{\varphi}_3^*. \end{aligned}$$

Recall that N_i is the tame conductor of π_i . Take an isomorphism $\mathcal{A}(\widetilde{\Pi}) \simeq \otimes_v \mathcal{V}_{\widetilde{\Pi}_v}$ with $\mathcal{V}_{\widetilde{\Pi}_v} = \mathcal{V}_{\widetilde{\pi}_{1,v}} \otimes \mathcal{V}_{\widetilde{\pi}_{2,v}} \otimes \mathcal{V}_{\widetilde{\pi}_{3,v}}$. We have a factorization $\widetilde{\phi}_F = \otimes_v \widetilde{\phi}_v$, where $\widetilde{\phi}_v = \widetilde{\varphi}_{1,v} \otimes \widetilde{\varphi}_{2,v} \otimes \widetilde{\varphi}_{3,v}$,

$$\begin{aligned} \widetilde{\phi}_{i,\infty} &\in \mathcal{V}_{\widetilde{\pi}_{i,\infty}}^{\text{new}}, \quad \widetilde{\phi}_{i,p} \in \mathcal{V}_{\widetilde{\pi}_{i,p}}^{\text{ord}}(\chi_{i,p} \omega_{Q_i}^{-1}); \\ \widetilde{\phi}_{i,v} &\in \widetilde{\pi}_{i,v} \left(\begin{pmatrix} 0 & 1 \\ -N_i & 0 \end{pmatrix} \right) \mathcal{V}_{\widetilde{\pi}_{i,v}}^{\text{new}} \quad \text{if } v \neq p, \infty. \end{aligned}$$

Moreover, $\widetilde{\phi}_F^* = \otimes_v \widetilde{\phi}_v^*$, where

$$(3.22) \quad \widetilde{\phi}_v^* = \begin{cases} \pi_{1,\infty}(\mathcal{J}_\infty) \widetilde{\varphi}_{1,\infty} \otimes \widetilde{\varphi}_{2,\infty} \otimes V_+^m \widetilde{\varphi}_{3,\infty} & \text{if } v = \infty, \\ \widetilde{\varphi}_{1,p} \otimes \widetilde{\varphi}_{2,p} \otimes \theta_p^k \widetilde{\varphi}_{3,p} & \text{if } v = p, \\ \widetilde{\mathcal{Q}}_{1,\ell}(V_\ell) \widetilde{\varphi}_{1,\ell} \otimes \widetilde{\mathcal{Q}}_{2,\ell}(V_\ell) \widetilde{\varphi}_{2,\ell} \otimes \widetilde{\mathcal{Q}}_{3,\ell}(V_\ell) \widetilde{\varphi}_{3,\ell} & \text{if } v = \ell \nmid p. \end{cases}$$

Here $\widetilde{\mathcal{Q}}_{i,\ell}(X) = \mathcal{Q}_{i,\ell}(\omega_i^{-1}(\varpi_\ell)X)$ for $i = 1, 2, 3$.

3.8.2. *Ichino's formula.* For $\underline{N} = (N_1, N_2, N_3)$, we put

$$\tau_{\underline{N}} = (\tau_{N_1}, \tau_{N_2}, \tau_{N_3}) \in \mathrm{GL}_2(\mathbf{A}_E).$$

Here τ_{N_i} is the matrix defined as in (2.16). For each place v of \mathbf{Q} , we choose a $\mathrm{GL}_2(E \otimes \mathbf{Q}_v)$ -equivariant map $\mathbf{b}_v : \mathcal{V}_{\Pi_v} \otimes \mathcal{V}_{\tilde{\Pi}_v} \rightarrow \mathbf{C}$ such that $\mathbf{b}_v(\phi_v, \phi_v) = 1$ for all but finitely many v . We introduce certain local zeta integrals that appear in our application of Ichino's formula. For each place v , we define the local zeta integral

$$(3.23) \quad I_v(\phi_v^* \otimes \tilde{\phi}_v^*) := \frac{L(1, \Pi_v, \mathrm{Ad})}{\zeta_v(2)^2 L(1/2, \Pi_v)} \int_{\mathrm{PGL}_2(\mathbf{Q}_v)} \frac{\mathbf{b}_v(\Pi_v(g_v)\phi_v^*, \tilde{\phi}_v^*)}{\mathbf{b}_v(\Pi_v(\tau_{\underline{N}, v})\phi_v, \tilde{\phi}_v)} dg_v.$$

Here dg_v is the Haar measure as in §2.4.1. At the place p , we will consider the local integral

$$(3.24) \quad I_p^{\mathrm{ord}}(\phi_p^* \otimes \tilde{\phi}_p^*, \mathbf{t}_n) := \frac{L(1, \Pi_p, \mathrm{Ad})}{\zeta_p(2)^2 L(1/2, \Pi_p)} \int_{\mathrm{PGL}_2(\mathbf{Q}_p)} \frac{\mathbf{b}_p(\Pi_p(g_p \mathbf{t}_n)\phi_p^*, \tilde{\Pi}_p(\mathbf{t}_n)\tilde{\phi}_p^*)}{\mathbf{b}_p(\Pi_p(\mathbf{t}_n)\phi_p, \tilde{\phi}_p)} dg_p.$$

Remark 3.8. The integrals $I_v(\phi_v^* \otimes \tilde{\phi}_v^*)$ and $I_p^{\mathrm{ord}}(\phi_p^* \otimes \tilde{\phi}_p^*, \mathbf{t}_n)$ do not depend any choice of the realizations $\mathcal{V}_{\Pi_v}, \mathcal{V}_{\tilde{\Pi}_v}$, the pairing \mathbf{b}_v and the new or ordinary vector ϕ_v in virtue of the irreducibility of Π_v and the multiplicity one for new vectors and ordinary vectors Proposition 2.2. This allows us to evaluate these local integrals by choosing favourable realizations of \mathcal{V}_{Π_v} .

Definition 3.9. Define the set

$$\Sigma_{fgh}^- = \{\ell \in \Sigma_f^0 \cap \Sigma_g^0 \cap \Sigma_h^0 \mid \varepsilon(1/2, \Pi_\ell) = -1\}.$$

From the rigidity of automorphic types in Remark 3.1, we can deduce that there is a subset Σ^- of primes dividing N such that

$$\Sigma^- = \Sigma_{\mathbf{f}_{Q_1} \mathbf{g}_{Q_2} \mathbf{h}_{Q_3}}^- = \left\{ \ell : \text{prime factors of } N \mid \varepsilon(\mathrm{WD}_\ell(\mathbf{V}_{\underline{Q}}^\dagger)) = -1 \right\}$$

for any arithmetic point $\underline{Q} \in \mathfrak{X}_{\mathcal{R}}^f$.

Proposition 3.10. *Suppose that $\Sigma^- = \emptyset$. Then*

$$\frac{I(\rho(\mathbf{t}_n)\phi_F^*)^2}{\prod_{i=1}^3 \langle \rho(\tau_{N_i} \mathbf{t}_n) \varphi_i, \tilde{\varphi}_i \rangle} = \frac{(-1)^{k_1} \zeta_{\mathbf{Q}}(2)}{8L(1, \Pi, \mathrm{Ad})} \cdot L\left(\frac{1}{2}, \Pi\right) \cdot I_p^{\mathrm{ord}}(\phi_p^* \otimes \tilde{\phi}_p^*, \mathbf{t}_n) \cdot \prod_{v \neq p} I_v(\phi_v^* \otimes \tilde{\phi}_v^*) \omega_{F, q}^{-1}(\mathbf{d}_f).$$

PROOF. Note that

$$I(\rho(\mathbf{t}_n)\phi_F^*)^2 = \omega_{1, \infty}(-1) I(\rho(\mathbf{t}_n)\phi_F^*) \cdot I(\rho(\mathbf{t}_n)\tilde{\phi}_F^*).$$

Applying [Ich08, Theorem 1.1, Remark 1.3], we obtain the proposition immediately in view of the decomposition of ϕ_F^* and $\tilde{\phi}_F^*$ into pure tensors. We remark that $\omega_{1, \infty}(-1) = (-1)^{k_1}$ and the product measure $\prod_v dg_v = \zeta_{\mathbf{Q}}(2) \cdot d^\tau g$ (cf. [III0, page 1403]), so the constant C in Remark 1.3 *loc.cit.* equals $\zeta_{\mathbf{Q}}(2)^{-1}$. \square

Lemma 3.11. (1) *If $q \nmid N$ is a finite prime, then $I_q(\phi_q^* \otimes \tilde{\phi}_q^*) = 1$;*

$$(2) \quad I_\infty(\phi_\infty^* \otimes \tilde{\phi}_\infty^*) = 2^{k_2 + k_3 - k_1 + 1}.$$

PROOF. Part (1) is [Ich08, Lemma 2.2]. Note that $\phi_q^* = \phi_q$ is a new vector in \mathcal{V}_{Π_q} for a finite prime $q \nmid N$. The formula of the archimedean zeta integral in part (2) is proved in [CC17]. For the reader's convenience, we sketch the proof. For $i = 1, 2, 3$, let $W_{k_i} = W_{\pi_i, \infty}$ be the Whittaker newform of the discrete series $\pi_{i, \infty} = \mathcal{D}_0(k_i)$ in (2.10). Define the matrix coefficient $\Phi_\infty : \mathrm{GL}_2(\mathbf{R}) \rightarrow \mathbf{C}$ by

$$\Phi_\infty(g) := \frac{\langle \rho(g \mathcal{J}_\infty) W_{k_1}, \rho(\mathcal{J}_\infty) W_{k_1} \rangle}{\langle \rho(\mathcal{J}_\infty) W_{k_1}, W_{k_1} \rangle} \cdot \frac{\langle \rho(g) W_{k_2}, W_{k_2} \rangle}{\langle \rho(\mathcal{J}_\infty) W_{k_2}, W_{k_2} \rangle} \cdot \frac{(8\pi)^{2m} \langle \rho(g) V_+^m W_{k_3}, V_+^m W_{k_3} \rangle}{\langle \rho(\mathcal{J}_\infty) W_{k_3}, W_{k_3} \rangle}$$

(recall that $m = \frac{k_1 - k_2 - k_3}{2}$). Note that Φ is right $\mathrm{SO}(2)(\mathbf{R})$ -invariant, and a lengthy computation shows that

$$\begin{aligned} \Phi_\infty\left(\begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix}\right) &= \mathbb{I}_{\mathbf{R}_+}(y) \cdot \frac{4^{k_1} \Gamma(k_3 + m)^2}{\Gamma(k_3)} \sum_{i, j=0}^m (-2)^{i+j} \binom{m}{i} \binom{m}{j} \frac{\Gamma(k_3 + i + j)}{\Gamma(k_3 + i) \Gamma(k_3 + j)} \\ &\quad \times \frac{(-y)^{k_1 - m + i}}{((1-a) - \sqrt{-1}x)^{k_1} ((1-y) + \sqrt{-1}x)^{k_1 - 2m + i + j}}. \end{aligned}$$

By definition,

$$I_\infty(\phi_\infty^* \otimes \tilde{\phi}_\infty^*) = \frac{L(1, \Pi_\infty, \text{Ad})}{\zeta_\infty(2)^2 L(1/2, \Pi_\infty)} \cdot (8\pi)^{-2m} I(\Phi_\infty),$$

where

$$I(\Phi_\infty) := \int_{\text{PGL}_2(\mathbf{R})} \Phi_\infty(g) dg = \int_{\mathbf{R}} \int_{\mathbf{R}^\times} \Phi_\infty\left(\begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix}\right) \frac{dy}{|y|} dx.$$

By a direct computation, we obtain

$$(3.25) \quad \begin{aligned} I(\Phi_\infty) &= \frac{4^{k_1} \Gamma(k_3 + m)^2}{\Gamma(k_3)} \sum_{i,j=0}^m (-2)^{i+j} \binom{m}{i} \binom{m}{j} \frac{\Gamma(k_3 + i + j)}{\Gamma(k_3 + i) \Gamma(k_3 + j)} \\ &\quad \times 2^{2-2k_1+2m-i-j} \cdot \pi \cdot \frac{\Gamma(k_1 - m + i - 1) \Gamma(k_3 - m + j)}{\Gamma(k_1 - 2m + i + j) \Gamma(k_1)} \\ &= \frac{4^{m+1} \pi \cdot \Gamma(k_3 + m)}{\Gamma(k_2) \Gamma(k_3)} \sum_{j=0}^m (-1)^j \binom{m}{j} \frac{\Gamma(k_1 - m + j)}{\Gamma(k_3 + j)} \cdot S_j, \end{aligned}$$

where

$$S_j := \Gamma(k_2 + m) \sum_{i=0}^m (-1)^i \binom{m}{i} \frac{\Gamma(k_2 + j + i)}{\Gamma(k_3 + i)} \cdot \frac{\Gamma(k_1 - m - 1 + i)}{\Gamma(k_1 - 2m_j + i)}.$$

Applying the combinatorial identity [Orl87, Lemma 3] to S_j , we find that

$$S_j = (-1)^m \cdot \frac{\Gamma(k_3 + j) \Gamma(k_1 - m - 1)}{\Gamma(k_1 - m + j)} \cdot \frac{\Gamma(k_1 - k_3 - m)}{\Gamma(k_1 - k_3 - 2m)} \cdot \frac{\Gamma(j + 1)}{\Gamma(j - m + 1)}.$$

Substituting the above expression to the last line of (3.25), we find that

$$I(\Phi_\infty) = 4^{m+1} \cdot \pi \cdot \frac{\Gamma(k_1 - m - 1) \Gamma(k_3 + m) \Gamma(k_2 + m) \Gamma(m + 1)}{\Gamma(k_1) \Gamma(k_2) \Gamma(k_3)}.$$

Hence, part (2) follows from the above expression of $I(\Phi_\infty)$ and

$$\frac{L(1, \Pi_\infty, \text{Ad})}{\zeta_\infty(2)^2 L(1/2, \Pi_\infty)} = \frac{\pi^{-3} \Gamma_{\mathbf{C}}(k_1) \Gamma_{\mathbf{C}}(k_2) \Gamma_{\mathbf{C}}(k_3)}{\pi^{-2} \cdot \Gamma_{\mathbf{C}}(k_1 - m - 1) \Gamma_{\mathbf{C}}(k_3 + m) \Gamma_{\mathbf{C}}(k_2 + m) \Gamma_{\mathbf{C}}(m + 1)}. \quad \square$$

To distinguish the contributions from each term in the formula of $\mathcal{L}_{\mathbf{F}}^f(\underline{Q})$, we introduce the normalized local zeta integrals. For each place v , define the local norm of Whittaker newforms for Π_v by

$$(3.26) \quad B_{\Pi_v} := B_{\pi_{1,v}} B_{\pi_{2,v}} B_{\pi_{3,v}}$$

with $B_{\pi_{i,v}}$ the local norm of $\pi_{i,v}$ defined as in (2.17). To each positive integer n , we associate the local norm $B_{\Pi_p^{\text{ord}}}^{[n]}$ of ordinary Whittaker functions for Π_p given by

$$(3.27) \quad B_{\Pi_p^{\text{ord}}}^{[n]} := \frac{\zeta_p(2)^3}{\zeta_p(1)^3 L(1, \Pi_p, \text{Ad})} \prod_{i=1}^3 \langle \rho(t_n) W_{\pi_{i,p}}^{\text{ord}}, W_{\pi_{i,p}}^{\text{ord}} \otimes \omega_{i,p}^{-1} \rangle.$$

We define the normalized local zeta integrals

$$(3.28) \quad \mathcal{I}_{\Pi_{\underline{Q},p}}^{\text{unb}} = I_p^{\text{ord}}(\phi_p^* \otimes \tilde{\phi}_p^*, \mathbf{t}_n) \cdot \frac{B_{\Pi_p^{\text{ord}}}^{[n]}}{\omega_{f,p}^{-1} \alpha_{f,p}^2 \cdot | \cdot |_p (-p^{2n})} \cdot \frac{\zeta_p(1)^2}{\zeta_p(2)^2};$$

$$(3.29) \quad \mathcal{I}_{\Pi_{\underline{Q},q}}^* = I_q(\phi_q^* \otimes \tilde{\phi}_q^*) \cdot B_{\Pi_q} \cdot \frac{\zeta_q(1)^2}{|N|_q^2 \zeta_q(2)^2} \cdot \omega_{F,q}^{-1}(\mathbf{d}_f) | \mathbf{d}_F^k |_q \text{ for } q \mid N.$$

Definition 3.12 (The canonical periods of Hida families). Define the canonical period $\Omega_{\mathbf{f}_Q}$ of the specialization \mathbf{f}_Q at an arithmetic point Q by

$$\Omega_{\mathbf{f}_Q} := 2^{k_Q} \cdot \| \mathbf{f}_Q^\circ \|_{\Gamma_0(N_Q)}^2 \cdot \frac{\mathcal{E}_p(\mathbf{f}_Q, \text{Ad})}{\eta_{\mathbf{f}_Q}},$$

where \mathbf{f}_Q° is the normalized newform associated with \mathbf{f}_Q of conductor N_Q and $\eta_{\mathbf{f}_Q}$ is the specialization of $\eta_{\mathbf{f}}$ at Q and $\mathcal{E}_p(\mathbf{f}_Q, \text{Ad})$ is the modified Euler factor in (3.10).

We summarize our computation in the following

Corollary 3.13. *Assume that $\Sigma^- = \emptyset$. For every $\underline{Q} = (Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathcal{R}}^f$, we have the interpolation formula*

$$\mathcal{L}_{\mathbf{F}}^f(\underline{Q})^2 = \frac{1}{4} \cdot \frac{L(1/2, \Pi_{\underline{Q}})}{(-1)^{k_{Q_1}} \Omega_{\mathbf{f}_{Q_1}}^2} \cdot \mathcal{I}_{\Pi_{\underline{Q}, p}}^{\text{unb}} \cdot \prod_{q|N} \mathcal{I}_{\Pi_{\underline{Q}, q}}^*.$$

PROOF. By Waldspurger's formula Proposition 2.7 and the equations

$$B_{\Pi_{\infty}} = 2^{-(k_1+k_2+k_3)-3}; \quad B_{\Pi_q} = 1 \text{ if } q \nmid N,$$

we have

$$\prod_{i=1}^3 \langle \rho(\tau_{N_i} t_n) \varphi_i, \tilde{\varphi}_i \rangle = \frac{8L(1, \Pi, \text{Ad})}{\zeta_{\mathbf{Q}}(2)^3} \cdot 2^{-(k_1+k_2+k_3)-3} B_{\Pi_p}^{[n]} \prod_{q|N} B_{\Pi_q}.$$

We thus obtain the corollary by combining Proposition 3.7, Proposition 3.10, Lemma 3.11 and

$$[\text{SL}_2(\mathbf{Z}) : \Gamma_0(N)] = \prod_{q|N} \frac{\zeta_q(1)}{|N|_q \zeta_q(2)}. \quad \square$$

4. THE BALANCED p -ADIC TRIPLE PRODUCT L -FUNCTIONS

4.1. Notation and conventions. Let D be a definite quaternion algebra over \mathbf{Q} of the absolute discriminant of N^- . Let $\nu : D^\times \rightarrow \mathbf{Q}^\times$ be the reduced norm. For any commutative \mathbf{Q} -algebra R , put

$$D^\times(R) = (D \otimes_{\mathbf{Q}} R)^\times.$$

If v is a place of \mathbf{Q} , let $D_v = D \otimes_{\mathbf{Q}} \mathbf{Q}_v$. For $x \in D^\times(\mathbf{A})$, denote by $x_v \in D_v^\times$ the local component of x at v . We fix an isomorphism $\Psi = \prod_{q \nmid N^-} \Psi_q : D^\times(\widehat{\mathbf{Q}}^{(N^-)}) \simeq \text{M}_2(\widehat{\mathbf{Q}}^{(N^-)})$ once and for all. Let \mathcal{O}_D be the maximal order of D such that $\Psi_q(\mathcal{O}_D \otimes \mathbf{Z}_q) = \text{M}_2(\mathbf{Z}_q)$ for all primes $q \nmid \infty N^-$. Let N^+ be a positive integer prime to N^- and let

$$N = N^+ N^-.$$

Denote by R_N the Eichler order of level N^+ in D with respect to Ψ . Put

$$U_1(N) = \left\{ g = (g_q)_q \in \widehat{R}_N^\times \mid \Psi_q(b_q) \equiv \begin{pmatrix} * & * \\ 0 & 1 \end{pmatrix} \pmod{N\mathbf{Z}_q} \text{ for } q \mid N^+ \right\}.$$

We shall frequently use the following notation in this section: let $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\widehat{\mathbf{Q}}^{(N^-)})$ act on $x \in \widehat{D}^\times$ by

$$x \begin{pmatrix} a & b \\ c & d \end{pmatrix} := x \cdot \Psi^{-1} \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right).$$

Let $d^r x$ be the Tamagawa measure on $\mathbf{A}^\times \backslash D^\times(\mathbf{A})$ with the volume $\text{vol}(\mathbf{A}^\times \backslash D^\times(\mathbf{A}), d^r x) = 2$. There exists a positive rational number $\text{vol}(\widehat{R}_N^\times)$ such that for any $f \in L^1(D^\times \backslash D^\times(\mathbf{A}) / D_\infty^\times \widehat{R}_N^\times)$, we have

$$(4.1) \quad \int_{\mathbf{A}^\times \backslash D^\times \backslash D^\times(\mathbf{A})} f(x) d^r x = \text{vol}(\widehat{R}_N^\times) \sum_{[x] \in D^\times \backslash \widehat{D}^\times / \widehat{R}_N^\times} f(x) \cdot (\#\Gamma_{N,x})^{-1},$$

where $[x]$ means the double coset $D^\times x \widehat{R}_N^\times$ and $\Gamma_{N,x} := (D^\times \cap x \widehat{R}_N^\times x^{-1}) \mathbf{Q}^\times / \mathbf{Q}^\times$. By Eichler's mass formula, we have

$$(4.2) \quad \begin{aligned} \text{vol}(\widehat{R}_N^\times) &= \frac{48}{N} \prod_{q|N^-} \zeta_q(1) \prod_{q|N^+} (1+q^{-1})^{-1} \\ &= \frac{48}{[\text{SL}_2(\mathbf{Z}) : \Gamma_0(N)]} \prod_{q|N^-} \frac{1+q^{-1}}{1-q^{-1}}. \end{aligned}$$

For a non-negative integer κ and a commutative ring A , let $L_\kappa(A) := A[X, Y]_{\deg=\kappa}$ be the space of two variable polynomials of degree κ over A . Let $\rho_\kappa : M_2(A) \rightarrow \text{End}_A L_\kappa(A)$ be the morphism $\rho_\kappa(g)P(X, Y) = P((X, Y)g)$. Let $\langle \cdot, \cdot \rangle_\kappa : L_\kappa(A) \times L_\kappa(A) \rightarrow A[\frac{1}{\kappa!}]$ be the pairing defined by

$$\langle X^i Y^{\kappa-i}, X^j Y^{\kappa-j} \rangle = \begin{cases} (-1)^i \binom{\kappa}{i}^{-1} & \text{if } i+j = \kappa, \\ 0 & \text{if } i+j \neq \kappa. \end{cases}$$

Let $g \mapsto g'$ be the main involution of $M_2(A)$ given by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}' = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

It is well-known that

$$(4.3) \quad \langle \rho_\kappa(g)P_1, P_2 \rangle_\kappa = \langle P_1, \rho_\kappa(g')P_2 \rangle_\kappa.$$

In this section, we shall freely identify Dirichlet characters χ with finite order Hecke characters $\chi_{\mathbf{A}}$ when no confusion may arise.

4.2. p -adic modular forms on definite quaternion algebras. Let $\mathcal{O} \subset \mathcal{O}_{\mathbf{C}_p}$ be a finite flat extension of \mathbf{Z}_p containing all $\phi(N)$ -th roots of unity. For an \mathcal{O} -algebra A and a A -valued (even) Hecke character $\chi : \mathbf{Q}^\times \backslash \widehat{\mathbf{Q}}^\times \rightarrow A^\times$, we let $\mathcal{S}_{\kappa+2}^D(N, \chi, A)$ be the space of p -adic modular forms on \widehat{D}^\times of weight $\kappa+2$, level N and branch character χ , consisting of vector-valued functions $f : \widehat{D}^\times \rightarrow L_\kappa(A)$ such that

$$f(\alpha x u z) = \rho_{\kappa,p}(u_p^{-1})f(x)z_p^{-\kappa}\chi^{-1}(z) \text{ for all } \alpha \in D^\times, u \in U_1(N^+), z \in \widehat{\mathbf{Q}}^\times.$$

Here u_p is the p -component of u and $\rho_{\kappa,p}(u_p) = \rho_\kappa(\Psi_p(u_p))$. For each integer d prime to pN^- , define the level raising operator $V_d : \mathcal{S}_{\kappa+2}^D(N, \chi, A) \rightarrow \mathcal{S}_{\kappa+2}^D(Nd, \chi, A)$ by

$$V_d f(x) = f\left(x \begin{pmatrix} d^{-1} & 0 \\ 0 & 1 \end{pmatrix}\right).$$

We recall the Hecke operators T_q and the operators \mathbf{U}_q acting on $f \in \mathcal{S}_{\kappa+2}^D(N, \chi, A)$. For each prime $q \mid N^-$, let $\varpi_{D_q} \in R_q^\times$ with $\nu(\varpi_{D_q}) = q$. The Hecke operator T_q for $q \nmid Np$ is given by

$$T_q f(x) = f\left(x \begin{pmatrix} 1 & 0 \\ 0 & \varpi_q \end{pmatrix}\right) + \sum_{b \in \mathbf{Z}_q/q\mathbf{Z}_q} f\left(x \begin{pmatrix} \varpi_q & b \\ 0 & 1 \end{pmatrix}\right)$$

and the operator \mathbf{U}_q for $q \mid MN^-p$ is given by

$$\begin{aligned} \mathbf{U}_q f(x) &= \sum_{b \in \mathbf{Z}_q/q\mathbf{Z}_q} f\left(x \begin{pmatrix} \varpi_q & b \\ 0 & 1 \end{pmatrix}\right) \text{ for } q \mid M, q \neq p, \quad \mathbf{U}_q f(x) = f(x\varpi_{D_q}) \text{ for } q \mid N^-, \\ \mathbf{U}_p f(x) &= \sum_{b \in \mathbf{Z}_p/p\mathbf{Z}_p} \rho_{\kappa,p}\left(\begin{pmatrix} \varpi_p & b \\ 0 & 1 \end{pmatrix}\right) f\left(x \begin{pmatrix} \varpi_p & b \\ 0 & 1 \end{pmatrix}\right). \end{aligned}$$

Here $\varpi_q = (\varpi_{q,\ell}) \in \widehat{\mathbf{Q}}^{(N^-)^\times}$ is the idele $\varpi_{q,q} = q$ and $\varpi_{q,\ell} = 1$ for $\ell \nmid N^-q$. If A is p -adically complete, then the ordinary projector $e = \lim_{n \rightarrow \infty} \mathbf{U}_p^{n!}$ converges to an idempotent in $\text{End}_{\mathcal{O}} \mathcal{S}_{\kappa+2}^D(N, \chi, A)$.

Inner products. Denote by $\varepsilon_{\text{cyc}} : \mathbf{Q}_+ \backslash \widehat{\mathbf{Q}}^\times \rightarrow \mathbf{Z}_p^\times$ the p -adic cyclotomic character defined by $\varepsilon_{\text{cyc}}(a) = |a|_{\mathbf{A}} a_p$. Assuming $6 \cdot \kappa! \in A^\times$, we have a perfect pairing

$$(\cdot, \cdot)_N : \mathcal{S}_{\kappa+2}^D(N, \chi, A) \times \mathcal{S}_{\kappa+2}^D(N, \chi^{-1}, A) \rightarrow A$$

given by

$$(f_1, f_2)_N := \sum_{[x] \in D^\times \backslash \widehat{D}^\times / \widehat{R}_N^\times} \langle f_1(x), f_2(x) \rangle_\kappa \cdot \varepsilon_{\text{cyc}}(\nu(x))^\kappa \cdot (\#\Gamma_{N,x})^{-1}.$$

Let $\tau_N^D = (\tau_{N,q}^D) \in \widehat{D}^\times$ be the element with $\tau_{N,q}^D = 1$ if $q \nmid N$ and $\tau_{N,q}^D = \Psi_q^{-1}\left(\begin{pmatrix} 0 & 1 \\ -N & 0 \end{pmatrix}\right)$ for $q \mid N^+$. Define the Atkin-Lehner involution $[\tau_N^D] : \mathcal{S}_{\kappa+2}^D(N, \chi, A) \rightarrow \mathcal{S}_{\kappa+2}^D(N, \chi^{-1}, A)$ by

$$[\tau_N^D]f(x) := \rho_{\kappa,p}(\tau_{N,p}^D)f(x\tau_N^D)\chi(\nu(x)).$$

We can define a new pairing $\langle \cdot, \cdot \rangle_N : \mathcal{S}_{\kappa+2}^D(N, \chi, A) \times \mathcal{S}_{\kappa+2}^D(N, \chi, A) \rightarrow A$ by

$$\langle f_1, f_2 \rangle_N = (f_1, [\tau_N^D]f_2)_N.$$

It is well known that this new pairing is Hecke equivariant and perfect (cf. [Hid06, Lemma 3.5]).

4.3. Automorphic forms on definite quaternion algebras. Fixing $\iota_p : \mathbf{C}_p \simeq \mathbf{C}$ once and for all, we choose an imbedding $\Psi_\infty : D_\infty \hookrightarrow M_2(\mathbf{C})$ such that $\Psi_\infty(\alpha) = \iota_p(\Psi_p(\alpha))$ for $\alpha \in D^\times$. Define the unitarized representation $\rho_\kappa^u : D_\infty^\times \rightarrow \text{Aut } L_\kappa(\mathbf{C})$ by $\rho_\kappa^u(x)P = |\nu(g)|_{\mathbf{A}}^{\kappa/2} \rho_\kappa(\Psi_\infty(g))P$ for $P \in L_\kappa(\mathbf{C})$.

For a finite order Hecke character ω modulo N^+ , let $\mathcal{A}_{\kappa+2}^D(N, \omega)$ be the space of $L_\kappa(\mathbf{C})$ -valued automorphic forms on $D^\times(\mathbf{A})$ of weight $\kappa + 2$, level N and character ω . In other words, $\mathcal{A}_{\kappa+2}^D(N, \omega)$ consists of functions $\varphi : D^\times(\mathbf{A}) \rightarrow L_\kappa(\mathbf{C})$ such that

$$\begin{aligned} \varphi(\alpha x u_\infty u_f z) &= \rho_\kappa^u(u_\infty^{-1})\varphi(x)\omega(z) \\ (\alpha \in D^\times, u_\infty \in D_\infty^\times, u_f \in U_1(N), z \in \mathbf{A}^\times). \end{aligned}$$

Here x_f denotes the finite part of x . To each p -adic modular form $f \in \mathcal{S}_{\kappa+2}^D(N, \chi, \mathcal{O})$, we associate the vector-valued automorphic form $\Phi(f) \in \mathcal{A}_{\kappa+2}^D(N, \chi^{-1})$ defined by

$$(4.4) \quad \Phi(f)(x) := \rho_\kappa(\Psi_\infty(x_\infty^{-1}))\iota_p(\rho_{\kappa,p}(x_p)f(x_f)) \cdot |\nu(x)|_{\mathbf{A}}^{\kappa/2}, \quad x \in D_{\mathbf{A}}^\times.$$

Let $\mathcal{A}^D(\omega)$ be the space of (scalar-valued) automorphic forms on $D^\times(\mathbf{A})$ with central character ω . For $\varphi, \varphi' \in \mathcal{A}^D(\omega)$, define

$$\langle \varphi, \varphi' \rangle = \int_{\mathbf{A}^\times D^\times \backslash D^\times(\mathbf{A})} \varphi(x)\varphi'(x)\omega^{-1}(\nu(x))d^\tau x.$$

Here $d^\tau x$ is the Tamagawa measure on $\mathbf{A}^\times \backslash D^\times(\mathbf{A})$. For $f \in \mathcal{S}_{\kappa+2}^D(N, \omega^{-1}, \mathbf{C}_p)$ and $\mathbf{u} \in L_\kappa(\mathbf{C}_p)$, let $\Phi(f)_{\mathbf{u}} \in \mathcal{A}^D(\omega)$ be the automorphic form given by $\Phi(f)_{\mathbf{u}}(x) := \langle \Phi(f)(x), \mathbf{u} \rangle_\kappa$. By (4.1) and Schur's orthogonal relation, we have

$$(4.5) \quad \langle \rho(\tau_N^D)\Phi(f)_{\mathbf{u}}, \Phi(f)_{\mathbf{v}} \rangle = \frac{\text{vol}(\widehat{R}_N^\times)}{(N^+)^{\kappa/2}(1+\kappa)} \cdot \langle f, f \rangle_N \cdot \langle \mathbf{u}, \mathbf{v} \rangle_\kappa.$$

4.4. Hida theory for quaternionic modular forms. In this subsection, we recall Hida theory for modular forms on definite quaternion algebras following [Hid88b]. Suppose that $p \nmid N$. For each positive integer α , let X_α be the finite set

$$X_\alpha = D^\times \backslash \widehat{D}^\times / U_1(Np^\alpha)$$

and let $\mathcal{O}[X_\alpha] = \bigoplus_{x \in X_\alpha} \mathcal{O}x$ be the finitely generated \mathcal{O} -module spanned by divisors of X_α . Recall that $\Lambda = \mathcal{O}[[1+p\mathbf{Z}_p]] = \mathcal{O}[[T]]$, where $T = \langle 1+p \rangle_\Lambda - 1$. For $z \in 1+p\mathbf{Z}_p$, let $\langle z \rangle_\Lambda$ act on $\mathcal{O}[X_\alpha]$ by $\langle z \rangle_\Lambda x := x \begin{pmatrix} z & 0 \\ 0 & z \end{pmatrix}$. Let $\Delta = (\mathbf{Z}/pN^+\mathbf{Z})^\times$. For $d \in \Delta$, the diamond operator σ_d acts on $\mathcal{O}[X_\alpha]$ as follows: decomposing $d = (d_1, d_2) \in (\mathbf{Z}/p\mathbf{Z})^\times \times (\mathbf{Z}/N^+\mathbf{Z})^\times$ and choosing an idele $\tilde{d} \in \widehat{\mathbf{Z}}^\times$ such that the p -component $\tilde{d}_p = \omega(d_1) \in \mathbf{Z}_p^\times$ is the Teichmüller lifting of d_1 and the prime-to- p component $\tilde{d}^{(p)} \in \widehat{\mathbf{Z}}^{(p)\times}$ is a lifting of d_2 , we define $\sigma_d x := x\tilde{d}$. Thus $\mathcal{O}[X_\alpha]$ is a finitely generated $\Lambda[\Delta]$ -module. Moreover, $\mathcal{O}[X_\alpha]$ is equipped with the usual Hecke operators T_q for $q \nmid Np$ given by

$$T_q x = x \begin{pmatrix} 1 & 0 \\ 0 & \varpi_q \end{pmatrix} + \sum_{b \in \mathbf{Z}_q/q\mathbf{Z}_q} x \begin{pmatrix} \varpi_q & b \\ 0 & 1 \end{pmatrix},$$

the operator \mathbf{U}_q for $q \mid Np$ defined by

$$\mathbf{U}_q x = \sum_{b \in \mathbf{Z}_q/q\mathbf{Z}_q} x \begin{pmatrix} \varpi_q & b \\ 0 & 1 \end{pmatrix} \text{ if } q \mid N^+p; \quad \mathbf{U}_q x = x\varpi_{D_q} \text{ if } q \mid N^-.$$

The ordinary projector $e = \varprojlim_n \mathbf{U}_p^n$ converges to an idempotent in $\text{End}_\Lambda(\mathcal{O}[X_\alpha])$.

We introduce the space of Λ -adic modular forms on definite quaternion algebras. Let $X_\infty := D^\times \backslash \widehat{D}^\times / U_1(Np^\infty)$, where

$$U_1(Np^\infty) = \left\{ g \in U_1(N) \mid g_p = \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}, a \in \mathbf{Z}_p^\times, b \in \mathbf{Z}_p \right\}.$$

We have a natural quotient map $X_\infty \rightarrow X_\beta \rightarrow X_\alpha$ for $\beta > \alpha$. Let P_α be the principal ideal of Λ generated by $(1+T)^{p^\alpha} - 1$.

Definition 4.1. Denote by $\mathbf{S}^D(N, \Lambda)$ the space of functions $\mathbf{f}: X_\infty \rightarrow \Lambda$ such that

- $\mathbf{f}(xz) = \mathbf{f}(x) \langle z \rangle^2 \langle z \rangle_\Lambda^{-1}$ for $z \in 1 + p\mathbf{Z}_p$;
- for any α sufficiently large, the function $\mathbf{f} \pmod{P_\alpha}: X_\infty \rightarrow \Lambda/P_\alpha$ factors through X_α .

We call $\mathbf{S}^D(N, \Lambda)$ the space of Λ -adic modular forms on D^\times of level N .

By definition, we have

$$(4.6) \quad \mathbf{S}^D(N, \Lambda) = \varprojlim_{\alpha} \mathrm{Hom}_{\Lambda}(\mathcal{O}[X_\alpha], \Lambda/P_\alpha) \otimes_{\Lambda, \iota_2} \Lambda,$$

where $\iota_2: \Lambda \rightarrow \Lambda$ is the \mathcal{O} -algebra homomorphism given by $\iota_2(T) = (1+T)^{-2}(1+p)^2 - 1$. It follows that we can make $\mathbf{S}^D(N, \Lambda)$ a compact Λ -module with the natural Hecke action given by $t\mathbf{f}(x) = \mathbf{f}(tx)$ for $t = T_q, \mathbf{U}_q$ and the diamond operators σ_d . In addition, the ordinary projector $e = \varprojlim_n \mathbf{U}_p^{n!}$ converges in $\mathrm{End}_{\Lambda} \mathbf{S}^D(N, \Lambda)$.

For a finite order Hecke character $\chi: \mathbf{Q}^\times \backslash \widehat{\mathbf{Q}}^\times \rightarrow \mathcal{O}^\times$ modulo N^+p , put

$$\begin{aligned} \mathbf{S}^D(N, \chi, \Lambda) &:= \{ \mathbf{f} \in \mathbf{S}^D(N, \Lambda) \mid \sigma_d \mathbf{f} = \chi^{-1}(d) \mathbf{f} \text{ for } d \in \Delta^\times \} \\ &= \left\{ \mathbf{f} \in \mathbf{S}^D(N, \Lambda) \mid \mathbf{f}(xz) = \mathbf{f}(x) \cdot \chi^{-1}(z) \langle \epsilon_{\mathrm{cyc}}(z) \rangle^2 \langle \epsilon_{\mathrm{cyc}}(z) \rangle_\Lambda^{-1} \text{ for } z \in \widehat{\mathbf{Q}}^\times \right\}. \end{aligned}$$

Let \mathbf{I} be a normal domain finite flat over Λ . We define $\mathbf{S}^D(N, \mathbf{I}) = \mathbf{S}^D(N, \Lambda) \otimes_{\Lambda} \mathbf{I}$ and $\mathbf{S}^D(N, \chi, \mathbf{I}) = \mathbf{S}^D(N, \chi, \Lambda) \otimes_{\Lambda} \mathbf{I}$.

Theorem 4.2 (Control Theorem). *Let $N_\chi := \sum_{d \in \Delta} \chi(d) \sigma_d \in \mathcal{O}[\Delta]$ and let P_χ be the ideal of $\Lambda[\Delta]$ generated by $\{\chi(d) \cdot \sigma_d - 1\}_{d \in \Delta}$. Suppose that $p > 3$. Then*

- (1) $\mathbf{S}^D(N, \chi, \mathbf{I})$ is a free \mathbf{I} -module, and the map $N_\chi: e\mathbf{S}^D(N, \mathbf{I})/P_\chi \simeq e\mathbf{S}^D(N, \chi, \mathbf{I})$ is an isomorphism.
- (2) For every arithmetic point $Q \in \mathfrak{X}_{\mathbf{I}}^+$, we have a Hecke equivariant isomorphism

$$\begin{aligned} e\mathbf{S}^D(N, \chi, \mathbf{I}) \otimes_{\mathbf{I}} \mathbf{I}/\wp_Q &\simeq e\mathcal{S}_{k_Q}^D(Np^\alpha, \chi \omega^{2-k_Q} \epsilon_Q, \mathcal{O}(Q)), \\ \mathbf{f} \pmod{\wp_Q} &\mapsto \mathbf{f}_Q, \end{aligned}$$

where $\alpha = \max\{1, c_p(\epsilon_Q)\}$ and \mathbf{f}_Q is the unique p -adic modular form such that

$$\mathbf{Q}(\mathbf{f}(x)) = \langle \mathbf{f}_Q(x), X^{k_Q-2} \rangle_{k_Q-2} \text{ for all } x \in \widehat{D}^\times.$$

PROOF. This is a reformation of Hida's control theorems for definite quaternion algebra. We give a sketch of proofs in [Hid88b] for the reader's convenience. We may assume $\mathbf{I} = \Lambda$ and $\mathcal{O} = \mathcal{O}(Q)$. Let Δ_p be the p -Sylow subgroup of Δ . We first show that $e\mathbf{S}^D(N, \Lambda)$ is a free $\Lambda[\Delta_p]$ -module. For any abelian group A , let $\mathrm{H}^0(X_\alpha, A)$ be the space of A -valued functions on X_α . Let $\mathcal{V}^{\mathrm{ord}}(N) := \varinjlim_{\alpha} \varprojlim_{\beta} e\mathrm{H}^0(X_\alpha, p^{-\beta}\mathcal{O}/\mathcal{O})$ be the discrete Λ -module $\mathcal{V}_0^{\mathrm{ord}}(0; U_1(N^+))$ defined in [Hid88b, Theorem 8.6]. Let $V^{\mathrm{ord}}(N) := \varprojlim_{\alpha} e \cdot \mathcal{O}[X_\alpha]$ be the Pontryagin dual of $\mathcal{V}^{\mathrm{ord}}(N)$. In virtue of (4.6),

$$e\mathbf{S}^D(N, \Lambda) = \mathrm{Hom}_{\Lambda}(V^{\mathrm{ord}}(N), \Lambda) \otimes_{\Lambda, \iota_2} \Lambda,$$

so it suffices to show that $V^{\mathrm{ord}}(N)$ is a free $\Lambda[\Delta_p]$ -module. For any positive integer α and character $\xi: (\mathbf{Z}/N^+p^\alpha)^\times \rightarrow \mathcal{O}_K^\times$ of p -power order with value in some finite extension K of $\mathrm{Frac} \mathcal{O}$, we define the \mathcal{O}_K -module

$$\mathrm{H}^0(X_\alpha, \xi, A) := \left\{ f \in \mathrm{H}^0(X_\alpha, A) \mid f(xz) = \xi(z)f(x), x \in X_\alpha, z \in \widehat{\mathbf{Z}}^\times \right\}, A = K/\mathcal{O}_K \text{ or } \mathcal{O}_K.$$

Since any finite order element in D^\times has order only divisible by 2 or 3 and $p > 3$, one verifies that the group $D^\times \cap xU_1(Np^\alpha)x^{-1} = \{1\}$ for any $x \in \widehat{D}^\times$ and that

$$\mathrm{H}^0(X_\alpha, \xi, K/\mathcal{O}_K) = \mathrm{H}^0(X_\alpha, \xi, \mathcal{O}_K) \otimes K/\mathcal{O}_K.$$

In particular, $\mathrm{H}^0(X_\alpha, \xi, K/\mathcal{O}_K)$ is p -divisible. Hence, the $\Lambda[\Delta_p]$ -freeness of $V^{\mathrm{ord}}(N)$ follows from [Hid88b, Corollary 10.1] (and the proof therein). From the $\Lambda[\Delta_p]$ -freeness of $e\mathbf{S}^D(N, \Lambda)$, we deduce that the map $\mathbf{f} \mapsto N_\chi \mathbf{f}$ induces an isomorphism

$$N_\chi: e\mathbf{S}^D(N, \Lambda)/P_\chi \simeq e\mathbf{S}^D(N, \Lambda)^{N_\chi=1} = e\mathbf{S}^D(N, \chi, \Lambda).$$

This proves part (1). We proceed to prove part (2). By [Hid88b, Theorem 9.4], we see that

$$e\mathbf{S}^D(N, \chi, \Lambda)/\wp_Q \simeq e\mathbf{S}^D(N, \Lambda)/(P_\chi, \wp_Q) \simeq e\mathcal{S}_{k_Q}^D(Np^n, \chi\omega^{2-k_Q}\epsilon_Q, \mathcal{O}).$$

The above isomorphism $\mathbf{f} \mapsto \mathbf{f}_Q$ is given by the dual map to the one ι in [Hid88b, (8.10)], whose explicit description is given in [Hid88b, line 9-11, page 375]. This finishes the proof of part (2). \square

A perfect pairing on the space of ordinary Λ -adic forms. For each positive integer α , put

$$X_0(Np^\alpha) = D^\times \backslash \widehat{D}^\times / \widehat{R}_{Np^\alpha}^\times.$$

To each finite order character $\chi : \mathbf{Q}^\times \backslash \widehat{\mathbf{Q}}^\times \rightarrow \mathcal{O}^\times$, we associate a universal \mathbf{I} -adic deformation defined by

$$\chi_{\mathbf{I}} : \mathbf{Q}^\times \backslash \widehat{\mathbf{Q}}^\times \rightarrow \mathbf{I}^\times, \quad \chi_{\mathbf{I}}(z) := \chi(z) \langle \varepsilon_{\text{cyc}}(z) \rangle^{-2} \langle \varepsilon_{\text{cyc}}(z) \rangle_{\mathbf{I}}.$$

For $\mathbf{f}, \mathbf{f}' \in e\mathbf{S}^D(N, \chi, \mathbf{I})$, put

$$\mathbf{B}_{N, \alpha}(\mathbf{f}, \mathbf{f}') := \sum_{[x] \in X_0(Np^\alpha)} \mathbf{U}_p^{-\alpha} \mathbf{f}(x\tau_{Np^\alpha}^D \mathbf{f}'(x)) \chi_{\mathbf{I}}(\nu(x)) \cdot (\#\Gamma_{Np^\alpha, x})^{-1} \pmod{P_\alpha} \in \mathbf{I}/P_\alpha.$$

One verifies that $\mathbf{B}_{N, \alpha+1}(\mathbf{f}, \mathbf{f}') \equiv \mathbf{B}_{N, \alpha}(\mathbf{f}, \mathbf{f}') \pmod{P_\alpha}$.

Definition 4.3. Let

$$\mathbf{B}_N : e\mathbf{S}^D(N, \chi, \mathbf{I}) \times e\mathbf{S}^D(N, \chi, \mathbf{I}) \rightarrow \mathbf{I}$$

be the Hecke-equivariant \mathbf{I} -bilinear pairing defined by

$$\mathbf{B}_N(\mathbf{f}, \mathbf{f}') := \varprojlim_{\alpha} \mathbf{B}_{N, \alpha}(\mathbf{f}, \mathbf{f}') \in \varprojlim_{\alpha} \mathbf{I}/P_\alpha = \mathbf{I}.$$

For every $Q \in \mathfrak{X}_{\mathbf{I}}^+$ with $k_Q = 2$, we have

$$\mathbf{B}_N(\mathbf{f}, \mathbf{f}')(Q) = \langle \mathbf{U}_p^{-\alpha} \mathbf{f}_Q, \mathbf{f}'_Q \rangle_{Np^\alpha}$$

for any $\alpha \geq \max\{1, c_p(\epsilon_Q)\}$. This in particular implies that the pairing \mathbf{B}_N is perfect.

Lemma 4.4. *For each arithmetic point Q in $\mathfrak{X}_{\mathbf{I}}^+$ and integer $\alpha \geq \max\{1, c_p(\epsilon_Q)\}$, we have*

$$\mathbf{B}_N(\mathbf{f}, \mathbf{f})(Q) = \langle \mathbf{U}_p^{-\alpha} \mathbf{f}_Q, \mathbf{f}_Q \rangle_{Np^\alpha}.$$

PROOF. Let $\kappa = k_Q - 2$ and let $\chi_Q = \chi_{\mathbf{I}} \pmod{Q} = \chi\omega^{-\kappa}\epsilon_Q \varepsilon_{\text{cyc}}^\kappa$. Let $f = \mathbf{f}_Q \in e\mathcal{S}_{k_Q}^D(Np^\alpha, \chi\omega^{-\kappa}\epsilon_Q, \mathcal{O}(Q))$. We first claim that the value $\langle \mathbf{U}_p^{-\beta} f, f \rangle_{Np^\beta}$ is independent of any integer $\beta \geq \alpha$. Choose a prime $\ell \nmid Np$ such that $\ell + 1 \not\equiv 0 \pmod{p}$ and ℓ is inert in $\mathbf{Q}(\sqrt{-1})$ and $\mathbf{Q}(\sqrt{-3})$. Then $D^\times \cap x\widehat{R}_{N\ell p^\alpha}^\times x^{-1} = \{\pm 1\}$ for all $x \in \widehat{D}^\times$. It follows that $(1 + \ell) \cdot \langle \mathbf{U}_p^{-\beta} f, f \rangle_{Np^\beta}$ equals

$$\begin{aligned} & \sum_{[x] \in X_0(N\ell p^\beta)} \langle [\tau_{Np^\beta}^D] \mathbf{U}_p^{-\beta} f(x), f(x) \rangle_\kappa \cdot \chi_Q(\nu(x)) \\ &= \sum_{[x] \in X_0(N\ell p^\alpha)} \sum_{b \in \mathbf{Z}_p/p^{\beta-\alpha} \mathbf{Z}_p} \langle [\tau_{Np^\beta}^D] \mathbf{U}_p^{-\beta} f(x \begin{pmatrix} 1 & 0 \\ p^\alpha b & 1 \end{pmatrix}), \rho_{\kappa, p}(\begin{pmatrix} 1 & 0 \\ -p^\alpha b & 1 \end{pmatrix}) f_Q(x) \rangle_\kappa \cdot \chi_Q(\nu(x)) \\ &= \sum_{[x] \in X_0(N\ell p^\alpha)} \sum_{b \in \mathbf{Z}_p/p^{\beta-\alpha} \mathbf{Z}_p} \langle \rho_{\kappa, p}(\begin{pmatrix} 0 & 1 \\ -p^\beta & bp^\alpha \end{pmatrix}) \mathbf{U}_p^{-\beta} f(x\tau_N^D \begin{pmatrix} 0 & 1 \\ -p^\beta & bp^\alpha \end{pmatrix}), f(x) \rangle_\kappa \cdot \chi_Q(\nu(x)) \\ &= \sum_{[x] \in D^\times \backslash \widehat{D}^\times / \widehat{R}_{N\ell p^\alpha}^\times} \sum_{b \in \mathbf{Z}_p/p^{\beta-\alpha} \mathbf{Z}_p} \langle \rho_{\kappa, p}(\tau_{p^\alpha}^D \begin{pmatrix} 1 & -p^{\beta-\alpha} b \\ 0 & 1 \end{pmatrix}) \mathbf{U}_p^{-\beta} f(x\tau_{Np^\alpha}^D \begin{pmatrix} 1 & -p^{\beta-\alpha} b \\ 0 & 1 \end{pmatrix}), f(x) \rangle \cdot \chi_Q(\nu(x)) \\ &= \sum_{[x] \in X_0(N\ell p^\alpha)} \langle [\tau_{Np^\alpha}^D] \mathbf{U}_p^{-\alpha} f(x), f(x) \rangle_\kappa \chi_Q(\nu(x)) = (1 + \ell) \cdot \langle \mathbf{U}_p^{-\alpha} f, f \rangle_{Np^\alpha}. \end{aligned}$$

This verifies the claim. We let $f^{[0]}(x) = \langle f(x), X^\kappa \rangle_\kappa$, $x \in \widehat{D}^\times$ be the specialization of $\mathbf{f}(x)$ at Q . For any positive integer m , there exists a sufficiently larger $\beta > m + v_p(\kappa!)$ such that

$$\begin{aligned} \mathbf{B}_N(\mathbf{f}, \mathbf{f})(Q) \pmod{p^m} &\equiv (1 + \ell)^{-1} \sum_{[x] \in X_0(N\ell p^\beta)} \mathbf{U}_p^{-\beta} f^{[0]}(x \tau_{Np^\beta}^D) f^{[0]}(x) \chi_Q(\nu(x)) \pmod{p^m} \\ &\equiv (1 + \ell)^{-1} \sum_{[x] \in X_0(N\ell p^\beta)} \langle \mathbf{U}_p^{-\beta} f_Q(x \tau_{Np^\beta}^D), \rho_{\kappa, p} \left(\begin{pmatrix} 0 & -1 \\ p^\beta & 0 \end{pmatrix} \right) f_Q(x) \rangle \cdot \chi_Q(\nu(x)) \pmod{p^m} \\ &= \langle \mathbf{U}_p^{-\beta} f, f \rangle_{Np^\beta} \pmod{p^m} = \langle \mathbf{U}_p^{-\alpha} f, f \rangle_{Np^\alpha} \pmod{p^m}. \end{aligned}$$

This proves the lemma. \square

4.5. Hecke algebras and primitive Λ -adic forms. Let $\mathbf{T}^D(N, \mathbf{I})$ be the sub-algebra of $\text{End}_{\mathbf{I}}(e\mathbf{S}^D(N, \mathbf{I}))$ generated by T_q , \mathbf{U}_q and the diamond operators $\langle d \rangle$ over \mathbf{I} and let $\mathbf{T}^D(N, \chi, \mathbf{I})$ be the holomorphic image of $\mathbf{T}^D(N, \mathbf{I})$ in $\text{End}_{\Lambda}(e\mathbf{S}^D(N, \chi, \mathbf{I}))$. Thanks to the Jacquet-Langlands correspondence, there is a surjective \mathbf{I} -algebra homomorphism $JL: \mathbf{T}(N, \mathbf{I}) \rightarrow \mathbf{T}^D(N, \mathbf{I})$ such that $JL(T_q) = T_q$ for $q \nmid Np$, $JL(\mathbf{U}_q) = \mathbf{U}_q$ for $q \mid N^+$, $JL(\mathbf{U}_q) = (-1)\mathbf{U}_q$ for $q \mid N^-$ and $JL(\sigma_d) = \sigma_d$; moreover, for an ordinary Λ -adic newform $\mathbf{f} \in e\mathbf{S}(N, \chi, \mathbf{I})$ of tame conductor N with $\text{supp } N^- \subset \Sigma_{\mathbf{f}}^0$, the corresponding homomorphism $\lambda_{\mathbf{f}}: \mathbf{T}(N, \mathbf{I}) \rightarrow \mathbf{I}$ factors through JL . We denote by $\lambda_{\mathbf{f}}^D: \mathbf{T}^D(N, \mathbf{I}) \rightarrow \mathbf{I}$ the morphism such that $\lambda_{\mathbf{f}} = \lambda_{\mathbf{f}}^D \circ JL$. Put

$$e\mathbf{S}^D(N, \mathbf{I})[\lambda_{\mathbf{f}}^D] := \{ \mathbf{f} \in e\mathbf{S}^D(N, \mathbf{I}) \mid t \cdot \mathbf{f} = \lambda_{\mathbf{f}}^D(t) \mathbf{f} \text{ for } t \in \mathbf{T}^D(N, \mathbf{I}) \}.$$

The multiplicity one theorem for $\text{GL}(2)$ implies that $\dim_{\text{Frac } \Lambda} e\mathbf{S}^D(N, \mathbf{I})[\lambda_{\mathbf{f}}^D] \otimes_{\Lambda} \text{Frac } \Lambda = 1$, but we do not have a notion of normalized eigenforms for quaternionic modular forms due to the lack of the q -expansion. Nonetheless, we have the following

Theorem 4.5. *If \mathbf{f} satisfies the Hypothesis (CR, $\text{supp}(N^-)$) in §1.4, then the \mathbf{I} -module $e\mathbf{S}^D(N, \mathbf{I})[\lambda_{\mathbf{f}}^D]$ is free of rank one. In this case, a generator \mathbf{f}^D of $e\mathbf{S}^D(N, \mathbf{I})[\lambda_{\mathbf{f}}^D]$ is called the primitive Jacquet-Langlands lift of \mathbf{f} . By definition, \mathbf{f}^D is unique up to a scalar in \mathbf{I}^\times .*

PROOF. Let \mathfrak{m} be the maximal ideal of $\mathbf{T}^D(N, \mathbf{I})$ containing $\text{Ker } \lambda_{\mathbf{f}}^D$. Under the Hypothesis (CR), we note that $e\mathbf{S}^D(N, \mathbf{I})_{\mathfrak{m}}$ is a free $\mathbf{T}^D(N, \mathbf{I})_{\mathfrak{m}}$ -module of rank one in virtue of [Wil95, Theorem, 2.1] and [Hel07, Corollary 8.11 and Remark 8.12] and Hida's control theorem (cf. [PW11, Proposition 6.4 and 6.5]). By Theorem 4.2 (1), we find that $e\mathbf{S}^D(N, \chi, \mathbf{I})_{\mathfrak{m}}$ is also a free $\mathbf{T}^D(N, \chi, \mathbf{I})_{\mathfrak{m}}$ -module of rank one which in turn implies that $\mathbf{T}^D(N, \chi, \mathbf{I})_{\mathfrak{m}}$ is Gorenstein as $e\mathbf{S}^D(N, \chi, \mathbf{I})_{\mathfrak{m}}$ is equipped with a Hecke-equivariant perfect pairing \mathbf{B}_N . It follows that $e\mathbf{S}^D(N, \mathbf{I})_{\mathfrak{m}}[\lambda_{\mathbf{f}}^D] = e\mathbf{S}^D(N, \chi, \mathbf{I})_{\mathfrak{m}}[\lambda_{\mathbf{f}}^D] \simeq \mathbf{T}^D(N, \chi, \mathbf{I})_{\mathfrak{m}}[\lambda_{\mathbf{f}}^D]$ is a free of rank one \mathbf{I} -module. \square

4.6. Regularized diagonal cycles and theta elements. Recall that $E = \mathbf{Q} \oplus \mathbf{Q} \oplus \mathbf{Q}$ is the totally split étale cubic algebra over \mathbf{Q} . Let $D_E = D \oplus D \oplus D$. For each positive integer n , let

$$U_{E,1}(Np^n) := U_1(Np^n) \times U_1(Np^n) \times U_1(Np^n)$$

be an open-compact subgroup of \widehat{D}_E^\times . Define the finite set

$$\begin{aligned} \mathbf{X}_n &:= D_E^\times \backslash \widehat{D}_E^\times / U_{E,1}(Np^n) \widehat{\mathbf{Q}}^\times \\ &= (X_n \times X_n \times X_n) / \widehat{\mathbf{Q}}^\times. \end{aligned}$$

The set \mathbf{X}_n is a zero dimensional analogue of the triple product of modular curves. Consider the finitely generated \mathbf{Z}_p -module $\mathbf{Z}_p[\mathbf{X}_n]$ equipped with the operator $\mathbf{U}_{E,p} := \mathbf{U}_p \otimes \mathbf{U}_p \otimes \mathbf{U}_p$ and the ordinary projector $e_E := e \otimes e \otimes e$. For each $(x_1, x_2, x_3) \in \widehat{D}_E^\times$, let $[(x_1, x_2, x_3)]$ denote the double coset $D_E^\times(x_1, x_2, x_3)U_{E,1}(Np^n)\widehat{\mathbf{Q}}^\times$. Set $\tau_{p^n} := \begin{pmatrix} 0 & 1 \\ -p^n & 0 \end{pmatrix} \in \text{GL}_2(\mathbf{Q}_p)$.

Definition 4.6 (Regularized diagonal cycles). Let $\Delta_n \in \mathbf{Z}_p[\mathbf{X}_n]$ be the twisted diagonal cycle given by

$$\Delta_n := \sum_{[x] \in X_0(Np^n)} \sum_{\substack{b \in \mathbf{Z}_p/p^n \mathbf{Z}_p, \\ z \in (\mathbf{Z}_p/p^n \mathbf{Z}_p)^\times}} \left[\left(x \begin{pmatrix} p^n & b \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^n & b+z \\ 0 & 1 \end{pmatrix}, x \tau_{p^n} \begin{pmatrix} 1 & 0 \\ 0 & z \end{pmatrix} \right) \right]$$

and define the regularized diagonal cycle Δ_n^\dagger by

$$\Delta_n^\dagger := \mathbf{U}_{E,p}^{-n}(e_E \Delta_n).$$

The following lemma allows us to define the Λ -adic diagonal cycle

$$\Delta_\infty^\dagger := \varprojlim_{n \rightarrow \infty} \Delta_n^\dagger \in \varprojlim_{n \rightarrow \infty} \mathbf{Z}_p[\mathbf{X}_n],$$

where the inverse limit is taken with respect to the natural homomorphism $N_{n+1,n} : \mathbf{Z}_p[\mathbf{X}_{n+1}] \rightarrow \mathbf{Z}_p[\mathbf{X}_n]$.

Lemma 4.7 (Distribution property). *For every $n \geq 1$,*

$$N_{n+1,n}(\Delta_{n+1}^\dagger) = \Delta_n^\dagger.$$

PROOF. By definition, it is equivalent to showing that

$$N_{n+1,n}(\Delta_{n+1}) = \mathbf{U}_{E,p} \Delta_n.$$

Let $S_n := (\mathbf{Z}_p/p^n \mathbf{Z}_p) \times (\mathbf{Z}_p/p^n \mathbf{Z}_p)^\times$. Then we have

$$\begin{aligned} & N_{n+1,n}(\Delta_{n+1}) \\ &= \sum_{[x] \in X_0(Np^{n+1})} \sum_{(b,z) \in S_n} \sum_{b_1, z_1 \in \mathbf{Z}_p/p\mathbf{Z}_p} [(x \begin{pmatrix} p^n & b \\ 0 & 1 \end{pmatrix} \begin{pmatrix} p & b_1 \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^n & b+z \\ 0 & 1 \end{pmatrix} \begin{pmatrix} p & b_1 + zz_1 \\ 0 & 1 \end{pmatrix}, x\tau_{p^{n+1}} \begin{pmatrix} 1 & 0 \\ 0 & z \end{pmatrix})] \\ &= \sum_{[x] \in X_0(Np^{n+1})} \sum_{(b,z) \in S_n} (\mathbf{U}_p \otimes \mathbf{U}_p \otimes \text{Id}) [(x \begin{pmatrix} p^n & b \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^n & b+z \\ 0 & 1 \end{pmatrix}, x\tau_{p^{n+1}} \begin{pmatrix} 1 & 0 \\ 0 & z \end{pmatrix})] \\ &= \sum_{[x] \in X_0(Np^n)} \sum_{(b,z) \in S_n} \sum_{c \in \mathbf{Z}_p/p\mathbf{Z}_p} (\mathbf{U}_p \otimes \mathbf{U}_p \otimes \text{Id}) [(x \begin{pmatrix} p^n & b \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^n & b+z \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} 1 & 0 \\ p^n c & 1 \end{pmatrix} \tau_{p^{n+1}} \begin{pmatrix} 1 & 0 \\ 0 & z \end{pmatrix})] \\ &= \sum_{[x] \in X_0(Np^n)} \sum_{(b,z) \in S_n} \sum_{c \in \mathbf{Z}_p/p\mathbf{Z}_p} (\mathbf{U}_p \otimes \mathbf{U}_p \otimes \text{Id}) [(x \begin{pmatrix} p^n & b \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^n & b+z \\ 0 & 1 \end{pmatrix}, x\tau_{p^n} \begin{pmatrix} p & -c \\ 0 & 1 \end{pmatrix} \tau_{p^{n+1}} \begin{pmatrix} 1 & 0 \\ 0 & z \end{pmatrix})] \\ &= (\mathbf{U}_p \otimes \mathbf{U}_p \otimes \mathbf{U}_p) \Delta_n. \end{aligned}$$

This proves the assertion. \square

Following the notation in §3.6, we let $\mathcal{R} = \mathbf{I}_1 \widehat{\otimes}_{\mathcal{O}} \mathbf{I}_2 \widehat{\otimes}_{\mathcal{O}} \mathbf{I}_3$ be a finite extension of $\mathcal{R}_0 = \mathcal{O}[[T_1, T_2, T_3]]$. For a triple of ordinary Λ -adic quaternionic forms

$$(\mathbf{f}, \mathbf{g}, \mathbf{h}) \in e\mathbf{S}^D(N, \psi_1, \mathbf{I}_1) \times e\mathbf{S}^D(N, \psi_2, \mathbf{I}_2) \times e\mathbf{S}^D(N, \psi_3, \mathbf{I}_3),$$

we let $\mathbf{F} = \mathbf{f} \boxtimes \mathbf{g} \boxtimes \mathbf{h} : D_E^\times \backslash \widehat{D}_E^\times \rightarrow \mathcal{R}$ be the triple product given by

$$\mathbf{F}(x_1, x_2, x_3) = \mathbf{f}(x_1) \otimes \mathbf{g}(x_2) \otimes \mathbf{h}(x_3).$$

Let $\chi_{\mathcal{R}}^* : \mathbf{Q}^\times \backslash \widehat{\mathbf{Q}}^\times \rightarrow \mathcal{R}^\times$ be the reciprocal of a square root of the character $(\psi_1)_{\mathbf{I}_1} \otimes (\psi_2)_{\mathbf{I}_2} \otimes (\psi_3)_{\mathbf{I}_3}$ defined by

$$\chi_{\mathcal{R}}^*(z) = \omega^a(z) \langle \varepsilon_{\text{cyc}}(z) \rangle^{-3} \langle \varepsilon_{\text{cyc}}(z) \rangle_{\mathbf{I}_1}^{1/2} \langle \varepsilon_{\text{cyc}}(z) \rangle_{\mathbf{I}_2}^{1/2} \langle \varepsilon_{\text{cyc}}(z) \rangle_{\mathbf{I}_3}^{1/2} \in \mathcal{R}^\times$$

and set

$$\mathbf{F}^*(x_1, x_2, x_3) := \mathbf{F}(x_1, x_2, x_3) \cdot \chi_{\mathcal{R}}^*(\nu(x_3)).$$

Then \mathbf{F}^* naturally induces a $\mathbf{Z}_p[[T_1, T_2, T_3]]$ -linear map

$$\mathbf{F}^* : \varprojlim_{n \rightarrow \infty} \mathbf{Z}_p[\mathbf{X}_n] \rightarrow \mathcal{R}.$$

The theta element $\Theta_{\mathbf{F}}$ attached to the triple product \mathbf{F} is then defined by the evaluation of \mathbf{F}^* at the Λ -adic diagonal cycle. In other words,

$$\Theta_{\mathbf{F}} := \mathbf{F}^*(\Delta_\infty^\dagger) \in \mathcal{R}.$$

4.7. **The construction of p -adic L -functions in the balanced case.** We let $\mathbf{F} = (\mathbf{f}, \mathbf{g}, \mathbf{h})$ be the triple of primitive Hida families of tame conductor (N_1, N_2, N_3) in §3.5. Recall that Σ^- is the finite subset of prime factors of $N = \text{lcm}(N_1, N_2, N_3)$ in Definition 3.9. Let $N^- = \prod_{\ell \in \Sigma^-} \ell$. In the remainder of this section, we assume that

- $\#(\Sigma^-)$ is odd,
- \mathbf{f}, \mathbf{g} and \mathbf{h} satisfy the Hypothesis (CR, Σ^-);
- N^- and N/N^- are relatively prime.

Let D be the definite quaternion algebra over \mathbf{Q} with the discriminant N^- and let

$$(\mathbf{f}^D, \mathbf{g}^D, \mathbf{h}^D) \in e\mathbf{S}^D(N_1, \psi_1, \mathbf{I}_1) \times e\mathbf{S}^D(N_2, \psi_2, \mathbf{I}_2) \times e\mathbf{S}^D(N_3, \psi_3, \mathbf{I}_3)$$

be the primitive Jacquet-Langlands lift of $(\mathbf{f}, \mathbf{g}, \mathbf{h})$ constructed in Theorem 4.5.

Definition 4.8. Let $N_i^+ = N_i/N^-$ for $i = 1, 2, 3$ and let $N^+ = \text{lcm}(N_1^+, N_2^+, N_3^+)$. Then $N = N^+N^-$. Define

$$(\mathbf{f}^{D^*}, \mathbf{g}^{D^*}, \mathbf{h}^{D^*}) \in e\mathbf{S}^D(N, \psi_1, \mathbf{I}_1) \times e\mathbf{S}^D(N, \psi_2, \mathbf{I}_2) \times e\mathbf{S}^D(N, \psi_3, \mathbf{I}_3)$$

by

$$\begin{aligned} \mathbf{f}^{D^*} &:= \sum_{I \subset \Sigma_{\mathbf{f},0}^{(\text{Ib})}} (-1)^{|I|} \beta_I(\mathbf{f})^{-1} \cdot V_{\mathbf{d}_{\mathbf{f}/n_{\mathbf{f}}}} \mathbf{f}^D, \\ \mathbf{g}^{D^*} &:= \sum_{I \subset \Sigma_{\mathbf{g},0}^{(\text{Ib})}} (-1)^{|I|} \beta_I(\mathbf{g})^{-1} \cdot V_{\mathbf{d}_{\mathbf{g}/n_{\mathbf{g}}}} \mathbf{g}^D, \\ \mathbf{h}^{D^*} &:= \sum_{I \subset \Sigma_{\mathbf{h},0}^{(\text{Ib})}} (-1)^{|I|} \beta_I(\mathbf{h})^{-1} \cdot V_{\mathbf{d}_{\mathbf{h}/n_{\mathbf{h}}}} \mathbf{h}^D. \end{aligned}$$

Define the triple product $\mathbf{F}^{D^*} : D_E^\times \backslash \widehat{D}_E^\times \rightarrow \mathcal{R}$ by

$$\mathbf{F}^{D^*} := \mathbf{f}^{D^*} \boxtimes \mathbf{g}^{D^*} \boxtimes \mathbf{h}^{D^*}.$$

Then \mathbf{F}^{D^*} is an eigenfunction of the operator $\mathbf{U}_{E,p}$ with the eigenvalue $\alpha_p(\mathbf{F}) := \mathbf{a}(p, \mathbf{f})\mathbf{a}(p, \mathbf{g})\mathbf{a}(p, \mathbf{h})$. We define the associated theta element $\Theta_{\mathbf{F}^{D^*}}$ to be the p -adic L -functions attached to the triple $(\mathbf{f}, \mathbf{g}, \mathbf{h})$ in the balanced range.

4.8. Global trilinear period integrals.

4.8.1. *The setting.* In this subsection, we relate the evaluations of the p -adic L -function $\Theta_{\mathbf{F}^{D^*}}$ at arithmetic points in the balanced range to certain global trilinear period integral on $D_{\mathbf{A}}^\times$. The set $\mathfrak{X}_{\mathcal{R}}^{\text{bal}}$ of arithmetic points in the balanced range, consisting of arithmetic points $\underline{Q} = (Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathbf{I}_1}^+ \times \mathfrak{X}_{\mathbf{I}_2}^+ \times \mathfrak{X}_{\mathbf{I}_3}^+$ such that

$$k_{Q_1} + k_{Q_2} + k_{Q_3} \equiv 0 \pmod{2}; \quad k_{Q_1} + k_{Q_2} + k_{Q_3} > 2k_{Q_i} \text{ for all } i = 1, 2, 3.$$

Let $\underline{Q} = (Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathcal{R}}^{\text{bal}}$. Put

$$k_i = k_{Q_i} \text{ and } \kappa_i = k_i - 2 \text{ for } i = 1, 2, 3.$$

We keep the notation in §3.8. Thus $F = (f, g, h)$ denotes the specialization $\mathbf{F}_{\underline{Q}} = (\mathbf{f}_{Q_1}, \mathbf{g}_{Q_2}, \mathbf{h}_{Q_3})$ of \mathbf{F} at \underline{Q} and $\omega_F^{1/2}$ is the square root of the central character $\omega_F = \omega_f \omega_g \omega_h$ defined in (3.13). Let $\Pi = \Pi_{\underline{Q}}$ be the automorphic representation of $\text{GL}_2(\mathbf{A}_E)$ defined by

$$\Pi_{\underline{Q}} = \pi_f \otimes \omega_F^{-1/2} \boxtimes \pi_g \boxtimes \pi_h.$$

Let $(f^D, g^D, h^D) = (\mathbf{f}_{Q_1}^D, \mathbf{g}_{Q_2}^D, \mathbf{h}_{Q_3}^D)$ be the specializations in the sense of Theorem 4.2 (2). We have

$$(f^D, g^D, h^D) \in \mathcal{S}_{\kappa_1+2}^D(N_1 p^n, \omega_f^{-1}, \mathcal{O}(\underline{Q})) \times \mathcal{S}_{\kappa_2+2}^D(N_2 p^n, \omega_g^{-1}, \mathcal{O}(\underline{Q})) \times \mathcal{S}_{\kappa_3+2}^D(N_3 p^n, \omega_h^{-1}, \mathcal{O}(\underline{Q})),$$

where

$$\omega_f = \psi_1^{-1} \omega^{\kappa_1} \epsilon_{Q_1}^{-1}, \quad \omega_g = \psi_2^{-1} \omega^{\kappa_2} \epsilon_{Q_2}^{-1} \text{ and } \omega_h = \psi_3^{-1} \omega^{\kappa_3} \epsilon_{Q_3}^{-1}.$$

Let $\varphi_{f^D} = \Phi(f^D)$, $\varphi_{g^D} = \Phi(g^D)$ and $\varphi_{h^D} = \Phi(h^D)$ be the associated vector-valued automorphic forms on $D^\times(\mathbf{A})$ as in (4.4). We have

$$(\varphi_{f^D}, \varphi_{g^D}, \varphi_{h^D}) \in \mathcal{A}_{\kappa_1+2}^D(N_1 p^n, \omega_f) \times \mathcal{A}_{\kappa_2+2}^D(N_2 p^n, \omega_g) \times \mathcal{A}_{\kappa_3+2}^D(N_3 p^n, \omega_h).$$

Let $\mathcal{Q}_{1,\ell}(X)$, $\mathcal{Q}_{2,\ell}(X)$ and $\mathcal{Q}_{3,\ell}(X)$ be the polynomials defined in (3.20) and put

$$(4.7) \quad \varphi_1^{D^*} = \prod_{\ell} \mathcal{Q}_{1,\ell}(V_{\ell})(\varphi_{f^D} \otimes \omega_F^{-1/2}), \quad \varphi_2^{D^*} = \prod_{\ell} \mathcal{Q}_{2,\ell}(V_{\ell})\varphi_{g^D}; \quad \varphi_3^{D^*} = \prod_{\ell} \mathcal{Q}_{3,\ell}(V_{\ell})\varphi_{h^D}.$$

Note that

$$(4.8) \quad \begin{aligned} \varphi_1^{D^*} &= \mathbf{d}_f^{\kappa_1/2} \omega_F^{1/2}(\widehat{\mathbf{d}}_f) \cdot \Phi(\mathbf{f}_{Q_1}^{D^*}) \otimes \omega_F^{-1/2}, \\ \varphi_2^{D^*} &= \mathbf{d}_g^{\kappa_2/2} \cdot \Phi(\mathbf{g}_{Q_2}^{D^*}); \quad \varphi_3^{D^*} = \mathbf{d}_h^{\kappa_3/2} \cdot \Phi(\mathbf{h}_{Q_3}^{D^*}). \end{aligned}$$

Let $L_{\underline{\kappa}}(A) := L_{\kappa_1}(A) \otimes L_{\kappa_2}(A) \otimes L_{\kappa_3}(A)$ for any commutative ring A and $\rho_{\underline{\kappa}} = \rho_{\kappa_1} \otimes \rho_{\kappa_2} \otimes \rho_{\kappa_3}$. Define $\rho_{\underline{\kappa}}^{\mathbf{u}}$ and $\rho_{\underline{\kappa},p}$ likewise. For any \mathbf{Q} -algebra R , let $D_E^{\times}(R) := D^{\times}(R) \times D^{\times}(R) \times D^{\times}(R)$ and let $\nu_E^{\underline{\kappa}} : D_E^{\times}(R) \rightarrow R^{\times}$ be the map $\nu_E^{\underline{\kappa}}(x_1, x_2, x_3) := \prod_{i=1}^3 \nu(x_i)^{\kappa_i}$. Define the vector-valued automorphic form

$$\begin{aligned} \vec{\phi}^{D^*} : D_E^{\times}(\mathbf{A}) &\rightarrow L_{\underline{\kappa}}(\mathbf{C}), \\ \vec{\phi}^{D^*}(x_1, x_2, x_3) &= \varphi_1^{D^*}(x_1) \otimes \varphi_2^{D^*}(x_2) \otimes \varphi_3^{D^*}(x_3) \quad (x_i \in D^{\times}(\mathbf{A})). \end{aligned}$$

Define $\mathbf{P}_{\underline{\kappa}} \in L_{\underline{\kappa}}(\mathbf{Z})$ by

$$(4.9) \quad \begin{aligned} \mathbf{P}_{\underline{\kappa}} &= (X_1 Y_2 - X_2 Y_1)^{\kappa_1^*} (X_3 Y_1 - X_1 Y_3)^{\kappa_2^*} (X_3 Y_2 - X_2 Y_3)^{\kappa_3^*}, \\ \kappa_i^* &:= \frac{\kappa_1 + \kappa_2 + \kappa_3}{2} - \kappa_i \quad (i = 1, 2, 3). \end{aligned}$$

Then $\mathbf{P}_{\underline{\kappa}}$ is a basis of the line $L_{\underline{\kappa}}(\mathbf{C})$ fixed by D_{∞}^{\times} under the action of $\rho_{\underline{\kappa}}^{\mathbf{u}}$. Define the automorphic form

$$(4.10) \quad \begin{aligned} \phi_F^{D^*} : D_E^{\times}(\mathbf{A}) &\rightarrow \mathbf{C}, \\ \phi_F^{D^*}(x_1, x_2, x_3) &= \langle \vec{\phi}^{D^*}(x_1, x_2, x_3), \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}}, \end{aligned}$$

where $\langle \cdot, \cdot \rangle_{\underline{\kappa}} = \langle \cdot, \cdot \rangle_{\kappa_1} \otimes \langle \cdot, \cdot \rangle_{\kappa_2} \otimes \langle \cdot, \cdot \rangle_{\kappa_3}$. One verifies that

$$(4.11) \quad \phi_F^{D^*}(x_1 u_{\infty}, x_2 u_{\infty}, x_3 u_{\infty}) = \phi_F^{D^*}(x_1, x_2, x_3) \text{ for } u_{\infty} \in D_{\infty}^{\times}.$$

4.8.2. *The global trilinear period integrals.* Let $n_{\underline{Q}} = \max\{c(\epsilon_{Q_1}), c(\epsilon_{Q_2}), c(\epsilon_{Q_3}), 1\}$ and let $n \geq n_{\underline{Q}}$ be a positive integer. Let $\check{\mathbf{t}}_n \in D_E^{\times}(\mathbf{Q}_p)$ be the matrix given by

$$\check{\mathbf{t}}_n = \left(\begin{pmatrix} 1 & p^{-n} \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & p^{-n} \\ -p^n & 0 \end{pmatrix} \right) \in \mathrm{GL}_2(E_p).$$

We shall relate the interpolation to the global trilinear period integral

$$I(\rho(\check{\mathbf{t}}_n)\phi_F^{D^*}) = \int_{D^{\times}\mathbf{A}^{\times}\backslash D_{\mathbf{A}}^{\times}} \phi_F^{D^*}\left(x \begin{pmatrix} 1 & p^{-n} \\ 0 & 1 \end{pmatrix}, x, x \begin{pmatrix} 0 & p^{-n} \\ -p^n & 0 \end{pmatrix}\right) d^{\tau}x.$$

Here $d^{\tau}x$ is the Tamagawa measure on $\mathbf{A}^{\times}\backslash D_{\mathbf{A}}^{\times}$.

Proposition 4.9. *For every $n \geq n_{\underline{Q}}$, we have*

$$\Theta_{\mathbf{F}^{D^*}}(\underline{Q}) = \frac{1}{\mathrm{vol}(\widehat{R}_N^{\times})} \cdot I(\rho(\check{\mathbf{t}}_n)\phi_F^{D^*}) \cdot \frac{\omega_{F,p}^{1/2}(p^n) |p^n|^{-\frac{\kappa_1 + \kappa_2 + \kappa_3}{2}}}{\alpha_p(F)^n \zeta_p(2)} \cdot \frac{1}{\omega_F^{1/2}(\widehat{\mathbf{d}}_f) \mathbf{d}_F^{\underline{\kappa}/2}},$$

where $\alpha_p(F) = \mathbf{a}(p, f)\mathbf{a}(p, g)\mathbf{a}(p, h)$ and $\mathbf{d}_F^{\underline{\kappa}/2} = \mathbf{d}_f^{\kappa_1/2} \mathbf{d}_g^{\kappa_2/2} \mathbf{d}_h^{\kappa_3/2}$ defined in (3.15).

PROOF. We begin with some notation. Let $\underline{Q}(\mathbf{F}^{D^*}) : D_E^{\times}\backslash \widehat{D}_E^{\times} \rightarrow \mathcal{O}_{\mathbf{C}_p}$ denote the value of \mathbf{F}^{D^*} at the point $\underline{Q} \in \mathrm{Spec} \mathcal{R}(\overline{\mathbf{Q}}_p)$. Namely,

$$\underline{Q}(\mathbf{F}^{D^*})(x_1, x_2, x_3) = Q_1(\mathbf{f}^{D^*}(x_1))Q_2(\mathbf{g}^{D^*}(x_2))Q_3(\mathbf{h}^{D^*}(x_3)).$$

Let $(f^{D^*}, g^{D^*}, h^{D^*}) = (\mathbf{f}_{Q_1}^{D^*}, \mathbf{g}_{Q_2}^{D^*}, \mathbf{h}_{Q_3}^{D^*})$ denote the specialization of $(\mathbf{f}^{D^*}, \mathbf{g}^{D^*}, \mathbf{h}^{D^*})$ as in Theorem 4.2 (2). Put

$$\widehat{F}^{D^*} := f^{D^*} \boxtimes g^{D^*} \boxtimes h^{D^*}, \quad \widehat{F}^{D^*}(x_1, x_2, x_3) = f^{D^*}(x_1) \otimes g^{D^*}(x_2) \otimes h^{D^*}(x_3).$$

By definition, we have

$$(4.12) \quad \underline{Q}(\mathbf{F}^{D^\star})(x_1, x_2, x_3) = \langle \widehat{F}^{D^\star}(x_1, x_2, x_3), X_1^{\kappa_1} X_2^{\kappa_2} X_3^{\kappa_3} \rangle_{\underline{\kappa}}.$$

Define $F^{D^\star} : D_E^\times(\mathbf{A}) \rightarrow L_{\underline{\kappa}}(\mathbf{C})$ to be the function

$$F^{D^\star}(x) = \rho_{\underline{\kappa}, p}(x_p) \widehat{F}^\star(x) \quad (x \in D_E^\times(\mathbf{A})).$$

Then one verifies that

$$\phi_F^{D^\star}(x) = d_F^{\kappa/2} \cdot \langle F^{D^\star}(x), \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}} \big| \nu_E^\kappa(x) \big|_{\mathbf{A}}^{1/2}.$$

Let m_k be the p -adic valuation of $(\kappa_1 + \kappa_2 + \kappa_3)!$ and let $m > m_k$ be a positive integer. For a number $A \in \mathbf{C}_p$, denote by $A \pmod{p^m}$ the residue class of A in \mathbf{C}_p modulo $p^m \mathcal{O}_{\mathbf{C}_p}$. By definition, for any sufficiently large integer $s \gg n + m + m_k \geq 1$,

$$(4.13) \quad \begin{aligned} & \Theta_{\mathbf{F}^{D^\star}}(\underline{Q}) \pmod{p^m} \equiv \underline{Q}((\mathbf{F}^{D^\star})^*)(\Delta_s^\dagger) \pmod{p^m} \\ & \equiv \alpha_p(F)^{-s} \sum_{[x] \in X_0(Np^s)} \sum_{\substack{b \in (\mathbf{Z}_p/p^s \mathbf{Z}_p), \\ z \in (\mathbf{Z}_p/p^s \mathbf{Z}_p)^\times}} \underline{Q}(\mathbf{F}^{D^\star})\left(x \begin{pmatrix} p^s & b \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^s & b+z \\ 0 & 1 \end{pmatrix}, x\tau_{p^s}\right) \\ & \quad \times \mathbb{k}_h(z) z^{\kappa_3^*} \cdot \chi_{\underline{Q}}^*(\nu(x)) \pmod{p^m}, \end{aligned}$$

where $\mathbb{k}_h(z) := \omega_F^{-1/2} \omega_h(z)$ for $z \in \mathbf{Z}_p^\times$ and $\chi_{\underline{Q}}^*$ is the specialization of $\chi_{\mathcal{R}}^*$ at \underline{Q}

$$\chi_{\underline{Q}}^* = \omega_F^{-1/2} \cdot \varepsilon_{\text{cyc}}^{r_\kappa} \quad (r_\kappa := \frac{\kappa_1 + \kappa_2 + \kappa_3}{2} = \frac{k_1 + k_2 + k_3}{2} - 3)$$

Putting

$$W'_s = \{(b_1, b_2) \in (\mathbf{Z}_p/p^s \mathbf{Z}_p)^2 \mid b_1 - b_2 \in (\mathbf{Z}_p/p^s \mathbf{Z}_p)^\times\},$$

we see from (4.13) and (4.12) that

$$(4.14) \quad \begin{aligned} & \Theta_{\mathbf{F}^{D^\star}}(\underline{Q}) \pmod{p^m} \equiv \alpha_p(F)^{-s} \sum_{\substack{x \in X_0(Np^s), \\ (b_1, b_2) \in W'_s}} (b_1 - b_2)^{\kappa_3^*} \mathbb{k}_h(b_1 - b_2) \chi_{\underline{Q}}^*(\nu(x)) \\ & \quad \times \underline{Q}(\mathbf{F}^{D^\star})\left(x \begin{pmatrix} p^s & b_1 \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^s & b_2 \\ 0 & 1 \end{pmatrix}, x\tau_{p^s}\right) \pmod{p^m} \\ & \equiv \alpha_p(F)^{-s} \sum_{x \in X_0(Np^s)} \sum_{c \in p^n \mathbf{Z}_p/p^s \mathbf{Z}_p} \sum_{(b_1, b_2) \in W'_s} \mathbb{k}_h(b_1 - b_2) \chi_{\underline{Q}}^*(\nu(x)) \\ & \quad \times \langle \widehat{F}^{D^\star}\left(x \begin{pmatrix} p^s & b_1 \\ cp^s & 1 + b_1c \end{pmatrix}\right), x \begin{pmatrix} p^s & b_2 \\ cp^s & 1 + b_2c \end{pmatrix}, x \begin{pmatrix} 0 & 1 \\ -p^s & c \end{pmatrix}\rangle, (b_1 - b_2)^{\kappa_3^*} X_1^{\kappa_1} X_2^{\kappa_2} X_3^{\kappa_3} \rangle_{\underline{\kappa}} \pmod{p^m}. \end{aligned}$$

To simplify the above expression, we note that by (4.3),

$$\langle \rho_{\underline{\kappa}}(x_p^{-1}) F^{D^\star}(xg_1, xg_2, xg_3), \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}} = \langle \widehat{F}^{D^\star}(xg_1, xg_2, xg_3), \rho_{\underline{\kappa}}(g'_1 \otimes g'_2 \otimes g'_3) \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}}$$

with

$$g_1 = \begin{pmatrix} p^s & b_1 \\ cp^s & 1 + b_1c \end{pmatrix}, g_2 = \begin{pmatrix} p^s & b_2 \\ cp^s & 1 + b_2c \end{pmatrix}, g_3 = \begin{pmatrix} 0 & 1 \\ -p^s & c \end{pmatrix},$$

we find the congruence relation

$$\begin{aligned} & \langle \rho_{\underline{\kappa}}(x_p^{-1}) F^{D^\star}(xg_1, xg_2, xg_3), \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}} \\ & \equiv \langle \widehat{F}^{D^\star}(xg_1, xg_2, xg_3), \rho_{\underline{\kappa}}\left(\begin{pmatrix} 1 + b_1c & -b_1 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 + b_2c & -b_2 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} c & -1 \\ 0 & 0 \end{pmatrix}\right) \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}} \pmod{p^m} \\ & \equiv \langle \widehat{F}^{D^\star}(xg_1, xg_2, xg_3), \begin{pmatrix} 1 + cb_1 & -b_1 \\ 1 + b_2c & -b_2 \end{pmatrix}^{\kappa_3^*} \begin{pmatrix} c & -1 \\ 1 + b_1c & -b_1 \end{pmatrix}^{\kappa_2^*} \begin{pmatrix} c & -1 \\ 1 + b_2c & -b_2 \end{pmatrix}^{\kappa_1^*} X_1^{\kappa_1} X_2^{\kappa_2} X_3^{\kappa_3} \rangle_{\underline{\kappa}} \pmod{p^m} \\ & \equiv \langle \widehat{F}^{D^\star}(xg_1, xg_2, xg_3), (b_1 - b_2)^{\kappa_3^*} X_1^{\kappa_1} X_2^{\kappa_2} X_3^{\kappa_3} \rangle_{\underline{\kappa}} \pmod{p^m}. \end{aligned}$$

Substituting the above to (4.14), we see that $\Theta_{\mathbf{F}^{D^*}}(\underline{Q}) \pmod{p^m}$ equals

$$\begin{aligned}
& \alpha_p(F)^{-s} \sum_{x \in X_0(Np^n)} \sum_{c \in p^n \mathbf{Z}_p / p^s \mathbf{Z}_p} \sum_{(b_1, b_2) \in W'_s} \mathbb{k}_h(b_1 - b_2) \chi_{\underline{Q}}^*(\nu(x)) \\
& \times \langle \rho_{\underline{\kappa}}(x_p^{-1}) F^{D^*} \left(x \begin{pmatrix} p^s & b_1 \\ cp^s & 1 + b_1c \end{pmatrix}, x \begin{pmatrix} p^s & b_2 \\ cp^s & 1 + b_2c \end{pmatrix}, x \begin{pmatrix} 0 & 1 \\ -p^s & c \end{pmatrix}, \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}} \pmod{p^m} \\
\equiv & \alpha_p(F)^{-s} \sum_{x \in X_0(Np^n)} \sum_{(b_1, b_2) \in W'_s} \mathbb{k}_h(b_1 - b_2) \chi_{\underline{Q}}^*(\nu(x)) \nu(x_p)^{-r_\kappa} \\
& \times \sum_{c \in p^n \mathbf{Z}_p / p^s \mathbf{Z}_p} \langle F^{D^*} \left(x \begin{pmatrix} p^s & b_1 \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^s & b_2 \\ 0 & 1 \end{pmatrix}, x \tau_{p^n} \begin{pmatrix} p^{s-n} & -p^{-n}c \\ 0 & 1 \end{pmatrix}, \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}} \pmod{p^m} \\
\equiv & \alpha_p(F)^{-n} \sum_{x \in X_0(Np^n)} \sum_{(b_1, b_2) \in W'_n} \mathbb{k}_h(b_1 - b_2) \omega_F^{-1/2} |\cdot|_{\mathbf{A}}^{r_\kappa}(\nu(x)) \cdot \langle F^{D^*} \left(x \begin{pmatrix} p^n & b_1 \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^n & b_2 \\ 0 & 1 \end{pmatrix}, x \tau_{p^n}, \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}} \pmod{p^m}.
\end{aligned}$$

The last congruence relation holds for any sufficiently large m , so we obtain the expression

$$\begin{aligned}
(4.15) \quad \Theta_{\mathbf{F}^{D^*}}(\underline{Q}) &= \alpha_p(F)^{-n} \sum_{x \in X_0(Np^n)} \sum_{\substack{b_1 \in (\mathbf{Z}_p/p^n \mathbf{Z}_p)^\times, \\ b_2 \in \mathbf{Z}_p/p^n \mathbf{Z}_p}} \mathbb{k}_h(b_1) \omega_F^{-1/2} |\cdot|_{\mathbf{A}}^{r_\kappa}(\nu(x)) \\
& \times \langle F^{D^*} \left(x \begin{pmatrix} p^n & b_1 + b_2 \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^n & b_2 \\ 0 & 1 \end{pmatrix}, x \tau_{p^n}, \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}}.
\end{aligned}$$

By the definition (4.8),

$$\overline{\phi}^{D^*}(x_1, x_2, x_3) = \mathbf{d}_F^{\kappa/2} \cdot F^{D^*}(x_1, x_2, x_3) \omega_F^{-1/2}(\nu(x_1)) |\nu_E^\kappa(x_1, x_2, x_3)|_{\mathbf{A}}^{1/2}$$

for $(x_1, x_2, x_3) \in \widehat{D}_E^\times$, and using (4.11), we obtain

$$\begin{aligned}
& \sum_{x \in X_0(Np^n)} \omega_F^{-1/2} |\cdot|_{\mathbf{A}}^{r_\kappa}(\nu(x)) \cdot \langle F^{D^*} \left(x \begin{pmatrix} p^n & b_1 + b_2 \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^n & b_2 \\ 0 & 1 \end{pmatrix}, x \tau_{p^n}, \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}} \\
&= \frac{\omega_{F,p}^{1/2}(p^n) |p^n|^{-r_\kappa}}{\text{vol}(\widehat{R}_{Np^n}^\times) \mathbf{d}_F^{\kappa/2}} \int_{\mathbf{A} \times D^\times \setminus D_{\mathbf{A}}^\times} \phi_F^{D^*} \left(x \begin{pmatrix} p^n & b_1 + b_2 \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^n & b_2 \\ 0 & 1 \end{pmatrix}, x \tau_{p^n} \right) d^T x.
\end{aligned}$$

Since $\mathbb{k}_h = \omega_F^{-1/2} \omega_h$, we find that the right hand side of the equation (4.15) equals

$$\begin{aligned}
& \frac{\omega_{F,p}^{1/2}(p^n) |p^n|^{-r_\kappa}}{\alpha_p(F)^n \text{vol}(\widehat{R}_{Np^n}^\times) \mathbf{d}_F^{\kappa/2}} \int_{\mathbf{A} \times D^\times \setminus D_{\mathbf{A}}^\times} \sum_{\substack{b_1 \in (\mathbf{Z}_p/p^n \mathbf{Z}_p)^\times, \\ b_2 \in \mathbf{Z}_p/p^n \mathbf{Z}_p}} \mathbb{k}_h(b_1) \cdot \phi_F^{D^*} \left(x \begin{pmatrix} p^n & b_1 + b_2 \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^n & b_2 \\ 0 & 1 \end{pmatrix}, x \tau_{p^n} \right) d^T x \\
&= \frac{p^{2n} (1 - p^{-1}) \omega_{F,p}^{1/2}(p^n) |p^n|^{-r_\kappa}}{\alpha_p(F)^n \text{vol}(\widehat{R}_{Np^n}^\times) \mathbf{d}_F^{\kappa/2}} \int_{\mathbf{A} \times D^\times \setminus D_{\mathbf{A}}^\times} \phi_F^{D^*} \left(x \begin{pmatrix} p^n & 1 \\ 0 & 1 \end{pmatrix}, x \begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix}, x \tau_{p^n} \right) d^T x.
\end{aligned}$$

Since $\text{vol}(\widehat{R}_N^\times) = \text{vol}(\widehat{R}_{Np^n}^\times) (1 + p^{-1}) p^n$, the proposition can be deduced from the last equation directly by making change of variable. \square

4.9. Ichino's formula. We now apply Ichino's formula to relate the global trilinear period $I(\rho(\check{\mathbf{t}}_n) \phi_F^{D^*})$ to a product of central L -values of triple L -functions, the local zeta integrals $I_q(\phi_q^* \otimes \check{\phi}_q^*)$ defined in (3.23) at primes $q \neq p$ and the following local zeta integral at p

$$(4.16) \quad I_p^{\text{ord}}(\phi_p \otimes \check{\phi}_p, \check{\mathbf{t}}_n) := \frac{L(1, \Pi_p, \text{Ad})}{\zeta_p(2)^2 L(1/2, \Pi_p)} \int_{\text{PGL}_2(\mathbf{Q}_p)} \frac{\mathbf{b}_p(\Pi_p(g_p \check{\mathbf{t}}_n) \phi_p \otimes \check{\Pi}_p(\check{\mathbf{t}}_n) \check{\phi}_p)}{\mathbf{b}_p(\Pi_p(t_n) \phi_p, \check{\phi}_p)} dg_p.$$

Here we recall that ϕ_p is any non-zero vector in the ordinary line $\mathcal{V}_{\pi_{1,p}}^{\text{ord}}(\chi_{1,p}) \otimes \mathcal{V}_{\pi_{2,p}}^{\text{ord}}(\chi_{2,p}) \otimes \mathcal{V}_{\pi_{3,p}}^{\text{ord}}(\chi_{3,p})$ with characters $\chi_{i,p}$ defined in (3.19) and $t_n = \begin{pmatrix} 0 & p^{-n} \\ -p^n & 0 \end{pmatrix} \in D_p^\times \hookrightarrow D_E^\times(\mathbf{Q}_p)$ for any integer $n \geq n_{\underline{Q}}$. For each

positive integer M , we shall use the notation $\widehat{M} \in \widehat{\mathbf{Q}}^\times$ to denote the idele with $\widehat{M}_\ell = \ell^{v_\ell(M)}$ at each finite prime ℓ .

Proposition 4.10. *We have*

$$\frac{I(\rho(\check{\mathfrak{t}}_n)\phi_F^{D*})^2}{\langle F^D, F^D \rangle} = 2^{\#\Sigma^- + 1} \text{vol}(\widehat{\mathcal{O}}_D^\times)^2 \cdot L(1/2, \Pi) \cdot \frac{\omega_F^{-1/2}(\widehat{N}_1^+) \cdot \omega_{F,p}^{-1}(p^n) \alpha_p(F)^{2n}}{L(1, \Pi, \text{Ad}) \prod_{i=1}^3 [\text{SL}_2(\mathbf{Z}) : \Gamma_0(N_i p^n)] (N_i^+ p^n)^{\kappa_i/2}}$$

$$\times I_p^{\text{ord}}(\phi_p \otimes \tilde{\phi}_p, \check{\mathfrak{t}}_n) \prod_{q \in \Sigma^-} \frac{\zeta_q(1)^3}{\zeta_q(2)^3} \cdot \prod_{q|N^+} I_q(\phi_q^* \otimes \tilde{\phi}_q^*),$$

where

$$\langle F^D, F^D \rangle = \langle \mathbf{U}_p^{-n} f^D, f^D \rangle_{N_1 p^n} \langle \mathbf{U}_p^{-n} g^D, g^D \rangle_{N_2 p^n} \langle \mathbf{U}_p^{-n} h^D, h^D \rangle_{N_3 p^n}.$$

PROOF. We begin with the explanation of the representation theoretic factorization for the automorphic form ϕ_F^{D*} . Let $(\pi_f^D, \pi_g^D, \pi_h^D)$ be the image of (π_f, π_g, π_h) under the Jacquet-Langlands correspondence and let

$$\pi_1^D = \pi_f^D \otimes \omega_F^{-1/2}, \quad \pi_2^D = \pi_g^D \quad \text{and} \quad \pi_3^D = \pi_h^D.$$

Let $\Pi^D = \pi_1^D \boxtimes \pi_2^D \boxtimes \pi_3^D$ be the Jacquet-Langlands transfer of Π and let $\mathcal{A}(\Pi^D)$ be the unique automorphic realization of Π^D . With the isomorphism $\Psi : \widehat{D}^{(N^-)^\times} \simeq \text{GL}_2(\widehat{\mathbf{Q}}^{(N^-)})$, we have a factorization

$$(4.17) \quad \mathcal{A}(\Pi^D) \simeq \bigotimes_{v \in \Sigma_D} \mathcal{V}_{\Pi_v^D} \bigotimes_{v \notin \Sigma_D} \mathcal{V}_{\Pi_v}.$$

Here $(\Pi_\infty^D, \mathcal{V}_{\Pi_\infty^D}) = (\rho_{\underline{\mathbf{k}}}^u, L_{\underline{\mathbf{k}}}(\mathbf{C}))$ and for finite prime $\ell \mid N^-$, $(\Pi_\ell^D, \mathcal{V}_{\Pi_\ell^D}) = (\mu_{E_\ell} \circ \nu, \mathbf{C} e_{\mu_{E_\ell}})$ is the one dimensional representation given by a unramified character $\mu_{E_\ell} = (\mu_{1,\ell}, \mu_{2,\ell}, \mu_{3,\ell}) : E_\ell^\times \rightarrow \mathbf{C}^\times$ with a basis $e_{\mu_{E_\ell}}$. Consider $\vec{\phi}_F^D = \varphi_1^D \boxtimes \varphi_2^D \boxtimes \varphi_3^D \in \mathcal{A}(\Pi^D) \otimes L_{\underline{\mathbf{k}}}(\mathbf{C})$. Let $X^{\underline{\mathbf{k}}} := X_1^{\kappa_1} X_2^{\kappa_2} X_3^{\kappa_3} \in L_{\underline{\mathbf{k}}}(\mathbf{C})$ and define $\phi_{X^{\underline{\mathbf{k}}}}^D \in \mathcal{A}(\Pi^D)$ by

$$\phi_{X^{\underline{\mathbf{k}}}}^D(x) := \langle \vec{\phi}^D(x), X^{\underline{\mathbf{k}}} \rangle_{\underline{\mathbf{k}}} \quad (x \in D_E^\times(\mathbf{A})).$$

Under the isomorphism (4.17), we have the factorization $\phi_{X^{\underline{\mathbf{k}}}}^D = \otimes_v \phi_v^D$, where

$$\begin{aligned} \phi_\infty^D &= X_1^{\kappa_1} X_2^{\kappa_2} X_3^{\kappa_3}, & \phi_\ell^D &= e_{\mu_{E_\ell}} \text{ for } \ell \mid N^-, \\ \phi_\ell^D &= \varphi_{1,\ell} \otimes \varphi_{2,\ell} \otimes \varphi_{3,\ell} \text{ for } \ell \notin \Sigma_D \end{aligned}$$

as in §3.8.1. Recall that $\varphi_{i,\ell} \in \mathcal{V}_{\pi_{i,v}}^{\text{new}}$ for $\ell \neq p$ is a new vector and $\varphi_{i,p} \in \mathcal{V}_{\pi_{i,p}}^{\text{ord}}(\chi_{i,p})$ is an ordinary vector. In view of the definition of ϕ_F^{D*} in (4.10), we obtain the factorization $\phi_F^{D*} = \otimes_v \phi_v^{D*}$, where

$$(4.18) \quad \phi_v^{D*} = \begin{cases} \mathbf{P}_{\underline{\mathbf{k}}} \in L_{\underline{\mathbf{k}}}(\mathbf{C}) & \text{if } v = \infty, \\ e_{\mu_{E_\ell}} & \text{if } v = \ell \in \Sigma^-, \\ \varphi_{1,p} \otimes \varphi_{2,p} \otimes \varphi_{3,p} (= \phi_p) & \text{if } v = p, \\ \mathcal{Q}_{1,\ell}(V_\ell) \varphi_{1,\ell} \otimes \mathcal{Q}_{2,\ell}(V_\ell) \varphi_{2,\ell} \otimes \mathcal{Q}_{3,\ell}(V_\ell) \varphi_{3,\ell} (= \phi_\ell^*) & \text{if } v = \ell \notin \{p\} \cup \Sigma^-. \end{cases}$$

Now consider the contragredient representation $\widetilde{\Pi}^D$. Let $\tilde{\varphi}_i^D = \varphi_i^D \otimes \omega_i^{-1}$ and $\tilde{\varphi}_i^{D*} = \varphi_i^{D*} \otimes \omega_i^{-1}$ for $i = 1, 2, 3$. Let $Y^{\underline{\mathbf{k}}} = Y^{\kappa_1} Y^{\kappa_2} Y^{\kappa_3} \in L_{\underline{\mathbf{k}}}(\mathbf{C})$. Define $\tilde{\phi}_{Y^{\underline{\mathbf{k}}}}^D$ and $\tilde{\phi}_F^{D*} \in \mathcal{A}(\widetilde{\Pi}^D)$ by

$$\tilde{\phi}_{Y^{\underline{\mathbf{k}}}}^D(x) := \langle \tilde{\varphi}_1^D \boxtimes \tilde{\varphi}_2^D \boxtimes \tilde{\varphi}_3^D(x), Y^{\underline{\mathbf{k}}} \rangle_{\underline{\mathbf{k}}}; \quad \tilde{\phi}_F^{D*}(x) = \langle \varphi_1^{D*} \boxtimes \varphi_2^{D*} \boxtimes \varphi_3^{D*}(x), \mathbf{P}_{\underline{\mathbf{k}}} \rangle_{\underline{\mathbf{k}}}$$

for $x \in D_E^\times(\mathbf{A})$. Fixing an isomorphism

$$\mathcal{A}(\widetilde{\Pi}^D) \simeq \bigotimes_{v \in \Sigma_D} \mathcal{V}_{\widetilde{\Pi}_v^D} \bigotimes_{v \notin \Sigma_D} \mathcal{V}_{\widetilde{\Pi}_v},$$

we then have a similar description for the factorizations $\tilde{\phi}_{Y^{\underline{\mathbf{k}}}}^D = \otimes_v \tilde{\phi}_v^D$ and $\tilde{\phi}_F^{D*} = \otimes_v \tilde{\phi}_v^{D*}$ likewise.

For $v \notin \{\infty\} \cup \Sigma^-$, let $\mathbf{b}_v : \mathcal{V}_{\Pi_v} \times \mathcal{V}_{\tilde{\Pi}_v} \rightarrow \mathbf{C}$ be a non-degenerate $\mathrm{GL}_2(E_v)$ -equivariant pairing such that $\mathbf{b}_v(\tilde{\phi}_v^D, \phi_v^D) = 1$ for all but finitely many v . For $v \in \{\infty\} \cup \Sigma^-$, let $\mathbf{b}_v^D : \mathcal{V}_{\Pi_v^D} \times \mathcal{V}_{\tilde{\Pi}_v^D} \rightarrow \mathbf{C}$ be a $D_E^\times(\mathbf{Q}_v)$ -equivariant pairing and define

$$I_v(\phi_v^D \otimes \tilde{\phi}_v^D) = \frac{L(1, \Pi_v, \mathrm{Ad})}{\zeta_v(2)^2 L(1/2, \Pi_v)} \int_{D_v^\times / \mathbf{Q}_v^\times} \frac{\mathbf{b}_v^D(\Pi_v^D(x_v) \phi_v^{D*} \otimes \tilde{\phi}_v^{D*})}{\mathbf{b}_v^D(\phi_v^D, \tilde{\phi}_v^D)} dx_v.$$

Here dx_v is the Haar measure with $\mathrm{vol}(\mathcal{O}_{D_v}^\times / \mathbf{Z}_v^\times, dx_v) = 1$. In the notation of [Ich08, page 282], we have

$$I(\rho(\check{\mathfrak{t}}_n) \phi_F^{D*})^2 = I(\rho(\check{\mathfrak{t}}_n) \phi_F^{D*}) \cdot I(\rho(\check{\mathfrak{t}}_n) \tilde{\phi}_F^{D*}).$$

Therefore, according to [Ich08, Theorem 1.1, Remark 1.3], we obtain

$$\begin{aligned} \frac{I(\rho(\check{\mathfrak{t}}_n) \phi_F^{D*})^2}{\langle \rho(\tau_{\underline{N}^+}^D t_n) \phi_{X^\kappa}^D, \tilde{\phi}_{Y^\kappa}^D \rangle} &= \frac{\mathrm{vol}(\widehat{\mathcal{O}}_D^\times)}{8} \cdot \frac{\zeta_{\mathbf{Q}}(2)^2 L(1/2, \Pi)}{L(1, \Pi, \mathrm{Ad})} \\ &\quad \times I_p^{\mathrm{ord}}(\phi_p \otimes \tilde{\phi}_p, \check{\mathfrak{t}}_n) \prod_{v \in \{\infty\} \cup \Sigma^-} I_v(\phi_v^D \otimes \tilde{\phi}_v^D) \prod_{q \notin \{p\} \cup \Sigma^-} I_q(\phi_q^* \otimes \tilde{\phi}_q^*), \end{aligned}$$

From (4.5) and (4.2), we find that $\langle \rho(\tau_{\underline{N}^+}^D t_n) \phi_{X^\kappa}^D, \tilde{\phi}_{Y^\kappa}^D \rangle$ equals

$$\begin{aligned} (4.19) \quad &\omega_F^{-1/2}(\widehat{N}_1^+) \omega_{F,p}^{-1}(p^n) \cdot \langle F^D, F^D \rangle \cdot \alpha_p(F)^{2n} \prod_{i=1}^3 \frac{\mathrm{vol}(\widehat{R}_{N_i p^{2n}}^\times)}{(N_i^+ p^{2n})^{\kappa_i/2} (\kappa_i + 1)} \\ &= \omega_F^{-1/2}(\widehat{N}_1^+) \langle F^D, F^D \rangle \cdot \frac{48^3 \cdot \omega_{F,p}^{-1}(p^n) \alpha_p(F)^{2n}}{\prod_{i=1}^3 (N_i^+ p^{2n})^{\kappa_i/2} [\mathrm{SL}_2(\mathbf{Z}) : \Gamma_0(N_i)]} \prod_{i=1}^3 \frac{1}{(\kappa_i + 1)} \prod_{q \in \Sigma^-} \frac{\zeta_q(1)^6}{\zeta_q(2)^3} \end{aligned}$$

We now proceed to compute the local zeta integrals $I_v(\phi_v^D \otimes \tilde{\phi}_v^D)$ for $v \in \{\infty\} \cup \Sigma^-$. Recall that the archimedean L -factors are given by

$$\begin{aligned} L(s, \Pi_\infty, \mathrm{Ad}) &= \pi^{-3} \Gamma_{\mathbf{C}}(s + \kappa_1 + 1) \Gamma_{\mathbf{C}}(s + \kappa_2 + 1) \Gamma_{\mathbf{C}}(s + \kappa_3 + 1); \\ L(s, \Pi_\infty) &= \Gamma_{\mathbf{C}}(s + \frac{\kappa_1 + \kappa_2 + \kappa_3 + 3}{2}) \Gamma_{\mathbf{C}}(s + \kappa_1^* + \frac{1}{2}) \Gamma_{\mathbf{C}}(s + \kappa_2^* + \frac{1}{2}) \Gamma_{\mathbf{C}}(s + \kappa_3^* + \frac{1}{2}), \end{aligned}$$

so we have

$$\begin{aligned} I_\infty(\phi_\infty^D \otimes \tilde{\phi}_\infty^D) &= \frac{L(1, \Pi_\infty, \mathrm{Ad})}{\zeta_\infty(2)^2 L(1/2, \Pi_\infty)} \int_{D^\times(\mathbf{R})/\mathbf{R}^\times} \frac{\langle \rho_{\underline{\kappa}}^u(x_\infty) \mathbf{P}_{\underline{\kappa}}, \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}}}{\prod_{i=1}^3 \langle X^{\kappa_i}, Y^{\kappa_i} \rangle_{\kappa_i}} dx_\infty \\ &= \frac{\Gamma(\kappa_1 + 2) \Gamma(\kappa_2 + 2) \Gamma(\kappa_3 + 2)}{4\pi^2 \Gamma(\frac{\kappa_1 + \kappa_2 + \kappa_3}{2} + 2) \Gamma(\kappa_1^* + 1) \Gamma(\kappa_2^* + 1) \Gamma(\kappa_3^* + 1)} \cdot \langle \mathbf{P}_{\underline{\kappa}}, \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}} \\ &= (4\pi^2)^{-1} (1 + \kappa_1)(1 + \kappa_2)(1 + \kappa_3). \end{aligned}$$

The last equality follows from Lemma 4.11 below. Now let q be a prime in Σ^- . According to [Pra90], $\pi_{i,q} = \mu_i \mathrm{St}$ for $i = 1, 2, 3$ are unramified special representations with $\mu_1 \mu_2 \mu_3(q) = 1$. Since

$$L(s, \Pi_q, \mathrm{Ad}) = \zeta_q(s + 1)^3; \quad L(s, \Pi_q) = \zeta_q(s + 1/2)^2 \zeta_q(s + 3/2),$$

we obtain

$$I_q(\phi_q^D \otimes \tilde{\phi}_q^D) = \frac{L(1, \Pi_q, \mathrm{Ad})}{\zeta_q(2)^2 L(1/2, \Pi_q)} \cdot (1 + \mu_{1,q} \mu_{2,q} \mu_{3,q}(q)) = 2\zeta_q(1)^{-2}.$$

Substituting (4.19) and the above computation of $I_q(\phi_q^D \otimes \tilde{\phi}_q^D)$ into Ichino's formula, we obtain

$$\begin{aligned} & \frac{I(\rho(\check{\mathfrak{t}}_n)\phi_F^{D^*})^2}{\langle F^D, F^D \rangle \omega_F^{-1/2}(\widehat{N}_1^+)} \\ &= \text{vol}(\widehat{\mathcal{O}}_D^\times)^2 \cdot \frac{N^-}{48} \cdot \frac{\zeta_{\mathbf{Q}}(2)^2 \cdot 48^3}{8 \cdot 4\pi^2} \cdot \frac{L(1/2, \Pi)}{L(1, \Pi, \text{Ad})} \cdot \frac{\omega_{F,p}^{-1}(p^n)\alpha_p(F)^{2n}}{\prod_{i=1}^3 [\text{SL}_2(\mathbf{Z}) : \Gamma_0(N_i p^n)](N_i^+ p^n)^{\kappa_i/2}} \prod_{q \in \Sigma^-} \frac{2\zeta_q(1)^3}{\zeta_q(2)^3} \\ & \quad \times I_p^{\text{ord}}(\phi_p \otimes \tilde{\phi}_p, \check{\mathfrak{t}}_n) \prod_{q \notin \{p, \infty\} \cup \Sigma^-} I_q(\phi_q^* \otimes \tilde{\phi}_q^*), \end{aligned}$$

and the proposition follows. \square

Lemma 4.11. *We have*

$$\langle \mathbf{P}_{\underline{\kappa}}, \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}} = \frac{\Gamma(\frac{\kappa_1 + \kappa_2 + \kappa_3}{2} + 2)\Gamma(\kappa_1^* + 1)\Gamma(\kappa_2^* + 1)\Gamma(\kappa_3^* + 1)}{\Gamma(\kappa_1 + 1)\Gamma(\kappa_2 + 1)\Gamma(\kappa_3 + 1)}.$$

PROOF. Let $v_1 = X_1^{\kappa_1} \otimes Y_2^{\kappa_2} \otimes X_3^{\kappa_1^*} Y_3^{\kappa_2^*}$ and $v_2 = Y_1^{\kappa_1} \otimes X_2^{\kappa_2} \otimes X_3^{\kappa_2^*} Y_3^{\kappa_1^*}$. Let du be the Haar measure on $\text{SU}(2)(\mathbf{R})$ with the volume $\text{vol}(\text{SU}(2)(\mathbf{R}), du) = 1$. More precisely, du is given by

$$\begin{aligned} \int_{\text{SU}(2)(\mathbf{R})} \Phi(u) du &= \frac{1}{4\pi^2} \int_0^{2\pi} \int_0^{2\pi} \int_0^{\pi/2} \Phi(u) \sin 2\theta d\theta d\varphi d\rho, \\ (u &= \begin{pmatrix} \alpha & \beta \\ -\beta & \alpha \end{pmatrix}), \alpha = \cos \theta e^{i\varphi}, \beta = \sin \theta e^{i\rho} \end{aligned}$$

for $\Phi \in L^1(\text{SU}(2)(\mathbf{R}))$. Write $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_{\underline{\kappa}}$ for simplicity. Since $L_{\underline{\kappa}}(\mathbf{C})^{\text{SU}(2)(\mathbf{R})} = \mathbf{C} \cdot \mathbf{P}_{\underline{\kappa}}$, we see that

$$(4.20) \quad \int_{\text{SU}(2)(\mathbf{R})} \langle \rho_{\underline{\kappa}}(u)v_1, v_2 \rangle du \cdot \langle \mathbf{P}_{\underline{\kappa}}, \mathbf{P}_{\underline{\kappa}} \rangle = \langle v_1, \mathbf{P}_{\underline{\kappa}} \rangle \cdot \langle \mathbf{P}_{\underline{\kappa}}, v_2 \rangle.$$

By definition,

$$\begin{aligned} \mathbf{P}_{\underline{\kappa}} &= \sum_{n_1=0}^{\kappa_1^*} \sum_{n_2=0}^{\kappa_2^*} \sum_{n_3=0}^{\kappa_3^*} \binom{\kappa_1^*}{n_1} \binom{\kappa_2^*}{n_2} \binom{\kappa_3^*}{n_3} (-1)^{\kappa_1^* + \kappa_2^* + \kappa_3^* - n_1 - n_2 - n_3} \\ & \quad \times X_1^{\kappa_2^* - n_2 + n_3} Y_1^{\kappa_3^* + n_2 - n_3} \otimes X_2^{\kappa_3^* + n_1 - n_3} Y_2^{\kappa_1^* - n_1 + n_3} \otimes X_3^{\kappa_1^* - n_1 + n_2} Y_3^{\kappa_2^* + n_1 - n_2}. \end{aligned}$$

Then

$$\langle v_1, \mathbf{P}_{\underline{\kappa}} \rangle = (-1)^{\kappa_1 + \kappa_2} \binom{\kappa_3}{\kappa_1^*}^{-1}, \quad \langle \mathbf{P}_{\underline{\kappa}}, v_2 \rangle = (-1)^{\kappa_1 + \kappa_1^* + \kappa_3} \binom{\kappa_3}{\kappa_1^*}^{-1}.$$

Let $r = \frac{\kappa_1 + \kappa_2 + \kappa_3}{2} = \kappa_1^* + \kappa_2^* + \kappa_3^*$. A direct computation shows that

$$\begin{aligned} \int_{\text{SU}(2)(\mathbf{R})} \langle \rho_{\underline{\kappa}}(u)v_1, v_2 \rangle du &= (-1)^r \binom{\kappa_3}{\kappa_1^*}^{-1} \sum_{j=0}^{\kappa_1^*} \binom{\kappa_1^*}{j} \binom{\kappa_2^*}{j} (-1)^j \int_{\text{SU}(2)(\mathbf{R})} |\alpha \bar{\alpha}|^{r-j} |\beta \bar{\beta}|^j du \\ &= 2(-1)^r (2r+2)^{-1} \binom{\kappa_3}{\kappa_1^*}^{-1} \sum_{j=0}^{\kappa_1^*} (-1)^j \frac{\binom{\kappa_1^*}{j} \binom{\kappa_2^*}{j}}{\binom{r}{j}} \\ &= (-1)^r (r+1)^{-1} \binom{\kappa_3}{\kappa_1^*}^{-1} \frac{\Gamma(\kappa_1 + 1)\Gamma(k_1^* + 1)}{\Gamma(r+1)} \sum_{j=0}^{k_1^*} (-1)^j \binom{\kappa_2^*}{j} \binom{r-j}{\kappa_1} \\ &= (-1)^r (r+1)^{-1} \binom{\kappa_3}{\kappa_1^*}^{-1} \frac{\Gamma(\kappa_1 + 1)\Gamma(k_1^* + 1)}{\Gamma(r+1)} \cdot \binom{r - \kappa_2^*}{\kappa_1 - \kappa_2^*}. \end{aligned}$$

Substituting the above to (4.20), we obtain

$$\langle \mathbf{P}_{\underline{\kappa}}, \mathbf{P}_{\underline{\kappa}} \rangle_{\underline{\kappa}} = \frac{\Gamma(r+2)}{\Gamma(\kappa_1 + 1)\Gamma(k_1^* + 1)} \cdot \frac{\Gamma(k_1^* + 1)\Gamma(k_2^* + 1)}{\Gamma(\kappa_3 + 1)} \cdot \frac{\Gamma(k_3^* + 1)\Gamma(k_1^* + 1)}{\Gamma(\kappa_2 + 1)},$$

and the lemma follows. \square

Definition 4.12 (The Gross periods of Hida families). Suppose that \mathcal{F} is a primitive \mathbf{I} -adic Hida family which satisfies (CR, Σ^-). Let \mathcal{F}^D be a primitive Jacquet-Langlands lift of \mathcal{F} with the tame conductor $N_{\mathcal{F}} = N^- N_{\mathcal{F}}^+$. Put

$$\eta_{\mathcal{F}^D} := \mathbf{B}_{N_{\mathcal{F}}}(\mathcal{F}^D, \mathcal{F}^D) \in \mathbf{I},$$

where $\mathbf{B}_{N_{\mathcal{F}}}$ is the Hecke-equivariant perfect pairing defined in Definition 4.3. For each arithmetic point $Q \in \mathfrak{X}_{\mathbf{I}}^+$, writing $\eta_{\mathcal{F}^D_Q}$ for the specialization of $\eta_{\mathcal{F}^D}$ at Q , define the Gross' period $\Omega_{\mathcal{F}^D_Q}$ of \mathcal{F}^D_Q by

$$\Omega_{\mathcal{F}^D_Q} = 2^{k_Q} \|\mathcal{F}^D_Q\|_{\Gamma_0(N_{\mathcal{F}^D_Q})}^2 \cdot \frac{\mathcal{E}_p(\mathcal{F}^D_Q, \text{Ad})}{\eta_{\mathcal{F}^D_Q}} \cdot \varepsilon^{\Sigma^-}(\mathcal{F}^D_Q),$$

where $\mathcal{E}_p(\mathcal{F}^D_Q, \text{Ad})$ is the modified p -Euler factor in (3.10) and

$$\varepsilon^{\Sigma^-}(\mathcal{F}^D_Q) := \prod_{\ell | N_{\mathcal{F}}^+} \varepsilon(1/2, \pi_{\mathcal{F}^D_Q, \ell}) | N_{\mathcal{F}}^+ |_{\ell}^{\frac{2-k_Q}{2}} \in \overline{\mathbf{Z}}_{(p)}^\times.$$

is the prime-to- Σ^- part of the root number of \mathcal{F}^D_Q .³ We will see from Remark 7.8 that the canonical period is an integral multiple of the Gross period in the sense that there exists a non-zero $u \in \mathbf{I}$ such that $\Omega_{\mathcal{F}^D_Q} = u(Q) \cdot \Omega_{\mathcal{F}^D_Q}$ for each arithmetic point Q .

Corollary 4.13. *For each $\underline{Q} = (Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathcal{R}}^{\text{bal}}$ in the balanced range, we have the interpolation formula*

$$\Theta_{\mathbf{F}^{D^*}}(\underline{Q})^2 = 2^{\#(\Sigma^-)+1} N \cdot \frac{L(1/2, \Pi_{\underline{Q}})}{\Omega_{\mathcal{F}^D_{Q_1}} \Omega_{\mathcal{G}^D_{Q_2}} \Omega_{\mathcal{H}^D_{Q_3}}} \cdot \mathcal{S}_{\Pi_{\underline{Q}, p}}^{\text{bal}} \cdot \prod_{q | N^+} \mathcal{S}_{\Pi_{\underline{Q}, q}}^*$$

where $\mathcal{S}_{\Pi_{\underline{Q}, p}}^{\text{bal}}$ is the normalized p -adic zeta integrals given by

$$(4.21) \quad \mathcal{S}_{\Pi_{\underline{Q}, p}}^{\text{bal}} = I_p^{\text{ord}}(\phi_p, \check{\mathfrak{t}}_n) \cdot B_{\Pi_p^{\text{ord}}}^{[n]} \cdot \frac{\omega_{F, p}^{1/2}(-p^{2n}) |p|_p^{-n(k_1+k_2+k_3)}}{\alpha_p(F)^{2n} \zeta_p(2)^2}$$

and $\mathcal{S}_{\Pi_{\underline{Q}, q}}^*$ are the local zeta integrals at q sdefined in (3.27).

PROOF. To simplify our notation, we let $f_1 = f$, $f_2 = g$ and $f_3 = h$. For a finite prime q , we put $B_{\Pi_{F, q}} = \prod_{i=1}^3 B_{\pi_{f_i, q}}$. By definition, we have $B_{\Pi_{F, q}} = \omega_{F, q}^{1/2}(N_f^+) B_{\Pi_q}$ if $q \neq p$ and $B_{\Pi_{F, q}} = 1$ if $q \nmid pN$. At the place p , from Lemma 2.8 and the definition of $\mathcal{E}_p(f_i, \text{Ad})$ in (3.10), we see that

$$\frac{B_{\Pi_p^{\text{ord}}}^{[n]}}{B_{\Pi_{F, p}}} = \omega_{F, p}^{1/2}(-p^{-2n}) \prod_{i=1}^3 \frac{\alpha_{f_i, p} | \cdot |_p^{\frac{1}{2}}(p^{2n})}{\varepsilon(1/2, \pi_{f_i, p})} \cdot \frac{[\text{SL}_2(\mathbf{Z}) : \Gamma_0(p^{c_i})]}{(1+p^{-1})} \cdot \mathcal{E}_p(f_i, \text{Ad}).$$

³Here $\Omega_{\mathcal{F}^D_Q}$ is called the Gross period for \mathcal{F}^D_Q as it first appeared in the Gross' special value formula for modular forms over imaginary quadratic fields.

Let f_i° be the associated newform of f_i and $c_i = c(\pi_{f_i, p})$. Write $\|f_i^\circ\|^2$ for the Petersson norm $\|f_i^\circ\|_{\Gamma_0(N_i p^{2n})}^2$. From the above equation and the Petersson norm formula (2.18), we find that

$$\begin{aligned} & \frac{\omega_F^{-1/2}(\widehat{N}_1^+) \omega_{F,p}(p^{-n}) \alpha_p(F)^{2n}}{L(1, \Pi, \text{Ad}) \prod_{i=1}^3 [\text{SL}_2(\mathbf{Z}) : \Gamma_0(N_i p^{2n})] (N_i^+ p^{2n})^{\kappa_i/2}} \\ &= \omega_F^{-1/2}(\widehat{N}_1^+) \omega_{F,p}(p^{-n}) \alpha_p(F)^{2n} \prod_{q|Np} B_{\Pi_{F,q}} \prod_{i=1}^3 \frac{[\text{SL}_2(\mathbf{Z}) : \Gamma_0(p^{c_i})] p^{-2n}}{2^{k_i} \omega(f_i^\circ) \|f_i^\circ\|^2 (1+p^{-1}) (N_i^+ p^{2n})^{\kappa_i/2}}, \\ &= \omega_F^{1/2}(\widehat{N}_1^-) \omega_{F,p}^{1/2}(-1) \alpha_p(F)^{2n} \cdot B_{\Pi_p^{[n]}} \cdot \prod_{q|N} B_{\Pi_q} \prod_{i=1}^3 \frac{\varepsilon(1/2, \pi_{f_i, p})}{\omega(f_i^\circ) (N_i^+)^{\kappa_i/2}} \cdot \frac{1}{\alpha_{f_i, p} \cdot \left| \frac{1-k_i}{p} \right| (p^{2n})^{2k_i} \|f_i^\circ\|^2 \mathcal{E}_p(f_i, \text{Ad})} \\ &= \omega_{F,p}^{1/2}(-1) B_{\Pi_p^{[n]}} \cdot \prod_{q|N} B_{\Pi_q} \prod_{i=1}^3 \frac{1}{\varepsilon^{\Sigma^-}(f_i) 2^{k_i} \|f_i^\circ\|^2 \mathcal{E}_p(f_i, \text{Ad})} \prod_{q \in \Sigma^-} \frac{\omega_{F,q}^{1/2}(q)}{\varepsilon(1/2, \pi_{f_1, q}) \varepsilon(1/2, \pi_{f_2, q}) \varepsilon(1/2, \pi_{f_3, q})} \\ &= (-1)^{\#\Sigma^-} \cdot \omega_{F,p}^{1/2}(-1) \cdot B_{\Pi_p^{[n]}} \cdot \prod_{q|N} B_{\Pi_q} \prod_{i=1}^3 \frac{\langle \mathbf{U}_p^{-n} f_i^D, f_i^D \rangle_{N_i p^n}}{\Omega_{f_i^D}}. \end{aligned}$$

In the last equality, we have used Lemma 4.4 and the fact that for $q \in \Sigma^-$,

$$\varepsilon(1/2, \Pi_q) = \omega_{F,q}^{-1/2}(q) \varepsilon(1/2, \pi_{f_1, q}) \varepsilon(1/2, \pi_{f_2, q}) \varepsilon(1/2, \pi_{f_3, q}) = -1.$$

Substituting the above equation and the definition of $\mathcal{S}_{\Pi_q}^*$ in (3.27) to Proposition 4.10, we deduce from Proposition 4.9 that

$$\Theta_{\mathbf{F}^D, \mathbf{Q}}(\underline{Q})^2 = \frac{\text{vol}(\widehat{\mathcal{O}}_D^\times)^2}{\text{vol}(\widehat{R}_N^\times)^2} \cdot (-2)^{\#\Sigma^-} 2N^- \cdot \frac{L(1/2, \Pi)}{\Omega_{f^D} \Omega_{g^D} \Omega_{h^D}} \cdot \mathcal{S}_p^{\text{bal}} \prod_{q \in \Sigma^-} B_{\Pi_q} \cdot \frac{\zeta_q(1)^3}{\zeta_q(2)^3} \prod_{q|N^+} \mathcal{S}_{\Pi_q}^* \cdot \frac{|N|_q^2 \zeta_q(2)^2}{\zeta_q(1)^2}.$$

Therefore, we obtain the corollary by noting that

$$\frac{\text{vol}(\mathcal{O}_D^\times)}{\text{vol}(\widehat{R}_N^\times)} = \prod_{q|N^+} \frac{\zeta_q(1)}{|N|_q \zeta_q(2)},$$

and that for $q \in \Sigma^-$,

$$B_{\Pi_q} = \prod_{i=1}^3 \frac{\zeta_q(2) \langle \rho(\tau_q) W_{\pi_i}, \widetilde{W}_{\pi_i} \rangle}{\zeta_q(1) L(1, \pi_i, \text{Ad})} = \prod_{i=1}^3 \varepsilon(1/2, \pi_{i, q}) \frac{\zeta_q(2)}{\zeta_q(1)} = (-1) \frac{\zeta_q(2)^3}{\zeta_q(1)^3}.$$

This finishes the proof. \square

5. THE CALCULATION OF LOCAL ZETA INTEGRALS (I)

5.1. Notation and conventions. Let q be a finite prime. Let $G = \text{GL}_2(\mathbf{Q}_q)$ and $Z = \mathbf{Q}_q^\times$ be the center of G . Let B be the group of the upper triangular matrices of G and let N be the unipotent radical of B . Let $K = \text{GL}_2(\mathbf{Z}_q)$. Let π be an irreducible unitary generic admissible representation of G , define $\lambda(\pi) \in \mathbf{R}$ by

$$\lambda(\pi) = \begin{cases} |\lambda| & \text{if } \pi = \chi_1 |\cdot|^\lambda \boxplus \chi_2 |\cdot|^{-\lambda} \text{ with } \chi_1, \chi_2 \text{ unitary and } \lambda \in \mathbf{R}, \\ -\frac{1}{2} & \text{if } \pi \text{ is a discrete series.} \end{cases}$$

Recall that $\mathcal{W}(\pi) = \mathcal{W}(\pi, \psi_{\mathbf{Q}_q})$ is the Whittaker model of π with respect to $\psi_{\mathbf{Q}_q}$. It is well known that for any $W \in \mathcal{W}(\pi)$ and $\epsilon > 0$, there exists a $\Phi_\epsilon \in \mathcal{S}(\mathbf{Q}_q)$ with

$$(5.1) \quad W\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) = |y|^{\frac{1}{2} - \lambda(\pi) - \epsilon} \Phi_\epsilon(y).$$

For characters $\chi, v : \mathbf{Q}_q^\times \rightarrow \mathbf{C}^\times$, let $\mathcal{B}(\chi, v)$ denote the induced representation given by

$$\mathcal{B}(\chi, v) = \left\{ \text{smooth functions } f : G \rightarrow \mathbf{C} \mid f\left(\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g\right) = \chi(a)v(d) \left| \frac{a}{d} \right|^{\frac{1}{2}} f(g) \right\}.$$

Let $\langle \cdot, \cdot \rangle : \mathcal{B}(\chi, v) \times \mathcal{B}(\chi^{-1}, v^{-1}) \rightarrow \mathbf{C}$ be the G -invariant perfect pairing given by

$$\langle f, f' \rangle := \int_K f(k) f'(k) dk,$$

where dk is the Haar measure with $\text{vol}(K, dk) = 1$. If $\chi v^{-1} \neq |\cdot|^{-1}$, then we let $\mathcal{B}(\chi, v)_0$ be the unique irreducible sub-representation of $\mathcal{B}(\chi, v)$ and let $\mathcal{B}(v, \chi)^0$ be the unique irreducible quotient of $\mathcal{B}(v, \chi)$. It is well known that $\mathcal{B}(\chi, v)_0 = \mathcal{B}(\chi, v)$ and $\mathcal{B}(v, \chi)^0 = \mathcal{B}(v, \chi)$ unless $\chi v^{-1} = |\cdot|^{-1}$. The above pairing $\langle \cdot, \cdot \rangle$ induces a G -invariant perfect pairing $\langle \cdot, \cdot \rangle : \mathcal{B}(\chi, v)_0 \times \mathcal{B}(\chi^{-1}, v^{-1})^0 \rightarrow \mathbf{C}$.

Intertwining operator. Define the normalized intertwining operator $M^*(v, \chi, s) : \mathcal{B}(v|\cdot|^s, \chi|\cdot|^{-s}) \rightarrow \mathcal{B}(\chi|\cdot|^{-s}, v|\cdot|^s)$ by

$$M^*(v, \chi, s)f := \gamma(2s, v\chi^{-1}) \int_{\mathbf{Q}_p} f\left(\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} g\right) dx \quad (g \in G).$$

Here $\gamma(s, -)$ is the γ -factor as in (2.9), and the integral in the right hand side is convergent absolutely for $\text{Re } s$ sufficiently large and has analytic continuation to all $s \in \mathbf{C}$ (cf. [Bum97, Proposition 4.5.7]). Let $\delta : G \rightarrow \mathbf{R}_+$

be the function given by $\delta\left(\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} k\right) = |ad^{-1}|$ for $k \in K$. If $\chi v^{-1} \neq |\cdot|^{-1}$, then $M^*(v, \chi, s)|_{s=0}$ factors through $\mathcal{B}(v, \chi)^0$, and hence we have a well-defined map $M^*(v, \chi) : \mathcal{B}(v, \chi)^0 \rightarrow \mathcal{B}(\chi, v)_0$ given by

$$(5.2) \quad M^*(v, \chi)f := M^*(v, \chi, s)(f\delta^s)|_{s=0}.$$

An integration formula. The following integration formula will be used frequently in our computation. For $F \in L^1(ZN \backslash G)$,

$$\begin{aligned} \int_{ZN \backslash G} F(g) dg &= \int_K \int_{\mathbf{Q}_q^\times} F\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} k\right) \frac{d^\times y}{|y|} dk \\ &= \frac{\zeta_q(2)}{\zeta_q(1)} \int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q} F\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}\right) dx \frac{d^\times y}{|y|} \end{aligned}$$

(cf. [MV10, 3.1.6, page 206]).

5.2. Local trilinear integrals and Rankin-Selberg integrals. Let π_1, π_2 and π_3 be irreducible unitary generic admissible representation of G with central characters ω_1, ω_2 and ω_3 . Suppose that $\omega_1 \omega_2 \omega_3 = 1$ and that π_3 is a constituent (an irreducible subquotient) of $\mathcal{B}(\chi_3, v_3)$. Assume further that the following condition holds for $(\pi_1, \pi_2; \pi_3)$:

$$(Hb) \quad \lambda(\pi_1) + \lambda(\pi_2) + |\lambda(\pi_3)| < 1/2.$$

Put

$$\mathcal{J} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \in \text{GL}_2(\mathbf{Q}_q).$$

For $(W_1, W_2, f_3) \in \mathcal{W}(\pi_1) \times \mathcal{W}(\pi_2) \times \mathcal{B}(\chi_3, v_3)$, define the local Rankin-Selberg integrals by

$$\begin{aligned} \Psi(W_1, W_2, f_3) &= \int_{ZN \backslash G} W_1(g) W_2(\mathcal{J}g) f_3(g) dg; \\ \tilde{\Psi}(\tilde{W}_1, \tilde{W}_2, \tilde{f}_3) &= \int_{ZN \backslash G} \tilde{W}_1(\mathcal{J}g) \tilde{W}_2(g) \tilde{f}_3(g) dg. \end{aligned}$$

The above integrals converge absolutely under the assumption (Hb). For $\tilde{W}_1 \in \mathcal{W}(\tilde{\pi}_1)$, $\tilde{W}_2 \in \mathcal{W}(\tilde{\pi}_2)$ and $\tilde{f}_3 \in \mathcal{B}(\chi_3^{-1}, v_3^{-1})$, define the local trilinear integral by

$$\mathcal{I}_q(W_1 \otimes W_2 \otimes f_3, \tilde{W}_1 \otimes \tilde{W}_2 \otimes \tilde{f}_3) := \int_{Z \backslash G} \langle \rho(g) W_1, \tilde{W}_1 \rangle \langle \rho(g) W_2, \tilde{W}_2 \rangle \langle \rho(g) f_3, \tilde{f}_3 \rangle dg.$$

The following result is a generalization of [MV10, Lemma 3.4.2]. We provide a different and more elementary proof and replace the assumption on the temperedness with a much weaker hypothesis (Hb).

Proposition 5.1. *With the assumption (Hb) for $(\pi_1, \pi_2; \pi_3)$, we have*

$$\mathcal{I}_q(W_1 \otimes W_2 \otimes f_3, \tilde{W}_1 \otimes \tilde{W}_2 \otimes \tilde{f}_3) = \zeta_q(1) \cdot \Psi(W_1, W_2, f_3) \cdot \tilde{\Psi}(\tilde{W}_1, \tilde{W}_2, \tilde{f}_3).$$

PROOF. Denote by $\Psi : ZN \rightarrow \mathbf{C}^\times$ the character $\omega_2 \boxtimes \psi_{\mathbf{Q}_q}$. Let $\langle\langle \cdot, \cdot \rangle\rangle : L^2(ZN \setminus G, \Psi) \otimes L^2(ZN \setminus G, \Psi^{-1}) \rightarrow \mathbf{C}$ be the G -equivariant bilinear pairing given by

$$\langle\langle F, F' \rangle\rangle = \int_{ZN \setminus G} F(g)F'(g)dg.$$

We put

$$\begin{aligned} F_1(g) &= W_1(g)f_3(g), \quad W_4(g) = \widetilde{W}_2(g) \in L^2(ZN \setminus G, \Psi); \\ F_2(g) &= \widetilde{W}_1(\mathcal{J}g)\widetilde{f}_3(g), \quad W_3(g) = W_2(\mathcal{J}g) \in L^2(ZN \setminus G, \Psi^{-1}). \end{aligned}$$

Then one verifies that

$$\langle\langle \rho(g)F_1, F_2 \rangle\rangle = \langle \rho(g)W_1, \widetilde{W}_1 \rangle \langle \rho(g)f_3, \widetilde{f}_3 \rangle,$$

and hence, it is equivalent to showing that

$$(5.3) \quad \mathcal{I}_q(W_1 \otimes W_2 \otimes f_3, \widetilde{W}_1 \otimes \widetilde{W}_2 \otimes \widetilde{f}_3) = \int_{ZN \setminus G} \langle\langle \rho(g)F_1, F_2 \rangle\rangle \langle \rho(g)W_3, W_4 \rangle dg = \zeta_q(1) \langle\langle F_1, W_3 \rangle\rangle \langle\langle F_2, W_4 \rangle\rangle.$$

Put

$$K_n = \bigcup_{i=0}^n K \begin{pmatrix} q^i & 0 \\ 0 & 1 \end{pmatrix} K \subset G.$$

First we claim that if $y_1, y_2 \in \mathbf{Q}_q^\times$, then

$$(5.4) \quad \begin{pmatrix} y_2^{-1}y_1 & y_2^{-1}x \\ 0 & 1 \end{pmatrix} \in ZK_n \iff q^{-n} \leq |y_2^{-1}y_1| \leq q^n \text{ and } |x^2| \leq q^n |y_1y_2|.$$

To see the claim, we note that if $|x| \leq 1$ or $|x| \leq |y|$, then

$$\begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix} \in K \begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} K,$$

and if $|x| > 1$ and $|x| > |y|$, then

$$\begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix} \in K \begin{pmatrix} x^{-2}y & 0 \\ 0 & 1 \end{pmatrix} K.$$

By the Cartan decomposition, we find $\begin{pmatrix} y_2^{-1}y_1 & y_2^{-1}x \\ 0 & 1 \end{pmatrix} \in ZK_n$ if and only if $|x| \leq \max\{|y_1|, |y_2|\}$ and $q^{-n} \leq |y_2^{-1}y_1| \leq q^n$ or $|x| > \max\{|y_1|, |y_2|\}$ and $q^{-n} \leq |x^{-2}y_1y_2| \leq 1$, and this proves the claim.

Now we proceed to prove the equation (5.3). Let $\mathbb{1}_{K_n}$ be the characteristic function of ZK_n and set

$$\mathcal{I}_n = \int_{Z \setminus G} \langle\langle \rho(g)F_1, F_2 \rangle\rangle \langle \rho(g)W_3, W_4 \rangle \mathbb{1}_{K_{2n}}(g) dg.$$

By a formal computation, we obtain

$$\begin{aligned} \mathcal{I}_n &= \int_{Z \setminus G} \int_{ZN \setminus G} F_1(hg)F_2(h) \cdot \langle \rho(g)W_3, W_4 \rangle \mathbb{1}_{K_{2n}}(g) dh dg \\ &= \int_{ZN \setminus G} \int_{Z \setminus G} F_1(hg)F_2(h) \cdot \langle \rho(g)W_3, W_4 \rangle \mathbb{1}_{K_{2n}}(g) dg dh \\ &= \int_{(ZN \setminus G)^2} \int_N \psi(u)F_1(g)F_2(h) \cdot \langle h^{-1}ugW_3, W_4 \rangle \cdot \mathbb{1}_{K_{2n}}(h^{-1}ug) dudgdh \\ &= \int_K \int_K \int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q} \psi_{\mathbf{Q}_q}(u)F_1\left(\begin{pmatrix} y_1 & 0 \\ 0 & 1 \end{pmatrix} k_1\right)F_2\left(\begin{pmatrix} y_2 & 0 \\ 0 & 1 \end{pmatrix} k_2\right) \langle \rho\left(\begin{pmatrix} y_2^{-1}y_1 & y_2^{-1}x \\ 0 & 1 \end{pmatrix} k_1\right)W_3, \rho(k_2)W_4 \rangle \\ &\quad \times \mathbb{1}_{K_{2n}}\left(\begin{pmatrix} y_2^{-1}y_1 & y_2^{-1}x \\ 0 & 1 \end{pmatrix}\right) dx \frac{d^\times y_1}{|y_1|} \frac{d^\times y_2}{|y_2|} dk_1 dk_2. \end{aligned}$$

To justify the computation, it suffices to show that the integral

$$\int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q} \psi_{\mathbf{Q}_q}(x)F_1'\left(\begin{pmatrix} y_1 & 0 \\ 0 & 1 \end{pmatrix}\right)F_2'\left(\begin{pmatrix} y_2 & 0 \\ 0 & 1 \end{pmatrix}\right) \cdot \langle \left(\begin{pmatrix} y_2^{-1}y_1 & y_2^{-1}x \\ 0 & 1 \end{pmatrix} W_3', W_4' \right) \mathbb{1}_{K_{2n}}\left(\begin{pmatrix} y_2^{-1}y_1 & y_2^{-1}x \\ 0 & 1 \end{pmatrix}\right) dx \frac{d^\times y_1}{|y_1|} \frac{d^\times y_2}{|y_2|}$$

is absolutely convergent, where $F'_1 = \rho(k_1)F_1$, $F'_2 = \rho(k_2)F'_2$, $W'_3 = \rho(k_1)W_3$ and $W'_4 = \rho(k_2)W_4$. By (5.4), we can deduce that for any $\epsilon > 0$ there exist constants C_ϵ and M such that

$$\begin{aligned} & \int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q} \left| F'_1 \left(\begin{pmatrix} y_1 & 0 \\ 0 & 1 \end{pmatrix} \right) F'_2 \left(\begin{pmatrix} y_2 & 0 \\ 0 & 1 \end{pmatrix} \right) \left| \left\langle \rho \left(\begin{pmatrix} y_2^{-1} y_1 & y_2^{-1} x \\ 0 & 1 \end{pmatrix} \right) W'_3, W'_4 \right\rangle \right| \mathbb{I}_{K_{2n}} \left(\begin{pmatrix} y_2^{-1} y_1 & y_2^{-1} x \\ 0 & 1 \end{pmatrix} \right) dx \frac{d^\times y_1}{|y_1|} \frac{d^\times y_2}{|y_2|} \\ & < C_\epsilon \iint_{\substack{|y_1| \leq M, |y_2| \leq M, \\ q^{-n} \leq |y_2^{-1} y_1| \leq q^n}} \int_{|x|^2 \leq |y_1 y_2| q^{2n}} |y_1 y_2|^{1-\lambda_1-\lambda_3-\epsilon} |y_2^{-1} y_1|^{\frac{1}{2}-\lambda_2-\epsilon} dx \frac{d^\times y_1}{|y_1|} \frac{d^\times y_2}{|y_2|} \\ & < C_\epsilon q^{n(3/2-\lambda_2)} \int_{|y_1| \leq M} \int_{|y_2| \leq M} |y_1 y_2|^{\frac{1}{2}-\lambda_1-\lambda_3-2\epsilon} d^\times y_1 d^\times y_2. \end{aligned}$$

For $(g, h) \in G \times G$, put

$$\mathcal{A}_n(g, h) := \int_{\mathbf{Q}_q} \psi(x) \left\langle \rho \left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} g \right) W_3, \rho(h) W_4 \right\rangle \mathbb{I}_{K_{2n}}(h^{-1} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} g) dx.$$

Then we have

$$\begin{aligned} \mathcal{A}_n \left(\begin{pmatrix} y_1 & 0 \\ 0 & 1 \end{pmatrix} k_1, \begin{pmatrix} y_2 & 0 \\ 0 & 1 \end{pmatrix} k_2 \right) &= \int_{\mathbf{Q}_q} \psi_{\mathbf{Q}_q}(x) \left\langle \rho \left(\begin{pmatrix} y_1 & x \\ 0 & 1 \end{pmatrix} k_1 \right) W_3, \rho \left(\begin{pmatrix} y_2 & 0 \\ 0 & 1 \end{pmatrix} k_2 \right) W_4 \right\rangle \varphi_{2n} \left(\begin{pmatrix} y_1 y_2^{-1} & y_2^{-1} x \\ 0 & 1 \end{pmatrix} \right) dx \\ &= \int_{\mathbf{Q}_q} \int_{\mathbf{Q}_q^\times} \psi_{\mathbf{Q}_q}((1-\nu)x) W_3 \left(\begin{pmatrix} \nu y_1 & 0 \\ 0 & 1 \end{pmatrix} k_1 \right) W_4 \left(\begin{pmatrix} \nu y_2 & 0 \\ 0 & 1 \end{pmatrix} k_2 \right) \mathbb{I}_{q^{r-n} \mathbf{Z}_q}(x) d^\times \nu dx \\ &= \int_{\mathbf{Q}_q^\times} W_3 \left(\begin{pmatrix} \nu y_1 & 0 \\ 0 & 1 \end{pmatrix} k_1 \right) W_4 \left(\begin{pmatrix} \nu y_2 & 0 \\ 0 & 1 \end{pmatrix} k_2 \right) \mathbb{I}_{1+q^{n-r} \mathbf{Z}_q}(\nu) |q^{r-n}| d^\times \nu, \end{aligned}$$

where $r = \lfloor \frac{v_p(y_1 y_2)}{2} \rfloor$. Therefore, there exists a positive integer m_0 such that if $v_p(y_1 y_2) < 2n - m_0$, then

$$\mathcal{A}_n \left(\begin{pmatrix} y_1 & 0 \\ 0 & 1 \end{pmatrix} k_1, \begin{pmatrix} y_2 & 0 \\ 0 & 1 \end{pmatrix} k_2 \right) = \zeta_q(1) W_3 \left(\begin{pmatrix} y_1 & 0 \\ 0 & 1 \end{pmatrix} k_1 \right) W_4 \left(\begin{pmatrix} y_2 & 0 \\ 0 & 1 \end{pmatrix} k_2 \right).$$

On the other hand, if $v_p(y_1 y_2) \geq 2n - m_0$, we have

$$\begin{aligned} \left| \mathcal{A}_n \left(\begin{pmatrix} y_1 & 0 \\ 0 & 1 \end{pmatrix} k_1, \begin{pmatrix} y_2 & 0 \\ 0 & 1 \end{pmatrix} k_2 \right) \right| &<_{W_2, \widetilde{W}_w, \epsilon} q^{n-r} |y_1 y_2|^{\frac{1}{2}-\lambda_2-\epsilon} \int_{\mathbf{Q}_q^\times} |\nu|^{1-2\lambda_2-2\epsilon} \mathbb{I}_{1+q^{n-r} \mathbf{Z}_q}(\nu) d^\times \nu \\ &< C_\epsilon \cdot q^{m_0/2} \cdot |y_1 y_2|^{\frac{1}{2}-\lambda_2-\epsilon}. \end{aligned}$$

We thus have

$$\begin{aligned} \mathcal{I}_n &= \int_K \int_K \int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q^\times} F_1 \left(\begin{pmatrix} y_1 & 0 \\ 0 & 1 \end{pmatrix} k_1 \right) F_2 \left(\begin{pmatrix} y_2 & 0 \\ 0 & 1 \end{pmatrix} k_2 \right) \cdot \mathcal{A}_n \left(\begin{pmatrix} y_1 & 0 \\ 0 & 1 \end{pmatrix} k_1, \begin{pmatrix} y_2 & 0 \\ 0 & 1 \end{pmatrix} k_2 \right) \frac{d^\times y_1 d^\times y_2}{|y_1 y_2|} dk_1 dk_2 \\ &= \zeta_q(1) \int_K \int_K \int_{\substack{q^{-2n} \leq |y_1 y_2^{-1}| \leq q^{2n}, \\ |y_1 y_2| > |q|^{2n-m_0}}} F_1 \otimes W_3 \left(\begin{pmatrix} y_1 & 0 \\ 0 & 1 \end{pmatrix} k_1 \right) F_2 \otimes W_4 \left(\begin{pmatrix} y_2 & 0 \\ 0 & 1 \end{pmatrix} k_2 \right) \frac{d^\times y_1 d^\times y_2}{|y_1 y_2|} dk_1 dk_2 + B_n, \end{aligned}$$

where

$$\begin{aligned} |B_n| &< C'_\epsilon \int_{\substack{q^{-2n} \leq |y_1 y_2^{-1}| \leq q^{2n}, \\ |y_1 y_2| \leq |q|^{2n-m_0}}} |y_1 y_2|^{\frac{1}{2}-\lambda_1-\lambda_2-\lambda_3-2\epsilon} d^\times y_1 d^\times y_2 \\ &< C''_\epsilon |q|^{2n(\frac{1}{2}-\lambda_1-\lambda_2-\lambda_3-2\epsilon)} (4n+1). \end{aligned}$$

It follows that

$$\int_{Z_N \backslash G} \langle \rho(g) F_1, F_2 \rangle \langle \rho(g) W_3, W_4 \rangle dg = \lim_{n \rightarrow \infty} \mathcal{I}_n = \zeta_q(1) \int_{Z_N \backslash G} F_1(g) W_3(g) dg \int_{Z_N \backslash G} F_2(h) W_4(h) dh.$$

This finishes the proof of (5.3). \square

Denote by $L(s, \pi_1 \otimes \pi_2)$ the local L -factor and by $\varepsilon(s, \pi_1 \otimes \pi_2) := \varepsilon(s, \pi_1 \otimes \pi_2, \psi_{\mathbf{Q}_q})$ the ε -factors attached to $\pi_1 \times \pi_2$ defined in [GJ78]. Define the γ -factor

$$(5.5) \quad \gamma(s, \pi_1 \otimes \pi_2) := \varepsilon(s, \pi_1 \otimes \pi_2) \frac{L(1-s, \tilde{\pi}_1 \otimes \tilde{\pi}_2)}{L(s, \pi_1 \otimes \pi_2)}.$$

The following corollary is the core of our calculations of local zeta integrals $I_v(\phi_v^* \otimes \tilde{\phi}_v^*)$ at the non-archimedean places.

Corollary 5.2. *Suppose that $(\pi_1, \pi_2; \pi_3)$ satisfies (Hb) and that $\chi_3 v_3^{-1} \neq |\cdot|$. If $\tilde{W}_1 = W_1 \otimes \omega_1^{-1}$, $\tilde{W}_2 = W_2 \otimes \omega_2^{-1}$ and $\tilde{f}_3 = M^*(\chi_3, v_3) f_3 \otimes \omega_3^{-1}$, then*

$$\mathcal{I}_q(W_1 \otimes W_2 \otimes f_3, \tilde{W}_1 \otimes \tilde{W}_2 \otimes \tilde{f}_3) = \zeta_q(1) \chi_3(-1) \cdot \gamma(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3) \cdot \Psi(W_1, W_2, f_3)^2.$$

PROOF. This is an immediate consequence of the local functional equation of $\mathrm{GL}(2) \times \mathrm{GL}(2)$ in [Jac72]. With the notation of [Jac72, page 12], we may assume that

$$f_3(g) = \chi_3 |\cdot|^{s+\frac{1}{2}} (\det g) \cdot z(\chi_3 v_3^{-1} |\cdot|^{2s+1}, \rho(g) \Phi) \cdot \frac{1}{L(2s+1, \chi_3 v_3^{-1})} \Big|_{s=0}$$

is the Godement section attached to a Bruhat-Schwartz function Φ on \mathbf{Q}_q^2 . Since $\chi_3 v_3^{-1} \neq |\cdot|$, one verifies that

$$M^*(\chi_3, v_3) f_3(g) = v_3 |\cdot|^{-s+\frac{1}{2}} (\det g) \cdot z(v_3 \chi_3^{-1} |\cdot|^{-2s+1}, \rho(g) \hat{\Phi}) \cdot \frac{1}{L(2s+1, \chi_3 v_3^{-1})} \Big|_{s=0},$$

where $\hat{\Phi}$ is the Fourier transform of Φ defined in [Jac72, Theorem 14.2 (3)]. Under the hypothesis (Hb), we have

$$\begin{aligned} \Psi(W_1, W_2, f_3) &= \frac{\Psi(s, W_1, W_2, \Phi)}{L(2s+1, \chi_3 v_3^{-1})} \Big|_{s=0}, \\ \tilde{\Psi}(W_1, W_2, M^*(\chi_3, v_3) f_3) &= \frac{\tilde{\Psi}(s, \tilde{W}_1, \tilde{W}_2, \hat{\Phi})}{L(2s+1, \chi_3 v_3^{-1})} \Big|_{s=0}, \end{aligned}$$

where $\Psi(s, W_1, W_2, \Phi)$ and $\tilde{\Psi}(s, W_1, W_2, \hat{\Phi})$ are defined in [Jac72, (14.5) and (14.6)].

Therefore, from [Jac72, Theorem 14.8] we can deduce that

$$\begin{aligned} \tilde{\Psi}(\tilde{W}_1, \tilde{W}_2, \tilde{f}_3) &= \omega_1(-1) v_3(-1) \Psi(W_1, W_2, M^*(\chi_3, v_3) f_3) \\ &= \omega_1 \omega_2 v_3(-1) \gamma(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3) \Psi(W_1, W_2, f_3). \end{aligned} \quad \square$$

5.3. The calculation of the p -adic zeta integrals.

5.3.1. Preliminaries. We follow the notation in §3.7. Let $(f, g, h) = (\mathbf{f}_{Q_1}, \mathbf{g}_{Q_2}, \mathbf{h}_{Q_3})$ be the specialization of the triple of Hida families at a classical point $\underline{Q} = (Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathcal{R}}^{\mathrm{cls}} := \mathfrak{X}_{\mathbf{I}_1}^{\mathrm{cls}} \times \mathfrak{X}_{\mathbf{I}_2}^{\mathrm{cls}} \times \mathfrak{X}_{\mathbf{I}_3}^{\mathrm{cls}}$. Let $\pi_1 = \pi_{f,p} \otimes \omega_{F,p}^{-1/2}$, $\pi_2 = \pi_{g,p}$ and $\pi_3 = \pi_{h,p}$ of the central characters $\omega_1 = \omega_{g,p}^{-1} \omega_{h,p}^{-1}$, $\omega_2 = \omega_{g,p}$ and $\omega_3 = \omega_{h,p}$ respectively. Let $\Pi_{\underline{Q},p} := \pi_1 \times \pi_2 \times \pi_3$. For $i = 1, 2, 3$, since $\pi_{i,p}$ contains a non-zero ordinary vector, by Proposition 2.2 π_i must be a constituent of the induced representation $\mathcal{B}(v_i, \chi_i)$ with $\mathcal{V}_{\pi_i}^{\mathrm{ord}}(\chi_i) \neq \{0\}$. In view of the discussion in Remark 2.5, we have $\chi_1 = \alpha_{f,p} \omega_{F,p}^{-1/2}$, $\chi_2 = \alpha_{g,p}$ and $\chi_3 = \alpha_{h,p}$ with $\alpha_{\gamma,p}$ unramified characters defined there, and the ordinary assumption implies that $\chi_i v_i^{-1} \neq |\cdot|^{-1}$. Recall that if we let $\xi_i \in \mathcal{V}_{\pi_i}^{\mathrm{ord}}(\chi_i)$ and $\tilde{\xi}_i \in \mathcal{V}_{\tilde{\pi}_i}^{\mathrm{ord}}(v_i^{-1})$ be nonzero ordinary vectors for $i = 1, 2, 3$, then

$$\phi_p = \xi_1 \otimes \xi_2 \otimes \xi_3 \quad \text{and} \quad \tilde{\phi}_p = \tilde{\xi}_1 \otimes \tilde{\xi}_2 \otimes \tilde{\xi}_3.$$

Put

$$w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}; \quad t_n = \begin{pmatrix} 0 & p^{-n} \\ -p^n & 0 \end{pmatrix} \in \mathrm{SL}_2(\mathbf{Q}_p).$$

We introduce the normalized ordinary section in the induced representations and compute its local pairing.

Lemma 5.3. *Let π be a constituent of the induced representation $\mathcal{B}(v, \chi)$ of $\mathrm{GL}_2(\mathbf{Q}_p)$ with the central character ω . Suppose that $\chi v^{-1} \neq |\cdot|^{-1}$. Let $f^{\mathrm{ord}} \in \mathcal{B}(v, \chi)$ be the unique section such that (i) f^{ord} is supported in $BwN(\mathbf{Z}_p)$ (ii) $f^{\mathrm{ord}}(g) = 1$ for all $g \in wN(\mathbf{Z}_p)$. Then*

$$f^{\mathrm{ord}} \in \mathcal{B}(v, \chi)^{\mathrm{ord}}(\chi).$$

We call f^{ord} the normalized ordinary section. Moreover, put

$$\tilde{f}^{\mathrm{ord}} := M^*(v, \chi) f^{\mathrm{ord}} \otimes \omega^{-1} \in \mathcal{B}(v^{-1}, \chi^{-1})^{\mathrm{ord}}(v^{-1}).$$

For $n \geq \max\{1, c(\pi_p)\}$, we have

$$\langle \rho(t_n) f^{\mathrm{ord}}, \tilde{f}^{\mathrm{ord}} \rangle = \frac{\omega(p^{-n}) \zeta_p(2) \chi|\cdot|^{\frac{1}{2}}(p^{2n})}{\zeta_p(1)} \cdot \gamma(0, v\chi^{-1}).$$

In particular, if W^{ord} is the normalized ordinary Whittaker function in Corollary 2.3, then

$$(5.6) \quad \frac{\langle \rho(t_n) W^{\mathrm{ord}}, W^{\mathrm{ord}} \otimes \omega^{-1} \rangle}{\langle \rho(t_n) f^{\mathrm{ord}}, \tilde{f}^{\mathrm{ord}} \rangle} = \frac{\chi(-1) \zeta_p(2)}{\zeta_p(1)}.$$

PROOF. It is straightforward to verify that $f^{\mathrm{ord}} \in \mathcal{B}^{\mathrm{ord}}(v, \chi)^{\mathrm{ord}}(\chi)$ is an \mathbf{U}_p -eigenfunction with eigenvalue $\chi|\cdot|^{-\frac{1}{2}}$. By the integration formula [MV10, (3.2) page 207], $\langle \rho(t_n) f^{\mathrm{ord}}, \tilde{f}^{\mathrm{ord}} \rangle$ equals

$$\begin{aligned} \langle f^{\mathrm{ord}}, \rho\left(\begin{pmatrix} 0 & -p^{-n} \\ p^n & 0 \end{pmatrix}\right) \tilde{f}^{\mathrm{ord}} \rangle &= \frac{\zeta_p(2)}{\zeta_p(1)} \int_{\mathbf{Q}_p} f^{\mathrm{ord}}\left(w \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}\right) \tilde{f}^{\mathrm{ord}}\left(w \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & -p^{-n} \\ p^n & 0 \end{pmatrix}\right) dx \\ &= \frac{\zeta_p(2)}{\zeta_p(1)} f^{\mathrm{ord}}(w) \tilde{f}^{\mathrm{ord}}\left(\begin{pmatrix} p^n & 0 \\ 0 & p^{-n} \end{pmatrix}\right) \\ &= \omega(p^{-n}) \frac{\zeta_p(2)}{\zeta_p(1)} \chi|\cdot|^{\frac{1}{2}}(p^{2n}) \gamma(0, v\chi^{-1}). \end{aligned}$$

The ratio of local pairings of ordinary Whittaker functions and ordinary sections is computed by the above and Lemma 2.8. \square

5.3.2. *The unbalanced case.* Suppose that Q is in the unbalanced range $\mathfrak{X}_{\mathcal{R}}^f$. We apply Corollary 5.2 to calculate the normalized p -adic zeta integral $\mathcal{S}_{\Pi_p}^{\mathrm{unb}}$ in (3.28).

Proposition 5.4 (p -adic zeta integral in the unbalanced case). *Define*

$$\mathcal{E}_f(\Pi_{Q,p}) := \frac{1}{\varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1)} \cdot \frac{L(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1)}{L(1/2, \pi_2 \otimes \pi_3 \otimes v_1)}.$$

Then

$$\mathcal{S}_{\Pi_{Q,p}}^{\mathrm{unb}} = \mathcal{E}_f(\Pi_{Q,p}) \cdot \frac{1}{L(1/2, \Pi_{Q,p})}.$$

PROOF. We write $\Pi_p = \Pi_{Q,p}$ for brevity. It is equivalent to proving that

$$(5.7) \quad L(1/2, \Pi_p) \cdot I_p^{\mathrm{ord}}(\phi_p^* \otimes \tilde{\phi}_p^*, \mathbf{t}_n) = \mathcal{E}_f(\Pi_p) \cdot \frac{\chi_1 v_1^{-1} |\cdot|(-p^{2n})}{B_{\Pi_p^{\mathrm{ord}}}^{[n]}} \cdot \frac{\zeta_p(2)^2}{\zeta_p(1)^2}$$

for $n \geq \max\{c(\pi_1), c(\pi_2), c(\pi_3), 1\}$, where $I_p^{\mathrm{ord}}(\phi_p^* \otimes \tilde{\phi}_p^*, \mathbf{t}_n)$ is the local zeta integral defined in (3.24). We first treat the case where either (i) π_1 is principal series or (ii) π_2 or π_3 is discrete series. Then it is known that $(\pi_2, \pi_3; \pi_1)$ satisfies (Hb) since each π_i is a local component of a cuspidal automorphic representation of $\mathrm{GL}_2(\mathbf{A})$. Consider the realizations

$$\mathcal{V}_{\Pi_p} := \mathcal{B}(v_1, \chi_1)^0 \boxtimes \mathcal{W}(\pi_2) \boxtimes \mathcal{W}(\pi_3); \quad \mathcal{V}_{\tilde{\Pi}_p} := \mathcal{B}(v_1^{-1}, \chi_1^{-1})_0 \boxtimes \mathcal{W}(\tilde{\pi}_2) \boxtimes \mathcal{W}(\tilde{\pi}_3)$$

of Π_p and the contragredient representation $\tilde{\Pi}_p$. For $i = 1, 2, 3$, let $W_i^{\mathrm{ord}} = W_{\pi_i}^{\mathrm{ord}} \in \mathcal{W}^{\mathrm{ord}}(\pi_i)(\chi_i)$ be the normalized ordinary Whittaker functions such that $W_{\pi_i}^{\mathrm{ord}}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) = \chi_i |\cdot|^{\frac{1}{2}}(y) \mathbb{I}_{\mathbf{Z}_p}(y)$ in Corollary 2.3; let $f_i^{\mathrm{ord}} \in \mathcal{B}(v_i, \chi_i)^{\mathrm{ord}}(\chi_i)$ be the normalized ordinary section in Lemma 5.3 and $\tilde{f}_i^{\mathrm{ord}} := M^*(v_i, \chi_i) f_i^{\mathrm{ord}} \otimes \omega_i^{-1} \in$

$\mathcal{B}(v_i^{-1}, \chi_i^{-1})_0^{\text{ord}}(v_i^{-1})$. First consider the case where π_1 is the principal series $\chi_1 \boxplus v_1$. Let $(f_1^{\text{ord}})^0$ be the holomorphic image of f_1^{ord} in $\mathcal{B}(v_1, \chi_1)^0$. In view of (3.21), we may take

$$(5.8) \quad \begin{aligned} \phi_p &:= (f_1^{\text{ord}})^0 \otimes W_2^{\text{ord}} \otimes W_3^{\text{ord}}, & \tilde{\phi}_p &:= \tilde{f}_1^{\text{ord}} \otimes \tilde{W}_2^{\text{ord}} \otimes \tilde{W}_3^{\text{ord}}, \\ \phi_p^* &:= (f_1^{\text{ord}})^0 \otimes W_2^{\text{ord}} \otimes \theta_p^{\mathbb{k}} W_3^{\text{ord}}, & \tilde{\phi}_p^* &:= \tilde{f}_1^{\text{ord}} \otimes \tilde{W}_2^{\text{ord}} \otimes \theta_p^{\mathbb{k}} \tilde{W}_3^{\text{ord}}, \end{aligned}$$

where \mathbb{k} is the Dirichlet character defined in (3.12) and $\theta_p^{\mathbb{k}}$ is the twisting operator in (2.12). According to the definition (3.24) and Corollary 5.2, we find that

$$(5.9) \quad \begin{aligned} I_p^{\text{ord}}(\phi_p^* \otimes \tilde{\phi}_p^*, \mathbf{t}_n) &= \frac{L(1, \Pi_p, \text{Ad})}{\zeta_p(2)^2 L(1/2, \Pi_p)} \cdot \frac{\mathcal{I}_p(\rho(t_n) W_2^{\text{ord}} \otimes \theta_p^{\mathbb{k}} W_3^{\text{ord}} \otimes f_1^{\text{ord}}, \rho(t_n) \tilde{W}_2^{\text{ord}} \otimes \theta_p^{\mathbb{k}} \tilde{W}_3^{\text{ord}} \otimes \tilde{f}_1^{\text{ord}})}{\langle \rho(t_n) W_2^{\text{ord}}, \tilde{W}_2^{\text{ord}} \rangle \langle \rho(t_n) W_3^{\text{ord}}, \tilde{W}_3^{\text{ord}} \rangle \langle \rho(t_n) f_1, \tilde{f}_1^{\text{ord}} \rangle} \\ &= \frac{I_p^*}{B_{\Pi_p^{\text{ord}}}^{[n]}} \cdot \frac{\langle \rho(t_n) W_1^{\text{ord}}, \tilde{W}_1^{\text{ord}} \rangle}{\langle \rho(t_n) f_1^{\text{ord}}, \tilde{f}_1^{\text{ord}} \rangle} \cdot \frac{\zeta_p(2)^3}{\zeta_p(1)^3} \cdot \frac{1}{L(1/2, \Pi_p)}, \end{aligned}$$

where

$$I_p^* = \frac{\zeta_p(1) v_1(-1) \gamma(1/2, \pi_2 \otimes \pi_3 \otimes v_1)}{\zeta_p(2)^2} \cdot \Psi(W_2^{\text{ord}}, \theta_p^{\mathbb{k}} W_3^{\text{ord}}, \rho(t_n) f_1^{\text{ord}})^2.$$

Note that the adelization $\mathbb{k}_{\mathbf{A}} = \check{\omega}_F^{-1} \omega_F^{1/2}$; hence

$$\mathbb{k}|_{\mathbf{Z}_p^\times} = \beta_p^{-1} \omega_F^{1/2}|_{\mathbf{Z}_p^\times} = v_1^{-1}|_{\mathbf{Z}_p^\times},$$

and a simple calculation shows that $\theta_p^{\mathbb{k}} W_3^{\text{ord}} \left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \right) = v_1^{-1}(a) \mathbb{I}_{\mathbf{Z}_p^\times}(a)$. We proceed to calculate the local Rankin-Selberg integral

$$\begin{aligned} &\Psi(W_2^{\text{ord}}, \theta_p^{\mathbb{k}} W_3^{\text{ord}}, \rho(t_n) f_1^{\text{ord}}) \\ &= \frac{\zeta_p(2)}{\zeta_p(1)} \int_{\mathbf{Q}_p^\times} \int_{\mathbf{Q}_p} W_2^{\text{ord}} \left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix} \right) \theta_p^{\mathbb{k}} W_3^{\text{ord}} \left(\begin{pmatrix} -y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix} \right) v_1 | \cdot |^{\frac{1}{2}}(y) f_1^{\text{ord}} \left(\begin{pmatrix} 0 & p^{-n} \\ -p^n & 0 \end{pmatrix} w \begin{pmatrix} 1 & p^{-2n} x \\ 0 & 1 \end{pmatrix} \right) dx \frac{d^\times y}{|y|} \\ &= \frac{\zeta_p(2) \chi_1 v_1^{-1} | \cdot | (-p^n)}{\zeta_p(1)} \int_{\mathbf{Q}_p^\times} W_2^{\text{ord}} \left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \right) \theta_p^{\mathbb{k}} W_3^{\text{ord}} \left(\begin{pmatrix} -y & 0 \\ 0 & 1 \end{pmatrix} \right) v_1 | \cdot |^{-\frac{1}{2}}(y) d^\times y \\ &= \frac{\zeta_p(2) \chi_1 v_1^{-1} | \cdot | (-p^n)}{\zeta_p(1)} \int_{\mathbf{Z}_p^\times} W_2^{\text{ord}} \left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \right) d^\times y = \frac{\zeta_p(2) \chi_1 v_1^{-1} | \cdot | (-p^n)}{\zeta_p(1)}. \end{aligned}$$

We thus obtain

$$(5.10) \quad \begin{aligned} I_p^* &= \frac{\zeta_p(1) v_1(-1)}{\zeta_p(2)^2} \cdot \frac{\chi_1 v_1^{-1} | \cdot | (p^{2n}) \zeta_p(2)^2 \gamma(1/2, \pi_2 \otimes \pi_3 \otimes v_1)}{\zeta_p(1)^2} \\ &= \frac{\chi_1 v_1^{-1} | \cdot | (p^{2n}) v_1(-1)}{\zeta_p(1)} \cdot \frac{\varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes v_1) L(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1)}{L(1/2, \pi_2 \otimes \pi_3 \otimes v_1)}. \end{aligned}$$

Substituting (5.6) and (5.10) to (5.9) and noting that

$$\varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes v_1) \varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1) = 1,$$

we immediately obtain (5.7).

Now we treat the remaining case, i.e. $\pi_1 = \chi_1 | \cdot |^{-\frac{1}{2}}$ St is special, and π_2 and π_3 are principal series. Thus $(\pi_1, \pi_3; \pi_2)$ satisfies (Hb). Consider the realizations

$$\mathcal{V}_{\Pi_p} := \mathcal{W}(\pi_1) \boxtimes \mathcal{W}(\pi_3) \boxtimes \mathcal{B}(v_2, \chi_2); \quad \mathcal{V}_{\tilde{\Pi}_p} := \mathcal{W}(\tilde{\pi}_1) \boxtimes \mathcal{W}(\tilde{\pi}_3) \boxtimes \mathcal{B}(v_2^{-1}, \chi_2^{-1}).$$

By Corollary 5.2, we have

$$(5.11) \quad \begin{aligned} L(1/2, \Pi_p) \cdot I_p^{\text{ord}}(\phi_p^* \otimes \tilde{\phi}_p^*, \mathbf{t}_n) &= \frac{\zeta_p(1) v_2(-1) \gamma(1/2, \pi_1 \otimes \pi_3 \otimes v_2)}{\zeta_p(2)^2 \cdot B_{\Pi_p^{\text{ord}}}^{[n]}} \cdot \Psi(\rho(t_n) W_1^{\text{ord}}, \theta_p^{\mathbb{k}} W_3^{\text{ord}}, f_2^{\text{ord}})^2 \\ &\quad \times \frac{\langle \rho(t_n) W_2^{\text{ord}}, \tilde{W}_2^{\text{ord}} \rangle}{\langle \rho(t_n) f_2^{\text{ord}}, \tilde{f}_2^{\text{ord}} \rangle} \cdot \frac{\zeta_p(2)^3}{\zeta_p(1)^3}. \end{aligned}$$

We calculate the local Rankin-Selberg integral in the right hand side

$$\begin{aligned}
& \Psi(\rho(t_n)W_1^{\text{ord}}, \theta_p^k W_3^{\text{ord}}, f_2^{\text{ord}}) \\
&= \frac{\zeta_p(2)}{\zeta_p(1)} \int_{\mathbf{Q}_p^\times} \int_{\mathbf{Q}_p} W_1^{\text{ord}}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} w \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} t_n\right) \theta_p^k W_3^{\text{ord}}\left(\begin{pmatrix} -y & 0 \\ 0 & 1 \end{pmatrix} w \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}\right) v_2 |\cdot|^{\frac{1}{2}}(y) \mathbb{I}_{\mathbf{Z}_p}(x) dx \frac{d^\times y}{|y|} \\
&= \frac{\zeta_p(2) \chi_1 v_1^{-1} |\cdot| (p^n) v_1 v_2 (-1)}{\zeta_p(1)} \int_{\mathbf{Q}_p^\times} \theta_p^k W_3^{\text{ord}}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} w\right) \chi_1 v_2(y) d^\times y \\
&= \frac{\zeta_p(2) \chi_1 v_1^{-1} |\cdot| (p^n) \chi_1 (-1)}{\zeta_p(1)} \gamma(1/2, \pi_3 \otimes v_1 \chi_2) \int_{\mathbf{Q}_p^\times} \theta_p^k W_3^{\text{ord}}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) v_1 \chi_2(y) d^\times y \\
&= \frac{\zeta_p(2) \chi_1 v_1^{-1} |\cdot| (p^n) \chi_1 (-1)}{\zeta_p(1)} \gamma(1/2, \pi_3 \otimes v_1 \chi_2).
\end{aligned}$$

Substituting the above equation and (5.6) into (5.11) and using the formulae of the local L -factor and ε -factor of $\pi_1 \otimes \pi_3 \otimes v_2$ in [GJ78, Proposition 1.4 (1.4.2)], we find that $L(1/2, \Pi_p) \cdot I_p(\phi_p^* \otimes \tilde{\phi}_p^*)$ equals

$$\begin{aligned}
& \frac{\chi_1 v_1^{-1} |\cdot| (p^{2n}) \zeta_p(2)^2}{B_{\Pi_p^{\text{ord}}}^{[n]} \zeta_p(1)^2} \cdot \frac{\varepsilon(1/2, \pi_1 \otimes \pi_3 \otimes v_2) L(1/2, \pi_1 \otimes \pi_3 \otimes \chi_2)}{L(1/2, \pi_1 \otimes \pi_3 \otimes v_2)} \cdot \varepsilon(1/2, \pi_3 \otimes v_1 \chi_2)^2 \frac{L(1/2, \pi_3 \otimes \chi_1 v_2)^2}{L(1/2, \pi_3 \otimes v_1 \chi_2)^2} \\
&= \frac{\zeta_p(2)^2}{\zeta_p(1)^2} \cdot \frac{\chi_1 v_1^{-1} |\cdot| (p^{2n}) \omega_2 \omega_3 (-1)}{B_{\Pi_p^{\text{ord}}}^{[n]}} \cdot \frac{\varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes v_1) L(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1)}{L(1/2, \pi_2 \otimes \pi_3 \otimes v_1)}.
\end{aligned}$$

This proves (5.7) in the remaining case. \square

Remark 5.5. Replacing $\phi_p^* \otimes \tilde{\phi}_p^*$ with $\phi_p \otimes \tilde{\phi}_p$ in (3.24), we define the improved p -adic zeta integral

$$\mathcal{I}_{\Pi_{\mathbf{Q}, p}}^* := I_p^{\text{ord}}(\phi_p \otimes \tilde{\phi}_p, \mathbf{t}_n) \cdot \frac{B_{\Pi_p^{\text{ord}}}^{[n]}}{\chi_1 v_1^{-1} | \cdot | (-p^{2n})} \cdot \frac{\zeta_p(1)^2}{\zeta_p(2)^2}.$$

If π_1 is principal series, then $v_1 \chi_2 \chi_3 \neq |\cdot|^{-\frac{1}{2}}$, and

$$\mathcal{I}_{\Pi_{\mathbf{Q}, p}}^* = \frac{1}{\varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1)} \cdot \frac{L(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1)}{L(1/2, \pi_2 \otimes \pi_3 \otimes v_1)} \cdot L(1/2, v_1 \chi_2 \chi_3)^2;$$

if π_1 is special and $v_1 \chi_2 \chi_3 = |\cdot|^{-\frac{1}{2}}$, then

$$\mathcal{I}_{\Pi_{\mathbf{Q}, p}}^* = \frac{1}{\varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes v_1)} \cdot \frac{(-1)}{L(1/2, v_1 \chi_2 v_3) L(1/2, v_1 v_2 \chi_3)}.$$

These equations will be used later for the interpolation formula of improved p -adic L -functions. It can be obtained by the same computation in the above proposition. We omit the details.

5.3.3. *The balanced case.* Now suppose that \underline{Q} is in the balanced range $\mathfrak{X}_{\mathcal{R}}^{\text{bal}}$. We shall compute the normalized p -adic zeta integral $\mathcal{I}_{\Pi_p}^{\text{bal}}$ in (4.21). Put

$$u_n = \begin{pmatrix} 1 & p^{-n} \\ 0 & 1 \end{pmatrix} \in \text{SL}_2(\mathbf{Q}_p); \quad \check{\mathbf{t}}_n = (u_n, 1, t_n) \in \text{GL}_2(E_p)$$

for $n \geq \max\{c(\pi_1), c(\pi_2), c(\pi_3), 1\}$. Observe that if $L : \pi_1 \otimes \pi_2 \otimes \pi_3 \rightarrow \mathbf{C}$ is any $\text{GL}_2(\mathbf{Q}_p)$ -invariant trilinear form, then

$$\begin{aligned}
L(\pi_1(u_n) \xi_1, \xi_2, \pi_3(t_n) \xi_3) &= L(\pi_1(u_n) \xi_1, \pi_2(t_n) \xi_2, \xi_3) \\
&= L(\xi_1, \pi_2(u_n) \xi_2, \pi_3(t_n) \xi_3),
\end{aligned}$$

Thus we may assume that

(Hb') either $\pi_3 = \chi_3 \boxplus v_3$ is principal series or each of π_1, π_2 and π_3 is special.

Proposition 5.6 (p -adic zeta integral in the balanced case). *Under the assumption (Hb'), we put*

$$\mathcal{E}_{\text{bal}}(\Pi_{\underline{Q},p}) := \frac{\varepsilon(1/2, v_1 v_2 \chi_3)}{\varepsilon(1/2, \chi_1 \chi_2 v_3)} \cdot \left(\frac{L(1/2, \chi_1 \chi_2 v_3)}{L(1/2, v_1 v_2 \chi_3)} \right)^2 \cdot \frac{L(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3)}{\varepsilon(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3) L(1/2, \pi_1 \otimes \pi_2 \otimes v_3)}.$$

Then we have

$$\mathcal{I}_{\Pi_{\underline{Q},p}}^{\text{bal}} = \mathcal{E}_{\text{bal}}(\Pi_{\underline{Q},p}) \cdot \frac{1}{L(1/2, \Pi_{\underline{Q},p})}.$$

PROOF. We write $\Pi_p = \Pi_{\underline{Q},p}$ as before. By definition, this is equivalent to proving

$$\begin{aligned} I_p^{\text{ord}}(\phi_p \otimes \tilde{\phi}_p, \check{\mathfrak{t}}_n) &= \alpha_p(F)^{2n} \omega_{F,p}^{-1/2}(-p^{2n}) |p^n|^{k_1+k_2+k_3} \cdot \mathcal{E}_{\text{bal}}(\Pi_p) \cdot \frac{\zeta_p(2)^2}{B_{\Pi_p^{\text{ord}}}^{[n]}} \cdot \frac{1}{L(1/2, \Pi_p)} \\ &= \chi_1 \chi_2 \chi_3 (-p^{2n}) |p|^{3n} \cdot \mathcal{E}_{\text{bal}}(\Pi_p) \cdot \frac{\zeta_p(2)^2}{B_{\Pi_p^{\text{ord}}}^{[n]}} \cdot \frac{1}{L(1/2, \Pi_p)}, \end{aligned}$$

where $I_p^{\text{ord}}(\phi_p \otimes \tilde{\phi}_p, \check{\mathfrak{t}}_n)$ is the local zeta integral in (4.16). The assumption (Hb') implies that $(\pi_1, \pi_2; \pi_3)$ satisfies (Hb), so we consider the realizations

$$\mathcal{V}_{\Pi_p} = \mathcal{W}(\pi_1) \boxtimes \mathcal{W}(\pi_2) \boxtimes \mathcal{B}(v_3, \chi_3)^0; \quad \mathcal{V}_{\tilde{\Pi}_p} = \mathcal{W}(\tilde{\pi}_1) \boxtimes \mathcal{W}(\tilde{\pi}_2) \boxtimes \mathcal{B}(v_3^{-1}, \chi_3^{-1})_0.$$

Let $W_i^{\text{ord}} = W_{\pi_i}^{\text{ord}}$ and $\tilde{W}_i^{\text{ord}} = W_i^{\text{ord}} \otimes \omega_i^{-1}$ be the normalized ordinary Whittaker functions for $i = 1, 2$. Let f_3^{ord} be the normalized ordinary section in $\mathcal{B}(v_3, \chi_3)^{\text{ord}}(\chi_3)$ in Lemma 5.3 and let $\tilde{f}_3^{\text{ord}} := M^*(v_3, \chi_3) f_3^{\text{ord}} \otimes \omega_3^{-1}$. Letting $(f_3^{\text{ord}})^0$ be the holomorphic image of f_3^{ord} in $\mathcal{B}(v_3, \chi_3)^0$ as before, we may take

$$\phi_p = W_1^{\text{ord}} \otimes W_2^{\text{ord}} \otimes (f_3^{\text{ord}})^0; \quad \tilde{\phi}_p = \tilde{W}_1^{\text{ord}} \otimes \tilde{W}_2^{\text{ord}} \otimes \tilde{f}_3^{\text{ord}}.$$

From the definition (4.16), Corollary 5.2 and (5.6), we deduce that

$$\begin{aligned} (5.12) \quad & I_p^{\text{ord}}(\phi_p \otimes \tilde{\phi}_p, \check{\mathfrak{t}}_n) \\ &= \frac{\mathcal{I}_p(\rho(u_n) W_1^{\text{ord}} \otimes W_2^{\text{ord}} \otimes \rho(t_n) f_3^{\text{ord}}, \rho(u_n) \tilde{W}_1^{\text{ord}} \otimes \tilde{W}_2^{\text{ord}} \otimes \rho(t_n) \tilde{f}_3^{\text{ord}})}{\zeta_p(2)^2 L(1/2, \Pi_p) \cdot B_{\Pi_p^{\text{ord}}}^{[n]}} \cdot \frac{\langle \rho(t_n) W_3^{\text{ord}}, \tilde{W}_3^{\text{ord}} \rangle}{\langle \rho(t_n) f_3^{\text{ord}}, \tilde{f}_3^{\text{ord}} \rangle} \cdot \frac{\zeta_p(2)^3}{\zeta_p(1)^3} \\ &= \frac{I_p^*}{B_{\Pi_p^{\text{ord}}}^{[n]}} \cdot \frac{\chi_3(-1) \zeta_p(1)^2}{\zeta_p(2)} \cdot \frac{\zeta_p(2)^3}{\zeta_p(1)^3} \cdot \frac{1}{L(1/2, \Pi_p)}, \end{aligned}$$

where

$$I_p^* = \frac{\zeta_p(1) v_3(-1) \gamma(1/2, \pi_1 \otimes \pi_2 \otimes v_3)}{\zeta_p(2)^2} \cdot \Psi(\rho(u_n) W_1^{\text{ord}}, W_2^{\text{ord}}, \rho(t_n) f_3^{\text{ord}})^2.$$

The local Rankin-Selberg integral $\Psi(\rho(u_n) W_1^{\text{ord}}, W_2^{\text{ord}}, \rho(t_n) f_3^{\text{ord}})$ equals

$$\begin{aligned} & \int_{ZN \setminus G} W_1^{\text{ord}}(g \begin{pmatrix} 1 & p^{-n} \\ 0 & 1 \end{pmatrix}) W_2^{\text{ord}}(\mathcal{J}g) f_3^{\text{ord}}(g \begin{pmatrix} 0 & p^{-n} \\ -p^n & 0 \end{pmatrix}) dg \\ &= \frac{\zeta_p(2)}{\zeta_p(1)} \int_{\mathbf{Q}_p^\times} \int_{\mathbf{Q}_p} W_1^{\text{ord}} \left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & p^{-n} \\ x & 1 + xp^{-n} \end{pmatrix} \right) W_2^{\text{ord}} \left(\begin{pmatrix} -y & 0 \\ x & 1 \end{pmatrix} \right) f_3^{\text{ord}} \left(\begin{pmatrix} p^{-n}y & 0 \\ 0 & p^n \end{pmatrix} w \begin{pmatrix} 1 & -p^{-2n}x \\ 0 & 1 \end{pmatrix} \right) \frac{d^\times y}{|y|} dx \\ &= \frac{\zeta_p(2) |p^{2n}| \chi_3 v_3^{-1}(p^n) \cdot |\cdot|^{\frac{1}{2}}(p^{-2n})}{\zeta_p(1)} \int_{\mathbf{Q}_p^\times} \int_{\mathbf{Z}_p} W_1^{\text{ord}} \left(\begin{pmatrix} y & yp^{-n}(1 + xp^n)^{-1} \\ 0 & 1 \end{pmatrix} \right) W_2^{\text{ord}} \left(\begin{pmatrix} -y & 0 \\ 0 & 1 \end{pmatrix} \right) v_3 \cdot |\cdot|^{-\frac{1}{2}}(y) dx d^\times y \\ &= \frac{\zeta_p(2) |p^n| \chi_3 v_3^{-1}(p^n) \chi_2(-1)}{\zeta_p(1)} \int_{\mathbf{Q}_p^\times} \psi(yp^{-n}) \mathbb{I}_{\mathbf{Z}_p}(y) \chi_1 \chi_2 v_3 \cdot |\cdot|^{\frac{1}{2}}(y) d^\times y \\ &= \frac{\zeta_p(2) |p^n| \chi_3 v_3^{-1}(p^n) \chi_2 v_3(-1)}{\zeta_p(1)} \cdot \frac{L(1/2, \chi_1 \chi_2 v_3)}{\varepsilon(1/2, \chi_1 \chi_2 v_3) L(1/2, \chi_1^{-1} \chi_2^{-1} v_3^{-1})} \int_{\mathbf{Q}_p^\times} \mathbb{I}_{p^{-n}(-1+p^n \mathbf{Z}_p)}(y) \chi_1^{-1} \chi_2^{-1} v_3^{-1} \cdot |\cdot|^{\frac{1}{2}}(y) d^\times y \\ &= \frac{\zeta_p(2) |p^n| \chi_1 \chi_2 \chi_3 (p^n) \chi_2 v_3(-1)}{\zeta_p(1)} \cdot \gamma(1/2, \chi_1^{-1} \chi_2^{-1} v_3^{-1}) \cdot \frac{|p|^{\frac{n}{2}}}{1-|p|}, \end{aligned}$$

so we find that

$$\begin{aligned} I_p^* &= v_3(-1)\varepsilon(1/2, \pi_1 \otimes \pi_2 \otimes v_3) \frac{L(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3)}{L(1/2, \pi_1 \otimes \pi_2 \otimes v_3)} \left(\zeta_p(2) |p|^{\frac{3n}{2}} \chi_1 \chi_2 \chi_3(p^n) \cdot \gamma(1/2, \chi_1^{-1} \chi_2^{-1} v_3^{-1}) \right)^2 \\ &= v_3(-1) \zeta_p(2)^2 \cdot \chi_1 \chi_2 \chi_3(p^{2n}) |p|^{3n} \cdot \chi_1 \chi_2 v_3(-1) \mathcal{E}_{\text{bal}}(\Pi_p). \end{aligned}$$

Substituting the above equation to (5.12), we obtain the desired formula. \square

Remark 5.7. Keep the notation in §1.3. For $\bullet \in \{\mathbf{f}, \text{bal}\}$, we put $U_{\underline{Q}} := \text{WD}_p(\text{Fil}_{\bullet}^+ \mathbf{V}_{\underline{Q}}^\dagger) \otimes_{\overline{\mathbf{Q}}_p, \iota_p} \mathbf{C}$ be the Weil-Deligne representation of $W_{\mathbf{Q}_p}$ associated with $\text{Fil}_{\bullet}^+ \mathbf{V}_{\underline{Q}}^\dagger$ by Fontaine [Fon94, (4.2.3)]. It is an easy exercise to show that

$$\mathcal{E}_{\bullet}(\Pi_{\underline{Q}, p}) = \frac{L(0, U_{\underline{Q}})}{\varepsilon(U_{\underline{Q}}) L(1, U_{\underline{Q}}^\vee)},$$

and hence

$$\mathcal{I}_{\Pi_{\underline{Q}, p}}^{\bullet} = \mathcal{E}_p(\text{Fil}_{\bullet}^+ \mathbf{V}_{\underline{Q}}^\dagger).$$

For example, if $\bullet = \text{bal}$ and $\pi_i = \chi_i |\cdot|^{-\frac{1}{2}} \text{St}$ are special for $i = 1, 2, 3$, then $\dim U_{\underline{Q}}^{\mathbf{N}=0} = 3$, where \mathbf{N} is the monodromy operator, and we have $L(s - \frac{1}{2}, U_{\underline{Q}}) = L(s, \chi_1 \chi_2 \chi_3) L(s, \chi_1 \chi_2 v_3)^2$ and $L(s - \frac{1}{2}, U_{\underline{Q}}^\vee) = L(s, v_1 v_2 \chi_3)^3$.

6. THE CALCULATION OF LOCAL ZETA INTEGRALS (II)

6.1. Setting. We continue to let $F = (f, g, h)$ be the specialization of $\mathbf{F} = (\mathbf{f}, \mathbf{g}, \mathbf{h})$ at a classical point $\underline{Q} = (Q_1, Q_2, Q_3)$. In this section, we assume the following hypothesis for the unitary automorphic representations (π_f, π_g, π_h) attached to (f, g, h)

Hypothesis 6.1. For each prime $q \mid N$, there exists a rearrangement $\{f_1, f_2, f_3\}$ of $\{f, g, h\}$ such that

- (1) $c_q(\pi_{f_1}) \leq \min\{c_q(\pi_{f_2}), c_q(\pi_{f_3})\}$,
- (2) the local components $\pi_{f_1, q}$ and $\pi_{f_3, q}$ are minimal,
- (3) either $\pi_{f_3, q}$ is a principal series or $\pi_{f_2, q}$ and $\pi_{f_3, q}$ are both discrete series.

Recall that an irreducible admissible representation π of $\text{GL}_2(\mathbf{Q}_q)$ is minimal if the conductor $c(\pi)$ is minimal among the twists $\pi \otimes \chi$ for all characters $\chi : \mathbf{Q}_q^\times \rightarrow \mathbf{C}^\times$.

Remark 6.2. Note that if the above hypothesis holds for (f, g, h) , then it also holds for specializations of $(\mathbf{f}, \mathbf{g}, \mathbf{h})$ at any classical point by Remark 3.1. Moreover, we observe that one can always find Dirichlet characters χ_1, χ_2 and χ_3 modulo some M with $M^2 \mid N$ such that $\chi_1 \chi_2 \chi_3 = 1$ and $(\pi_f \otimes \chi_1, \pi_g \otimes \chi_2, \pi_h \otimes \chi_3)$ satisfies Hypothesis 6.1.

As before, we let $\pi_1 = \pi_{f, q} \otimes \omega_{F, q}^{-1/2}$, $\pi_2 = \pi_{g, q}$ and $\pi_3 = \pi_{h, q}$; let $\Pi_q = \Pi_{\underline{Q}, q} = \pi_1 \times \pi_2 \times \pi_3$. Let q be a prime factor of N . Suppose that

$$\varepsilon(1/2, \Pi_q) = +1 \quad (q \notin \Sigma^-).$$

The purpose of this section is to evaluate the local zeta integral defined in (3.23)

$$I_q(\phi_q^* \otimes \tilde{\phi}_q^*) = \frac{L(1, \Pi_q, \text{Ad})}{\zeta_q(2)^2 L(1/2, \Pi_q)} \int_{\text{PGL}_2(\mathbf{Q}_q)} \frac{\mathbf{b}_q(\Pi_q(g_q) \phi_q^*, \tilde{\phi}_q^*)}{\mathbf{b}_q(\Pi_q(\tau_{\underline{N}, q}) \phi_q, \tilde{\phi}_q)} dg_q$$

under Hypothesis 6.1. For $i = 1, 2, 3$, let $c_i = c(\pi_i)$ be the exponent of the conductors. Note that $\omega_{F, q}^{1/2}$ is unramified, so under Hypothesis 6.1 and the condition (sf), we may assume by symmetry that

$$c_1 \leq \min\{c_2, c_3, 1\}; \quad \pi_3 \text{ is minimal,}$$

and that $\{\pi_1, \pi_2, \pi_3\}$ satisfies one of the following conditions:

- Case (Ia): $\pi_3 = \chi_3 \boxplus v_3$ is a principal series with χ_3 unramified character of \mathbf{Q}_q^\times .
- Case (Ib): π_1, π_2 and π_3 are discrete series.
- Case (IIa): π_1 is a principal series; π_2 and π_3 are discrete series with $L(s, \pi_2 \otimes \pi_3) \neq 1$.
- Case (IIb): π_1 is a principal series; π_2 and π_3 are discrete series with $L(s, \pi_2 \otimes \pi_3) = 1$.

For $i = 1, 2, 3$, let $\xi_i \in \mathcal{V}_{\pi_i}^{\text{new}}$ and $\tilde{\xi}_i \in \tilde{\pi}_i(\tau_{c_i})\mathcal{V}_{\tilde{\pi}_i}^{\text{new}}$ be new vectors. Set

$$c^* = \max\{c_2, c_3\} > 0.$$

We recall the following choices of local test vectors $\phi_q^* \in \mathcal{V}_{\Pi_q}$ and $\tilde{\phi}_q^* \in \mathcal{V}_{\tilde{\Pi}_q}$ in (3.21) and (3.22) according to the polynomials $\mathcal{Q}_{i,q}(X)$ for $i = 1, 2, 3$ in (3.20). Put

$$w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \eta = \begin{pmatrix} q^{-1} & 0 \\ 0 & 1 \end{pmatrix} \text{ and } \tau_n = \begin{pmatrix} 0 & 1 \\ -q^n & 0 \end{pmatrix} \text{ for } n \in \mathbf{Z}.$$

- Case (Ia) and (Ib):

$$\begin{aligned} \phi_q^* &= \xi_1 \otimes \pi_2(\eta^{c^*-c_2})\xi_2 \otimes \pi_3(\eta^{c^*-c_3})\xi_3, \\ \tilde{\phi}_q^* &= \omega_2(q^{c_2-c^*})\omega_3(q^{c_3-c^*}) \cdot \tilde{\xi}_1 \otimes \tilde{\pi}_2(\eta^{c^*-c_2})\tilde{\xi}_2 \otimes \tilde{\pi}_3(\eta^{c^*-c_3})\tilde{\xi}_3. \end{aligned}$$

- Case (IIa): Let $r = \lceil \frac{c^*}{2} \rceil$. Then

$$\phi_q^* = \pi_1(\eta^r)\xi_1 \otimes \xi_2 \otimes \xi_3, \quad \tilde{\phi}_q^* = \omega_1(q^{-r}) \cdot \tilde{\pi}_1(\eta^r)\tilde{\xi}_1 \otimes \tilde{\xi}_2 \otimes \tilde{\xi}_3.$$

- Case (IIb): If $c_1 = 0$, then let $v_1 : \mathbf{Q}_q^\times \rightarrow \mathbf{C}^\times$ be the unramified character with $v_1(q) = \beta_q(f) |q|^{\frac{k_1-1}{2}}$, where $\beta_q(f)$ is the specialization of $\beta_q(\mathbf{f})$ at Q_1 in Definition 3.3 and we have

$$\begin{aligned} \phi_q^* &= (\pi_1(\eta^{c^*})\xi_1 - v_1^{-1}|\cdot|^{\frac{1}{2}}(q)\pi_1(\eta^{c^*-1})\xi_1) \otimes \xi_2 \otimes \xi_3, \\ \tilde{\phi}_q^* &= \omega_1(q^{-c^*}) \cdot (\tilde{\pi}_1(\eta^{c^*})\tilde{\xi}_1 - \omega_1 v_1^{-1}|\cdot|^{\frac{1}{2}}(q)\tilde{\pi}_1(\eta^{c^*-1})\tilde{\xi}_1) \otimes \tilde{\xi}_2 \otimes \tilde{\xi}_3. \end{aligned}$$

If $c_1 > 0$, then

$$\phi_q^* = \pi_1(\eta^{c^*-c_1})\xi_1 \otimes \xi_2 \otimes \xi_3, \quad \tilde{\phi}_q^* = \omega_1(q^{c_1-c^*}) \cdot \tilde{\pi}_1(\eta^{c^*-c_1})\tilde{\xi}_1 \otimes \tilde{\xi}_2 \otimes \tilde{\xi}_3.$$

In what follows, we let $W_i = W_{\pi_i} \in \mathcal{W}(\pi_i)^{\text{new}}$ be the normalized Whittaker newforms and let $\tilde{W}_i = W_{\tilde{\pi}_i} \otimes \omega_i^{-1}$ for $i = 1, 2, 3$. For a non-negative integer n , put

$$\mathcal{U}_0(q^n) = \text{GL}_2(\mathbf{Z}_q) \cap \begin{pmatrix} \mathbf{Z}_q & \mathbf{Z}_q \\ q^n \mathbf{Z}_q & \mathbf{Z}_q \end{pmatrix}.$$

6.2. The ramified case (Ia). In the case (Ia), $\pi_3 = \chi_3 \boxplus v_3$ is a principle series with $c(\chi_3) = 0$.

Proposition 6.3. *In case (Ia), we have*

$$I_q(\phi_q^* \otimes \tilde{\phi}_q^*) = \varepsilon(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3) \cdot \chi_3^{-2} |\cdot| (q^{c^*}) \omega_3(-1) \varepsilon(1/2, \pi_3)^2 \cdot \frac{1}{B_{\Pi_q}} \cdot \frac{\zeta_q(2)^2}{\zeta_q(1)^2}.$$

PROOF. In this case, $c_3 = c(\omega_3) = c(\omega_1 \omega_2) \leq c_2$, so $c^* = c_2$. We use the realizations

$$\mathcal{V}_{\Pi_q} = \mathcal{W}(\pi_1) \boxtimes \mathcal{W}(\pi_2) \boxtimes \mathcal{B}(\chi_3, v_3); \quad \mathcal{V}_{\tilde{\Pi}_q} = \mathcal{W}(\tilde{\pi}_1) \boxtimes \mathcal{W}(\tilde{\pi}_2) \boxtimes \mathcal{B}(\chi_3^{-1}, v_3^{-1}).$$

Let $f_3 \in \mathcal{B}(\chi_3, v_3)^{\text{new}}$ be the new section normalized so that $f_3(1) = 1$ and $\tilde{f}_3 = M^*(\chi_3, v_3)f_3 \otimes \omega_3^{-1}$. Let

$$f_3^* = \rho \left(\begin{pmatrix} q^{c_3-c_2} & 0 \\ 0 & 1 \end{pmatrix} \right) f_3; \quad \tilde{f}_3^* = \omega_3(q^{c_3-c_2}) \cdot \rho \left(\begin{pmatrix} q^{c_3-c_2} & 0 \\ 0 & 1 \end{pmatrix} \right) \tilde{f}_3 = M^*(\chi_3, v_3)f_3^* \otimes \omega_3^{-1}.$$

We thus have

$$\phi_q^* = W_1 \otimes W_2 \otimes f_3^*; \quad \tilde{\phi}_q^* = \tilde{W}_1 \otimes \tilde{W}_2 \otimes \tilde{f}_3^*.$$

By Corollary 5.2,

$$\begin{aligned} (6.1) \quad I_q(\phi_q^* \otimes \tilde{\phi}_q^*) &= \frac{\mathcal{I}_q(W_1 \otimes W_2 \otimes f_3^*, \tilde{W}_1 \otimes \tilde{W}_2 \otimes \tilde{f}_3^*)}{\zeta_q(2)^2 L(1/2, \Pi_q) \cdot B_{\Pi_q}} \cdot \frac{\langle \rho(\tau_{c_3})W_3, \tilde{W}_3 \rangle}{\langle \rho(\tau_{c_3})f_3, \tilde{f}_3 \rangle} \cdot \frac{\zeta_q(2)^3}{\zeta_q(1)^3} \\ &= \frac{I_q^*}{B_{\Pi_q}} \cdot \frac{\langle \rho(\tau_{c_3})W_3, \tilde{W}_3 \rangle}{\langle \rho(\tau_{c_3})f_3, \tilde{f}_3 \rangle} \cdot \frac{\zeta_q(2)^3}{\zeta_q(1)^3}, \end{aligned}$$

where

$$I_q^* = \frac{\zeta_q(1) \cdot \chi_3(-1) \gamma(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3)}{\zeta_q(2)^2 L(1/2, \Pi_q)} \cdot \Psi(W_1, W_2, f_3)^2$$

There are three subcases:

- (a) v_3 is ramified,
- (b) v_3 is unramified and $L(s, \pi_2) = L(s, \chi_2)$ for some unramified character χ_2 ,
- (c) v_3 is unramified and $L(s, \pi_2) = 1$.

Subcase (a): In this case, $f_3 \in \mathcal{B}(\chi_3, v_3)$ is given by

$$f_3\left(\begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}\right) = \mathbb{I}_{q^{c_3} \mathbf{Z}_q}(x)$$

by [Sch02, Prop. 2.1.2]. We have $W_2\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) = \mathbb{I}_{\mathbf{Z}_q^\times}(y)$ if $L(s, \pi_2) = 1$ and $W_2\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) = \chi_2|\cdot|^{\frac{1}{2}}(y)\mathbb{I}_{\mathbf{Z}_q}(y)$ if $L(s, \pi_2) = L(s, \chi_2)$ for some unramified character χ_2 . In any case, the integral $\Psi(W_1, W_2, f_3^*)$ equals

$$\begin{aligned} & \frac{\zeta_q(2)\chi_3|\cdot|^{\frac{1}{2}}(q^{c_3-c_2})}{\zeta_q(1)} \int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q} W_1\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}\right) W_2\left(\begin{pmatrix} -y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}\right) \chi_3|\cdot|^{-\frac{1}{2}}(y) f_3\left(\begin{pmatrix} 1 & 0 \\ q^{c_3-c_2}x & 1 \end{pmatrix}\right) dx d^\times y \\ &= \frac{\zeta_q(2)\chi_3|\cdot|^{\frac{1}{2}}(q^{c_3-c_2})}{\zeta_q(1)} |q|^{c_2} \int_{\mathbf{Q}_q^\times} W_1\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) W_2\left(\begin{pmatrix} -y & 0 \\ 0 & 1 \end{pmatrix}\right) \chi_3|\cdot|^{-\frac{1}{2}}(y) d^\times y. \end{aligned}$$

Note that π_1 and π_2 can not be both unramified special representations as $c(\pi_1) \leq 1$ and v_3 is ramified. A standard calculation together with the recipe of local L -factors for $\mathrm{GL}(2) \times \mathrm{GL}(2)$ in [GJ78, Proposition 1.4] shows that

$$\int_{\mathbf{Q}_q^\times} W_1\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) W_2\left(\begin{pmatrix} -y & 0 \\ 0 & 1 \end{pmatrix}\right) \chi_3|\cdot|^{-\frac{1}{2}}(y) d^\times y = L(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3).$$

We obtain

$$\Psi(W_1, W_2, f_3^*) = \frac{\zeta_q(2)\chi_3|\cdot|^{\frac{1}{2}}(q^{c_3-c_2})}{\zeta_q(1)} |q|^{c_2} L(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3),$$

and hence

$$I_q^* = \chi_3^{-2}|\cdot|(q^{c_2}) \cdot \chi_3^2|\cdot|(q^{c_3}) \cdot \varepsilon(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3).$$

Substituting the above equation and the formula Lemma 6.4 below to (6.1), we obtain the expression of $I_q(\phi_q^* \otimes \tilde{\phi}_q^*)$ as claimed in this subcase.

Subcase (b) and (c): Next we consider the case v_3 is unramified, so π_1 and π_3 are spherical ($c_1 = c_3 = 0$). Note that in Subcase (b) where $L(s, \pi_2) = L(s, \chi_2)$ for χ_2 a unramified character, we must have $\pi_2 = \chi_2|\cdot|^{-\frac{1}{2}} \mathrm{St}$ is a unramified special representation. Define the function $\mathcal{F} : ZN \backslash G/K_0(q^{c_2}) \rightarrow \mathbf{C}$ by

$$\mathcal{F}(g) = W_1(g) W_2\left(\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} g\right) f_3\left(g \begin{pmatrix} q^{-c_2} & 0 \\ 0 & 1 \end{pmatrix}\right).$$

We have

$$\begin{aligned} \Psi(W_1, W_2, f_3^*) &= \frac{\zeta_q(2)}{\zeta_q(1)} \int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q} \mathcal{F}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}\right) dx \frac{d^\times y}{|y|} \\ &= \frac{\zeta_q(2)}{\zeta_q(1)} \cdot (J_0^- + J_{c_2}^+ + \sum_{n=1}^{c_2-1} J_n), \end{aligned}$$

where

$$\begin{aligned} J_0^- &= \int_{\mathbf{Q}_q^\times} \int_{|x| \geq 1} \mathcal{F}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}\right) dx \frac{d^\times y}{|y|}, \\ J_n &= \int_{\mathbf{Q}_q^\times} \int_{q^n \mathbf{Z}_q^\times} \mathcal{F}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}\right) dx \frac{d^\times y}{|y|}, \\ J_{c_2}^+ &= \int_{\mathbf{Q}_q^\times} \int_{|x| \leq |q|^{c_2}} \mathcal{F}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}\right) dx \frac{d^\times y}{|y|}. \end{aligned}$$

Using the identity

$$\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix} = (-x) \cdot \begin{pmatrix} yx^{-2} & x^{-1} \\ 0 & 1 \end{pmatrix} w \begin{pmatrix} 1 & x^{-1} \\ 0 & 1 \end{pmatrix}$$

and the formula

$$W_2\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} w\right) = \begin{cases} -|\varpi|\chi_2|\cdot|^{\frac{1}{2}}(y)\mathbb{I}_{\mathbf{Z}_q}(y) & \text{in subcase (b),} \\ \varepsilon(1/2, \pi_2) \cdot \omega_2(q^{-c_2})\mathbb{I}_{q^{-c_2}\mathbf{Z}_q^\times}(y) & \text{in subcase (c),} \end{cases}$$

we find that

$$\begin{aligned} J_0^- &= \int_{\mathbf{Q}_q^\times} \mathcal{F}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} w\right) \frac{d^\times y}{|y|} \\ &= v_3^{-1}|\cdot|^{\frac{1}{2}}(q^{c_2}) \int_{\mathbf{Q}_q^\times} W_1\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) W_2\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} w\right) \chi_3|\cdot|^{-\frac{1}{2}}(y) d^\times y \\ &= -|q| \cdot v_3^{-1}|\cdot|^{\frac{1}{2}}(q^{c_2}) \cdot \begin{cases} L(1/2, \pi_1 \otimes \chi_2 \chi_3) & \text{in subcase (b),} \\ 0 & \text{in subcase (c).} \end{cases} \end{aligned}$$

On the other hand, it is easy to see that

$$J_{c_2}^+ = |q|^{c_2} \int_{\mathbf{Q}_q^\times} \mathcal{F}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) \frac{d^\times y}{|y|} = \chi_3^{-1}|\cdot|^{\frac{1}{2}}(q^{c_2})L(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3).$$

It remains to calculate J_n in subcase (c). We have

$$J_n = \sum_{m \in \mathbf{Z}} (1 - |q|) |q|^n \chi_3|\cdot|^{\frac{1}{2}}(q^{-c_2}) W_1\left(\begin{pmatrix} q^m & 0 \\ 0 & 1 \end{pmatrix}\right) \chi_3|\cdot|^{-\frac{1}{2}}(q^m) f_3\left(\begin{pmatrix} 1 & 0 \\ q^{n-c_2} & 1 \end{pmatrix}\right) A_n^{(m)}(\mathbf{1}),$$

where

$$A_n^{(m)}(\mathbf{1}) = \int_{\mathbf{Z}_q^\times} W_2\left(\begin{pmatrix} q^m y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ q^n & 1 \end{pmatrix}\right) d^\times y.$$

By Lemma 6.5 below, we find that $J_n = 0$ unless $n = c_2 - 1$ and

$$J_{c_2-1} = \chi_3^{-1}|\cdot|^{\frac{1}{2}}(q^{c_2}) \cdot \chi_3 v_3^{-1}|\cdot|(q).$$

Combining the above calculations, in either subcase (b) or subcase (c), we obtain

$$\Psi(W_1, W_2, f_3) = \frac{\zeta_q(2)}{\zeta_q(1)} \cdot (J_0^- + J_{c_2}^+ + \sum_{n=1}^{c_2-1} J_n) = \frac{\zeta_q(2)\chi_3^{-1}|\cdot|^{\frac{1}{2}}(q^{c_2})}{\zeta_q(1)} \cdot \frac{L(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3)}{L(1, \chi_3 v_3^{-1})}.$$

This shows that

$$I_q^* = \frac{\zeta_q(1)\gamma(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3)\Psi(W_1, W_2, f_3^*)^2}{\zeta_q(2)^2 L(1/2, \Pi_q)} = \frac{\chi_3^{-2}|\cdot|(q^{c_2})\varepsilon(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3)}{\zeta_q(1)L(1, \chi_3 v_3^{-1})^2}.$$

The above equation with Lemma 6.4 below and (6.1) yield that

$$I_q(\phi_q^* \otimes \tilde{\phi}_q^*) = \chi_3^{-2}|\cdot|(q^{c_2})\varepsilon(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3) \cdot \frac{1}{B_{\Pi_q}} \cdot \frac{\zeta_q(2)^2}{\zeta_q(1)^2}.$$

This completes the proof. \square

Lemma 6.4. *Let π be a constituent of $\mathcal{B}(\chi, v)$ of central character ω . Suppose that χ is unramified. Let $c = c(\pi)$ be the exponent of the conductor. Let W_π be the new vector in $\mathcal{W}(\pi)^{\text{new}}$ with $W_\pi(1) = 1$ and $\widetilde{W}_\pi = W_\pi \otimes \omega^{-1}$. Let $f \in \mathcal{B}(v, \chi)$ and $\tilde{f} = M^*(\chi, v)f \otimes \omega^{-1}$.*

(1) *Suppose that π is a principal series and $f \in \mathcal{B}(\chi, v)^{\text{new}}$ is the new section with $f(1) = 1$. Then*

$$\frac{\langle \rho(\tau_c)W_\pi, \widetilde{W}_\pi \rangle}{\langle \rho(\tau_c)f, \tilde{f} \rangle} = \chi^2|\cdot|(q^{-c})\varepsilon(1/2, \pi)^2\omega(-1) \cdot L(1, \chi v^{-1})^2 \cdot \frac{\zeta_q(1)^2}{\zeta_q(2)}.$$

(2) *Suppose that π is a unramified special representation with $\chi v^{-1} = |\cdot|^{-1}$, i.e. $\pi = v|\cdot|^{-\frac{1}{2}}\text{St}$. Let f be the section in $\mathcal{B}(\chi, v)^{U_0(q)}$ with $f(w) = 1$. Then*

$$\frac{\langle \rho(\tau_c)W_\pi, \widetilde{W}_\pi \rangle}{\langle \rho(\tau_c)f, \tilde{f} \rangle} = \frac{\zeta_q(1)^2}{\zeta_q(2)}.$$

PROOF. We first consider the case π is a principal series. Suppose that $c = 0$. Then we have

$$\begin{aligned} \langle f, M^*(\chi, v)f \otimes \omega^{-1} \rangle &= \gamma(0, \chi v^{-1}) \frac{L(0, \chi v^{-1})}{L(1, \chi v^{-1})} = \frac{L(1, \chi^{-1}v)}{L(1, \chi v^{-1})}, \\ \langle W_\pi, W_\pi \rangle &= \frac{L(1, \pi, \text{Ad})\zeta_q(1)}{\zeta_q(2)}, \end{aligned}$$

and hence

$$\frac{\langle W_\pi, \widetilde{W}_\pi \rangle}{\langle f, \widetilde{f} \rangle} = L(1, \chi v^{-1})^2 \cdot \frac{\zeta_q(1)^2}{\zeta_q(2)}.$$

Suppose that $c > 0$. Then v is ramified and f is supported in $B\mathcal{U}_0(q^c)$ (cf. [Sch02, Proposition 2.1.2]), and hence $\langle \rho(\tau_c)f, \widetilde{f} \rangle = \langle f, \rho(\tau_c^{-1})\widetilde{f} \rangle$ equals

$$\begin{aligned} \int_K f(k)\widetilde{f}(k\tau_c^{-1})dk &= \text{vol}(\mathcal{U}_0(q^c)) \cdot \widetilde{f}(\tau_c^{-1}) \\ &= \text{vol}(\mathcal{U}_0(q^c)) \cdot \omega(q^c) \cdot M^*(\chi, v)f(\tau_c^{-1}) \\ &= \text{vol}(\mathcal{U}_0(q^c)) \cdot \omega(q^c) \cdot \gamma(0, \chi v^{-1})f\left(\begin{pmatrix} 1 & 0 \\ 0 & q^{-c} \end{pmatrix}\right) \\ &= |q|^c (1 + |q|)^{-1} \varepsilon(1/2, \chi v^{-1}) \cdot \chi(q^c). \end{aligned}$$

In addition, $\langle \rho(\tau_c)W_\pi, \widetilde{W}_\pi \rangle = \varepsilon(1/2, \pi)\zeta_q(1)$, so we obtain that

$$\frac{\langle \rho(\tau_c)W_\pi, \widetilde{W}_\pi \rangle}{\langle \rho(\tau_c)f, \widetilde{f} \rangle} = \frac{\zeta_q(1)^2}{\zeta_q(2)} \chi^2 \cdot |(q^{-c})\varepsilon(1/2, \pi)^2 \omega(-1).$$

Now we consider the case π is a unramified special representation. Then $c = 1$ and we may assume $f(w) = 1$, i.e. f is supported in $Bw\mathcal{U}_0(q)$. An elementary computation shows that

$$M^*(\chi, v)f(1) = \zeta_q(2)(1 - |q|^{-1}); \quad M^*(\chi, v)f(w) = \zeta_q(2).$$

Then $\langle \rho(\tau_1)f, \widetilde{f} \rangle$ equals

$$\int_{w\mathcal{U}_0(q)} f(k\tau_1)\widetilde{f}(k)dk = \text{vol}(\mathcal{U}_0(q)) \cdot f(\tau_1)M^*(\chi, v)f(1) = (-v|\cdot|^{-\frac{1}{2}}(q)) \cdot \frac{\zeta_q(2)^2}{\zeta_q(1)^2}.$$

Combined with the formulas

$$\langle \rho(\tau_1)W_\pi, \widetilde{W}_\pi \rangle = \varepsilon(1/2, \pi)\zeta_q(2) = (-v|\cdot|^{-\frac{1}{2}}(q)) \cdot \zeta_q(2),$$

the lemma in this case follows. \square

Lemma 6.5. *Let π is an irreducible admissible generic representation of $\text{GL}_2(\mathbf{Q}_q)$ and let $W_\pi \in \mathcal{W}(\pi)^{\text{new}}$ be the normalized Whittaker newform with $W_\pi(1) = 1$. Let $\chi : \mathbf{Q}_q^\times \rightarrow \mathbf{C}^\times$ with $\chi(q) = 1$. Suppose that $L(s, \pi) = L(s, \pi \otimes \chi) = 1$. Put*

$$A_n^{(m)}(\chi) := \int_{\mathbf{Z}_q^\times} W_\pi\left(\begin{pmatrix} q^m y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ q^n & 1 \end{pmatrix}\right) \chi(y) d^\times y.$$

If $\chi \neq 1$, then $A_n^{(m)}(\chi) = 0$ unless $m = c(\pi) - c(\pi \otimes \chi)$ and $n = c(\pi) - c(\chi)$; in this case

$$A_{c(\pi) - c(\chi)}^{(c(\pi) - c(\pi \otimes \chi))}(\chi) = \varepsilon(1, \chi) \cdot \frac{\varepsilon(1/2, \pi)}{\varepsilon(1/2, \pi \otimes \chi)} \cdot \chi(-1)\zeta_q(1).$$

If $\chi = \mathbf{1}$ is the trivial character, then $A_n^{(m)}(\mathbf{1}) = 0$ unless $m = 0$ and $n \geq c(\pi) - 1$; in this case,

$$A_{c(\pi) - 1}^{(m)}(\mathbf{1}) = -|q|\zeta_q(1) \text{ and } A_n^{(m)}(\mathbf{1}) = 1 \text{ if } n \geq c(\pi).$$

PROOF. Let $A_n^{(m)} = A_n^{(m)}(\chi)$ and $c = c(\pi)$. Let $\varphi_n(a) := W_\pi\left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ q^n & 1 \end{pmatrix}\right)$ for $a \in \mathbf{Q}_q^\times$. Then φ_n belongs to the Krillov model $\mathcal{K}(\pi)$ of π with respect to $\psi_{\mathbf{Q}_q}$. Since $L(s, \pi) = 1$, $\varphi := \mathbb{1}_{\mathbf{Z}_q^\times}$ is a new vector

in $\mathcal{K}(\pi)$ and $\mathcal{K}(\tilde{\pi})$ (cf. [Sch02, §2.4]). Then $\pi\left(\begin{pmatrix} 0 & 1 \\ -q^c & 0 \end{pmatrix}\right)\varphi(a)\omega^{-1}(a) = \alpha \cdot \varphi(a)$ for some $\alpha \in \mathbf{C}^\times$. By the functional equation, we have

$$\int_{\mathbf{Q}_q^\times} \varphi_n(a)\chi(a) |a|^{s-\frac{1}{2}} d^\times a = \gamma(s, \pi \otimes \chi)^{-1} \int_{\mathbf{Q}_q^\times} \pi(w)\varphi_n(a)\omega^{-1}(a)\chi^{-1}(a) |a|^{\frac{1}{2}-s} d^\times a,$$

where

$$\gamma(s, \pi \otimes \chi)^{-1} = \frac{L(s, \pi \otimes \chi)}{L(1-s, \tilde{\pi} \otimes \chi^{-1})\varepsilon(s, \pi \otimes \chi)}.$$

By the relation

$$\begin{aligned} \pi(w)\varphi_n(a) &= \pi\left(\begin{pmatrix} 1 & -q^n \\ 0 & 1 \end{pmatrix} w\right)\varphi(a) = \psi_{\mathbf{Q}_q}(-aq^n)\pi\left(\begin{pmatrix} 1 & 0 \\ 0 & q^{-c} \end{pmatrix}\right)\left(\begin{pmatrix} 0 & 1 \\ -q^c & 0 \end{pmatrix}\right)\varphi(a) \\ &= \alpha \cdot \psi_{\mathbf{Q}_q}(-aq^n) \cdot \varphi(q^c a)\omega(a), \end{aligned}$$

we find that $\alpha = \varepsilon(1/2, \pi)$ and

$$\begin{aligned} &\sum_{m \in \mathbf{Z}} \int_{\mathbf{Z}_q^\times} \varphi_n(q^m y)\chi(y) d^\times y \cdot \chi(q^m) |q^m|^{s-\frac{1}{2}} \\ &= \gamma(s, \pi \otimes \chi)^{-1} \varepsilon(1/2, \pi) \cdot |q^c|^{s-\frac{1}{2}} \cdot \int_{\mathbf{Q}_q^\times} \psi_{\mathbf{Q}_q}\left(-\frac{a}{q^{c-n}}\right)\chi^{-1}|\cdot|^{\frac{1}{2}-s}(a)\varphi(a) d^\times a. \end{aligned}$$

Let $t = |q|^s$. From the above equation, we deduce that

$$\begin{aligned} &\sum_{m \in \mathbf{Z}} A_m^{(n)} \cdot \chi(q^m) |q^m|^{-\frac{1}{2}} \cdot t^m \\ &= \gamma(s, \pi \otimes \chi)^{-1} \cdot \varepsilon(1/2, \pi)\chi(-1) |q^c|^{-\frac{1}{2}} \cdot t^c \cdot \begin{cases} 0 & \text{if } c-n \neq c(\chi) > 0 \text{ or } c-n \geq 2, c(\chi) = 0, \\ \chi(q^{-c(\chi)})\varepsilon(1, \chi)\zeta_q(1) & \text{if } c-n = c(\chi) > 0, \\ 1 & \text{if } c-n \leq 0, c(\chi) = 0, \\ -|q|\zeta_q(1) & \text{if } c-n = 1, c(\chi) = 0. \end{cases} \end{aligned}$$

Since $L(s, \pi) = L(s, \pi \otimes \chi) = 1$, we have

$$\gamma(s, \pi \otimes \chi)^{-1} = \varepsilon(0, \pi \otimes \chi)^{-1} t^{-c(\pi \otimes \chi)}.$$

Comparing the coefficients of t^m , if $\chi \neq \mathbf{1}$, we find that $A_n^{(m)} \neq 0$ only when $c-n = c(\chi)$, and $m = c - c(\pi \otimes \chi)$. In this case

$$A_n^{(m)} = \chi(-q^{-m-c(\chi)}) |q|^{\frac{m-c}{2}} \varepsilon(1/2, \pi) \cdot \frac{\varepsilon(1, \chi)}{\varepsilon(0, \pi \otimes \chi)} \zeta_q(1).$$

If $\chi = \mathbf{1}$, and $A_n^{(m)} = 0$ unless $m = 0$, and

$$A_n^{(0)} = \begin{cases} 1, & \text{if } c-n \leq 0 \\ -|q|\zeta_q(1), & \text{if } c-n = 1 \\ 0 & \text{if } c-n \geq 2. \end{cases}$$

This completes the proof. \square

6.3. The case (Ib). In this case $\pi_1 = \chi_1|\cdot|^{-\frac{1}{2}}$ St is an unramified special representation, and π_2 and π_3 are discrete series with the local root number $\varepsilon(1/2, \Pi_q) = 1$. We first remark that if $L(s, \pi_2 \otimes \pi_3) \neq 1$, then by the minimality of π_3 combined with [GJ78, Proposition (1.2)], this implies that $\pi_3 = \tilde{\pi}_2 \otimes \sigma$ for some unramified character σ of \mathbf{Q}_q^\times and π_2 is also minimal. Hence, in view of [Pra90, Proposition 8.5] π_2 and π_3 must be unramified special in case (Ib) if $L(s, \pi_2 \otimes \pi_3) \neq 1$.

Proposition 6.6. *In case (Ib),*

(1) *if $L(s, \pi_2 \otimes \pi_3) = 1$, then we have*

$$I_q(\phi_q^* \otimes \tilde{\phi}_q^*) = \chi_1^2 | \cdot | (q^{c^*}) \varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes \nu_1) \varepsilon(1/2, \pi_2)^2 \varepsilon(1/2, \pi_3)^2 \cdot \frac{1}{B_{\Pi_q}} \cdot \frac{\zeta_q(2)^2}{\zeta_q(1)^2};$$

(2) if $L(s, \pi_2 \otimes \pi_3) \neq 1$, then $c_1 = c_2 = c_3 = 1$ and

$$I_q(\phi_q^* \otimes \tilde{\phi}_q^*) = I_q(\phi_q \otimes \tilde{\phi}_q) = \frac{2|q|}{B_{\Pi_q}} \cdot \frac{\zeta_q(2)^2}{\zeta_q(1)^2}.$$

PROOF. Now we suppose that $\pi_1 = \chi_1 |\cdot|^{-\frac{1}{2}} \text{St}$ is unramified special. Let $v_1 = \chi_1 |\cdot|^{-1}$. We use the realizations

$$\mathcal{V}_{\Pi_q} = \mathcal{B}(v_1, \chi_1)^0 \boxtimes \mathcal{W}(\pi_2) \boxtimes \mathcal{W}(\pi_3); \quad \mathcal{V}_{\tilde{\Pi}_q} = \mathcal{B}(v_1^{-1}, \chi_1^{-1})_0 \boxtimes \mathcal{W}(\tilde{\pi}_2) \boxtimes \mathcal{W}(\tilde{\pi}_3).$$

Here $\mathcal{B}(v_1, \chi_1)^0$ is the unique irreducible quotient space of $\mathcal{B}(v_1, \chi_1)$ and $\mathcal{B}(v_1^{-1}, \chi_1^{-1})_0$ is the unique irreducible sub-representation of $\mathcal{B}(v_1^{-1}, \chi_1^{-1})$ as in §5.1. Let $f_1 \in \mathcal{B}(v_1, \chi_1)^{\mathcal{U}_0(q)}$ be the unique function supported in $BwN(\mathbf{Z}_q)$ with $f_1(1) = 1$. Then the holomorphic image f_1^0 of f_1 in $\mathcal{V}_{\pi_1} = \mathcal{B}(v_1, \chi_1)^0$ is a new vector. Let $\tilde{f}_1 = M^*(v_1, \chi_1)f \otimes \omega_1^{-1}$. We may assume that $c_2 \geq c_3$ (so $c^* = c_2$). Let $W_3^* = \rho\left(\begin{pmatrix} q^{c_3-c_2} & 0 \\ 0 & 1 \end{pmatrix}\right)W_3$ and $\tilde{W}_3^* = W_3^* \otimes \omega_3^{-1}$. Then

$$\phi_q^* = f_1^0 \otimes W_2 \otimes W_3^*; \quad \tilde{\phi}_q = \tilde{f}_1 \otimes \tilde{W}_2 \otimes \tilde{W}_3^*.$$

By Corollary 5.2 and Lemma 6.4 (2), we obtain

$$(6.2) \quad \begin{aligned} I_q(\phi_q^* \otimes \tilde{\phi}_q^*) &= \frac{\mathcal{I}_q(W_2 \otimes W_3^* \otimes f_1, \tilde{W}_2 \otimes \tilde{W}_3^* \otimes \tilde{f}_1)}{\zeta_q(2)^2 L(1/2, \Pi_q) \cdot B_{\Pi_q}} \cdot \frac{\langle \rho(\tau_{c_1})W_1, \tilde{W}_1 \rangle}{\langle \rho(\tau_{c_1})f_1, \tilde{f}_1 \rangle} \cdot \frac{\zeta_q(2)^3}{\zeta_q(1)^3} \\ &= \frac{\gamma(1/2, \pi_2 \otimes \pi_3 \otimes v_1) \Psi(W_2, W_3^*, f_1)^2}{L(1/2, \Pi_q)} \cdot \frac{1}{B_{\Pi_q}}. \end{aligned}$$

In what follows, if $L(s, \pi_2 \otimes \pi_3) \neq 1$, then we write $\pi_2 = \chi_2 |\cdot|^{-\frac{1}{2}} \text{St}$ and $\pi_3 = \chi_3 |\cdot|^{-\frac{1}{2}} \text{St}$ with χ_2, χ_3 unramified. Using the integration formula

$$\int_{ZN \backslash G} F(g) dg = \frac{\zeta_q(2)}{\zeta_q(1)} \int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q} F\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} w \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}\right) dx \frac{d^\times y}{|y|}$$

for $F \in L^1(ZN \backslash G)$, we find that $\Psi(W_2, W_3^*, f_1)$ equals

$$\begin{aligned} &\frac{\zeta_q(2)}{\zeta_q(1)} \int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q} W_2\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} w \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}\right) W_3\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} w \begin{pmatrix} q^{c_3-c_2} & x \\ 0 & 1 \end{pmatrix}\right) v_1 |\cdot|^{\frac{1}{2}}(y) \mathbb{I}_{\mathbf{Z}_q}(x) dx \frac{d^\times y}{|y|} \\ &= \frac{\zeta_q(2) \chi_1 |\cdot|^{\frac{1}{2}}(q^{c_2})}{\zeta_q(1)} \int_{\mathbf{Q}_q^\times} W_2\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \tau_{c_2}\right) W_3\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \tau_{c_3}\right) v_1 |\cdot|^{\frac{1}{2}}(y) \mathbb{I}_{\mathbf{Z}_q}(x) dx \frac{d^\times y}{|y|} \\ &= \frac{\zeta_q(2)}{\zeta_q(1)} \cdot \chi_1 |\cdot|^{\frac{1}{2}}(q^{c_2}) \cdot \varepsilon(1/2, \pi_2) \varepsilon(1/2, \pi_3) \begin{cases} 1 & \text{if } L(s, \pi_2 \otimes \pi_3) = 1, \\ L(-1/2, \chi_1 \chi_2 \chi_3) & \text{if } L(s, \pi_1 \otimes \pi_3) \neq 1. \end{cases} \end{aligned}$$

If $L(s, \pi_2 \otimes \pi_3) = 1$, then one verifies easily that $\gamma(1/2, \pi_2 \otimes \pi_3 \otimes v_1) = \varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes v_1)$ and $L(s, \Pi_q) = 1$, so we obtain the claimed expression of $I_q(\phi_q^* \otimes \tilde{\phi}_q^*)$ in this case by substituting the above equation into (6.2).

Suppose that $L(s, \pi_2 \otimes \pi_3) \neq 1$. Then $c_1 = c_2 = c_3 = 1$ and $\varepsilon(1/2, \pi_i) = -\chi_i |\cdot|^{-\frac{1}{2}}(q)$ for $i = 1, 2, 3$. Hence, $W_3^* = W_3$ and

$$\Psi(W_2, W_3^*, f_1)^2 = \Psi(W_2, W_3, f_1)^2 = |q|^2 \cdot \frac{\zeta_q(2)^2}{\zeta_q(1)^2} \cdot L(-1/2, \chi_1 \chi_2 \chi_3)^2.$$

On the other hand, by [Pra90, Proposition 8.6], $\varepsilon(1/2, \Pi_q) = 1$ implies that

$$\chi_1 \chi_2 \chi_3(q) = -|q|^{\frac{3}{2}}.$$

By [GJ78, Proposition 1.4], $\varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes v_1) = |q|^{-1}$ and

$$\frac{L(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1)}{L(1/2, \pi_2 \otimes \pi_3 \otimes v_1)} = \frac{L(1/2, \chi_1 \chi_2 \chi_3)}{L(-3/2, \chi_1 \chi_2 \chi_3)} = 2L(1/2, \chi_1 \chi_2 \chi_3),$$

and a simple computation of the Langlands parameter for Π_q shows

$$L(s, \Pi_q) = L(s, \chi_1 \chi_2 \chi_3) L(s-1, \chi_1 \chi_2 \chi_3)^2.$$

We thus obtain

$$\gamma(1/2, \pi_2 \otimes \pi_3 \otimes v_1) = 2|q|^{-1} \cdot \frac{L(1/2, \Pi_q)}{L(-1/2, \chi_1 \chi_2 \chi_3)^2}.$$

The desired formula of $I_q(\phi_q^* \otimes \tilde{\phi}_q^*) = I_q(\phi_q \otimes \tilde{\phi}_q)$ in this case can be deduced immediately by combining (6.2) with the above formulae of $\Psi(W_2, W_3^*, f_1)$ and the γ -factor. \square

Remark 6.7. In the case where $L(s, \pi_2 \otimes \pi_3) \neq 1$, i.e. π_i are special unramified, the integral $I_q(\phi_q^* \otimes \tilde{\phi}_q^*)$ was computed in [II10, page 1405-1406], from which we have $I_q(\phi_q^* \otimes \tilde{\phi}_q^*) = 2|q|(1+|q|)$. Our computation agrees with the result therein (note that $B_{\Pi_q} = \zeta_q(2)^3 \zeta_q(1)^{-3}$).

6.4. The ramified case (IIa). In this case, π_2, π_3 are discrete series and $L(s, \pi_2 \otimes \pi_3) \neq 1$. As we have remarked in the previous subsection, $\pi_3 \simeq \tilde{\pi}_2 \otimes \sigma$ for some unramified character σ of \mathbf{Q}_q^\times and π_1 must be spherical. Let $\tau_{\mathbf{Q}_{q^2}}$ be the quadratic character associated with the unramified quadratic field extension \mathbf{Q}_{q^2} of \mathbf{Q}_q . We say a discrete series π is of type **1** if $\pi \simeq \pi \otimes \tau_{\mathbf{Q}_{q^2}}$ and is of type **2** if $\pi \not\simeq \pi \otimes \tau_{\mathbf{Q}_{q^2}}$.

The following lemma for minimal supercuspidal representations should be well-known to experts. We include a proof here for the reader's convenience.

Lemma 6.8. *Let π be a minimal supercuspidal representation with central character ω .*

(1) *Let χ be a charatcer of \mathbf{Q}_q^\times . Then we have the following conductor formula*

$$c(\pi \otimes \chi) = \begin{cases} c(\pi) & \text{if } c(\pi) \geq 2c(\chi), \\ 2c(\chi) & \text{if } c(\pi) < 2c(\chi). \end{cases}$$

Here recall that $c(?)$ denotes the exponent of the conductor of $?$.

(2) *If π is of type **1**, then $c(\pi)$ is even and $L(s, \pi \otimes \tilde{\pi}) = \zeta_q(2s)$. If π is of type **2**, then $c(\pi)$ is odd and $L(s, \pi \otimes \tilde{\pi}) = \zeta_q(s)$.*

PROOF. Let $c = c(\pi) \geq 2$. To prove the first assertion, we begin with an immediate consequence of [JL70, Proposition 2.11 (i)]. Let $\chi_0 = \chi|_{\mathbf{Z}_q^\times}$ and $\omega_0 = \omega|_{\mathbf{Z}_q^\times}$. If $\chi_0 \omega_0 \neq 1$, then there exists a character σ such that

$$(6.3) \quad c(\pi \otimes \sigma) = c + c(\pi \otimes \chi) - 2c(\chi\omega)$$

and if $\chi_0 \neq 1, \omega_0^{-1}$, then either of the following condition holds:

(i) $\sigma|_{\mathbf{Z}_q^\times} \neq 1, \chi_0$ and

$$c(\sigma) = c - c(\chi\omega), \quad c(\sigma\chi^{-1}) = c(\pi \otimes \chi) - c(\chi\omega),$$

(ii) $\sigma|_{\mathbf{Z}_q^\times} = 1, c(\chi) = c(\pi \otimes \chi) - c(\chi\omega)$ and $c(\chi\omega) - c \geq -1$;

(iii) $\sigma|_{\mathbf{Z}_q^\times} = \chi_0, c(\chi) = c(\pi) - c(\chi\omega)$ and $c(\chi\omega) - c(\pi \otimes \chi) \geq -1$.

To see it, we set $\rho = \chi_0^{-1} \omega_0^{-1}, \nu = \omega_0^{-1}, m = c(\chi\omega), p = m - c(\pi \otimes \chi)$ and $n = m - c(\pi)$ in the equality proved in [JL70, Proposition 2.11 (i)], from which we see immediately that the equality shows the existence of desired σ by noting that $C_n(\rho^{-1} \omega^{-1}) \neq 0$ if and only if $n = c(\pi \otimes \rho)$. Note that (6.3) implies that

$$c(\pi \otimes \chi) \geq 2c(\chi\omega) \text{ for all } \chi$$

by the minimality of π . In particular, $c(\omega) \leq c/2$. Suppose that $c(\chi) > c/2$. Then $c(\chi\omega) = c(\chi)$ and σ satisfies either (i) or (ii). In case (ii), we have $c(\pi \otimes \chi) = 2c(\chi)$. In case (i), $c(\sigma) = c - c(\chi) < c/2$, and hence we also have $c(\pi \otimes \chi) = c(\chi) + c(\sigma\chi^{-1}) = 2c(\chi)$. Now we suppose that $c(\chi) \leq c/2$. If $\chi_0 = \omega_0^{-1}$, then $c(\pi \otimes \chi) = c(\tilde{\pi}) = c$, so we may assume $\chi_0 \neq \omega_0^{-1}$. It suffices to show $c(\pi \otimes \chi) \leq c$. Note that $c(\chi\omega) \leq c/2$. In case (iii), $c(\pi \otimes \chi) \leq c(\chi\omega) + 1 \leq c$, and in the case (ii), $c(\pi \otimes \chi) = c(\chi) + c(\chi\omega) \leq c$. We consider case (i). We have $c(\sigma) = c - c(\chi\omega) \geq c/2$. If $c(\sigma) > c/2$, then

$$c(\pi \otimes \chi) = c(\chi\omega) + c(\sigma^{-1}\chi) = c(\chi\omega) + c(\sigma) = c.$$

If $c(\sigma) = c/2$, then we also have $c(\pi \otimes \chi) \leq c/2 + c/2 = c$. This finishes the proof of the first assertion.

We proceed to show the second assertion. This is [Hid90, Proposition 6.1]. We give a more elementary proof. The local L -factor of $L(s, \pi \otimes \tilde{\pi})$ is given in [GJ78, Corollary (1.3)]. To see the parity of the conductor, we note that $\pi \simeq \pi \otimes \tau_{\mathbf{Q}_{q^2}}$ if and only if $\varepsilon(s, \pi \otimes \chi) = \varepsilon(s, \pi \otimes \chi \tau_{\mathbf{Q}_{q^2}})$ for all character $\chi : \mathbf{Q}_q^\times \rightarrow \mathbf{C}^\times$ as π is supercuspidal. Since $\tau_{\mathbf{Q}_{q^2}}$ is unramified, this is equivalent to saying $(-1)^{c(\pi \otimes \chi)} = 1$ for all χ . It follows from part (1) that π is of type **1** if and only if $c(\pi)$ is even. \square

Proposition 6.9. *Let $r = \left\lceil \frac{c(\pi_2)}{2} \right\rceil$. We have*

$$I_q(\phi_q^* \otimes \tilde{\phi}_q^*) = \chi_1^{-2} |\cdot| (q^r) \cdot \varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1) \cdot \frac{1}{B_{\Pi_q}} \cdot \frac{\zeta_q(2)^2}{\zeta_q(1)^2} \cdot \begin{cases} (1 + |q|) & \text{if } \pi_2 \text{ is of type } \mathbf{1}, \\ 1 & \text{if } \pi_2 \text{ is of type } \mathbf{2}. \end{cases}$$

PROOF. After a unramified twist, we may assume that $\pi_1 = \chi_1 \boxplus v_1$ with $\chi_1 = |\cdot|^{s-\frac{1}{2}}$ and $v_1 = |\cdot|^{\frac{1}{2}-s}$ for some $s \in \mathbf{C}$ and $\pi_3 = \tilde{\pi}_2$. Let $\pi = \pi_2$ be a minimal discrete series. We use the realizations as in (6.5). Let f_1 be the normalized new vector in $\mathcal{B}(|\cdot|^{s-\frac{1}{2}}, |\cdot|^{\frac{1}{2}-s})$ and let $f_1^* = \rho\left(\begin{pmatrix} q^{-r} & 0 \\ 0 & 1 \end{pmatrix}\right)f_1$. As in the previous cases, by Corollary 5.2 we obtain

$$(6.4) \quad \begin{aligned} I_q(\phi_q^* \otimes \tilde{\phi}_q^*) &= \frac{\zeta_q(1) \mathcal{I}_q(W_2 \otimes W_3 \otimes f_1^*, \tilde{W}_2 \otimes \tilde{W}_3 \otimes \tilde{f}_1^*)}{\zeta_q(2)^2 L(1/2, \Pi_q) B_{\Pi_q}} \cdot \frac{\langle W_{\pi_1}, \tilde{W}_{\pi_1} \rangle}{\langle f_1, \tilde{f}_1 \rangle} \cdot \frac{\zeta_q(2)^3}{\zeta_q(1)^3} \\ &= \frac{\zeta_q(1) \gamma(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1) \cdot \Psi(W_2, W_3, f_1^*)^2}{\zeta_q(2)^2 L(1/2, \Pi_q)} \cdot \frac{\langle W_{\pi_1}, \tilde{W}_{\pi_1} \rangle}{\langle f_1, \tilde{f}_1 \rangle} \cdot \frac{1}{B_{\Pi_q}} \cdot \frac{\zeta_q(2)^3}{\zeta_q(1)^3}. \end{aligned}$$

Define the function $\mathbf{W} : ZN \backslash G \rightarrow \mathbf{C}$ by

$$\mathbf{W}(g) := W_2(g) W_3\left(\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} g\right).$$

We compute $\Psi(W_2, W_3, f_1^*)$ in the following two subcases.

Subcase (a): $\pi_i = \chi_i |\cdot|^{-\frac{1}{2}} \text{St}$ are unramified special for $i = 2, 3$. Then π_2 is of type $\mathbf{2}$ and $r = 1$. We have

$$\Psi(W_2, W_3, f_1^*) = \text{vol}(K_0(q))(J_1 + J_2),$$

where

$$\begin{aligned} J_1 &= |q|^{-s} \int_{\mathbf{Q}_q^\times} \mathbf{W}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) |y|^{s-1} d^\times y, \\ J_2 &= \sum_{x \in \mathbf{Z}/q\mathbf{Z}} |q|^s \int_{\mathbf{Q}_q^\times} \mathbf{W}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} w\right) |y|^{s-1} d^\times y. \end{aligned}$$

By a direct calculation, we find that

$$\begin{aligned} J_1 &= |q|^{-s} L(\mathbf{s}, \chi_2 \chi_3), \\ J_2 &= |q|^s \cdot q \cdot |q|^2 \cdot \chi_2^{-1} \chi_3^{-1} |\cdot|^{-s}(q) \cdot L(\mathbf{s}, \chi_1 \chi_2) = |q| \chi_1 \chi_2 (q^{-1}) L(\mathbf{s}, \chi_2 \chi_3). \end{aligned}$$

Note that $\omega_2 \omega_3 = \chi_2^2 \chi_3^2 |\cdot|^{-2} = 1$. Hence

$$\begin{aligned} \Psi(W_2, W_3, f_1^*) &= \frac{1}{1+q} |q|^{-s} \cdot (1 + \chi_2 \chi_3 |\cdot|^{s-1}(q)) \cdot L(\mathbf{s}, \chi_2 \chi_3) \\ &= \frac{\zeta_q(2)}{\zeta_q(1)} |q|^{1-s} \frac{L(\mathbf{s}, \chi_2 \chi_3) L(\mathbf{s}-1, \chi_2 \chi_3)}{\zeta_q(2\mathbf{s})} \\ &= \frac{\zeta_q(2)}{\zeta_q(1)} |q|^{1-s} \frac{L(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1)}{L(1, \chi_1 v_1^{-1})}. \end{aligned}$$

Subcase (b): π_2 and π_3 are supercuspidal. In this case,

$$\begin{aligned} \Psi(W_2, W_3, f_1^*) &= \frac{\zeta_q(2)}{\zeta_q(1)} \int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q} \mathbf{W}\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}\right) |y|^{s-1} f_1\left(\begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}\right) \begin{pmatrix} q^{-r} & 0 \\ 0 & 1 \end{pmatrix} dx d^\times y \\ &= \frac{\zeta_q(2)}{\zeta_q(1)} |q|^{-rs} \sum_{n \in \mathbf{Z}} J_n, \end{aligned}$$

where

$$\begin{aligned}
J_n &= \int_{\mathbf{Q}_q^\times} \int_{q^n \mathbf{Z}_q^\times} \mathbf{W} \left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ q^n & 1 \end{pmatrix} \right) |y|^{s-1} f_3 \left(\begin{pmatrix} 1 & 0 \\ q^{n-r} & 1 \end{pmatrix} \right) dx dy \\
&= |q^n| (1 - |q|) \sum_{m \in \mathbf{Z}} |q^m|^{s-1} \cdot f_1 \left(\begin{pmatrix} 1 & 0 \\ q^{n-r} & 1 \end{pmatrix} \right) \int_{\mathbf{Z}_q^\times} \mathbf{W} \left(\begin{pmatrix} q^m u & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ q^n & 1 \end{pmatrix} \right) d^\times u \\
&= |q^n| (1 - |q|) \sum_{m \in \mathbf{Z}} |q^m|^{s-1} f_1 \left(\begin{pmatrix} 1 & 0 \\ q^{n-r} & 1 \end{pmatrix} \right) \sum_{\chi \in \widehat{\mathbf{Z}}_q^\times} A_{\pi_2, n}^{(m)}(\chi) A_{\pi_3, n}^{(m)}(\chi^{-1}) \chi(-1),
\end{aligned}$$

where

$$A_{\pi_i, n}^{(m)}(\chi) := \int_{\mathbf{Z}_q^\times} W_i \left(\begin{pmatrix} q^m u & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ q^n & 1 \end{pmatrix} \right) d u.$$

In the case $\chi \neq \mathbf{1}$, by Lemma 6.5, we have

$$A_{\pi_2, n}^{(m)}(\chi) A_{\pi_3, n}^{(m)}(\chi^{-1}) \chi(-1) = |q|^{c-n} \zeta_q(1)^2$$

if $n = c - c(\chi)$ and

$$m = c - c(\pi \otimes \chi) = \begin{cases} 0 & \text{if } n \geq r, \\ 2n - c & \text{if } n < r, \end{cases}$$

by Lemma 6.8 (2), and $A_{\pi_2, n}^{(m)}(\chi) A_{\pi_3, n}^{(m)}(\chi^{-1}) = 0$ otherwise. If $\chi = \mathbf{1}$, then

$$A_{\pi_2, n}^{(m)}(\mathbf{1}) A_{\pi_3, n}^{(m)}(\mathbf{1}) = |q|^{2n} \zeta_q(1)^2$$

if $c - n = 1$. Therefore, if $n < r$, then

$$\begin{aligned}
J_n &= (1 - |q|) |q|^n |q|^{(2n-c)(s-1)} |q|^{2(r-n)s} \cdot |q|^{c-n} \zeta_q(1)^2 \# \{ \chi \mid c(\chi) = c - n \} \\
&= (1 - |q|) |q|^{(2r-c)s + c - n}.
\end{aligned}$$

If $r \leq n < c - 1$, then

$$J_n = (1 - |q|) |q|^n.$$

If $n = c - 1$, then

$$J_{c-1} = (1 - |q|) |q|^{c-1} (|q| (q - 1 - 1) + |q|^2) \zeta_q(1)^2 = (1 - |q|) |q|^{c-1}.$$

If $n \geq c$, then $J_{\geq c} = |q|^c$. Combining the above equations, we find that $\Psi(W_2, W_3, f_1^*)$ equals

$$\begin{aligned}
\frac{\zeta_q(2)}{\zeta_q(1)} |q|^{-rs} \sum_{n \in \mathbf{Z}} J_n &= \frac{\zeta_q(2)}{\zeta_q(1)} |q|^{-rs} \left(J_{r-1}^- + \sum_{n=1}^r J_n + J_c^+ \right) \\
&= \frac{\zeta_q(2)}{\zeta_q(1)} |q|^{-rs} (|q|^{(2r-c)s} |q|^{c+1-r} + |q|^r - |q|^c + |q|^c) \\
&= \frac{\zeta_q(2)}{\zeta_q(1)} |q|^{-rs} \begin{cases} |q|^{\frac{s}{2}} (1 + |q|) & \text{if } c \text{ is even } (\pi_2 \text{ is of type } \mathbf{1}), \\ |q|^{\frac{c+1}{2}} (1 + |q|^s) & \text{if } c \text{ is odd } (\pi_2 \text{ is of type } \mathbf{2}). \end{cases}
\end{aligned}$$

On the other hand, when π_2 and π_3 are supercuspidal, it is easy to see that

$$\frac{L(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1)}{L(1, \chi_1 v_1^{-1})} = \begin{cases} 1 & \text{if } \pi_2 \text{ is of type } \mathbf{1}, \\ 1 + |q|^s & \text{if } \pi_2 \text{ is of type } \mathbf{2}. \end{cases}$$

We thus conclude that in either subcase (a) or subcase (b),

$$\Psi(W_2, W_3, f_1^*) = \frac{\zeta_q(2)}{\zeta_q(1)} |q|^{r(1-s)} \frac{L(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1)}{L(1, \chi_1 v_1^{-1})} \begin{cases} 1 + |q| & \text{if } \pi_2 \text{ is of type } \mathbf{1}, \\ 1 & \text{if } \pi_2 \text{ is of type } \mathbf{2}. \end{cases}$$

Substituting the above equation and Lemma 6.4 into (6.4), we obtain

$$\begin{aligned} I_q(\phi_q^* \otimes \tilde{\phi}_q^*) &= \frac{\zeta_q(2)\gamma(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1) \cdot \Psi(W_2, W_3, f_1^*)^2}{\zeta_q(1)^2 L(1/2, \Pi_q)} \cdot \frac{\zeta_q(1)^2 L(1, \chi_1 v_1^{-1})^2}{\zeta_q(2)} \\ &= \varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1) \chi_1^{-2} |\cdot| (q^r) \cdot \frac{1}{B_{\Pi_q}} \cdot \frac{\zeta_q(2)^2}{\zeta_q(1)^2} \cdot \begin{cases} (1 + |q|) & \text{if } \pi_2 \text{ is of type } \mathbf{1}, \\ 1 & \text{if } \pi_2 \text{ is of type } \mathbf{2} \end{cases} \end{aligned}$$

by noting that $L(s, \Pi_q) = L(s, \pi_2 \otimes \pi_3 \otimes \chi_1) L(s, \pi_2 \otimes \pi_3 \otimes v_1)$. This finishes the proof. \square

6.5. The ramified case (IIb). Finally, we consider the case where π_2 and π_3 are discrete series, π_3 is minimal and $L(s, \pi_2 \otimes \pi_3) = 1$. It is also assumed that $\pi_1 = \chi_1 \boxplus v_1$ is a principal series with $c(\chi_1) = 0$ and $c(v_1) \leq 1$.

Proposition 6.10. *Let $c^* = \max\{c_2, c_3\}$. We have*

$$I_q(\phi_q^* \otimes \tilde{\phi}_q^*) = \omega_1(-1) \chi_1^{-2} |\cdot| (q^{c^*}) \varepsilon(1/2, \pi_1)^2 \cdot \varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1) \cdot \frac{1}{B_{\Pi_q}} \cdot \frac{\zeta_q(2)^2}{\zeta_q(1)^2}.$$

PROOF. In this case, we use the realizations

$$(6.5) \quad \mathcal{V}_{\Pi_q} = \mathcal{B}(\chi_1, v_1) \boxtimes \mathcal{W}(\pi_2) \boxtimes \mathcal{W}(\pi_3); \quad \mathcal{V}_{\tilde{\Pi}_q} = \mathcal{B}(\chi_1^{-1}, v_1^{-1}) \boxtimes \mathcal{W}(\tilde{\pi}_2) \boxtimes \mathcal{W}(\tilde{\pi}_3).$$

Let $f_1 \in \mathcal{B}(\chi_1, v_1)^{\text{new}}$ be the new vector with $f_1(1) = 1$. Define the section $f_1^* \in \mathcal{B}(\chi_1, v_1)^{U_0(q^{c^*})}$ by

$$f_1^* = \rho\left(\begin{pmatrix} q^{-c^*} & 0 \\ 0 & 1 \end{pmatrix}\right) f_1 - v_1^{-1} |\cdot|^{\frac{1}{2}}(q) \rho\left(\begin{pmatrix} q^{1-c^*} & 0 \\ 0 & 1 \end{pmatrix}\right) f_1 \text{ if } c_1 = c(v_1) = 0$$

and $f_1^* = \rho\left(\begin{pmatrix} q^{1-c^*} & 0 \\ 0 & 1 \end{pmatrix}\right) f_1$ if $c_1 = 1$. Then f_1^* is the section supported in the $BU_0(q^{c^*})$ with $f_1^*(1) = \chi_1 |\cdot|^{\frac{1}{2}}(q^{c_1 - c^*}) L(1, \chi_1 v_1^{-1})^{-1}$. Let $\tilde{f}_1 = M^*(\chi_1, v_1) f_1 \otimes \omega_1^{-1}$. Then we have

$$\begin{aligned} \tilde{f}_1 &= \rho\left(\begin{pmatrix} q^{-c^*} & 0 \\ 0 & 1 \end{pmatrix}\right) \tilde{f}_1 \cdot \omega_1(p^{-c^*}) - v_1^{-1} |\cdot|^{\frac{1}{2}}(q) \rho\left(\begin{pmatrix} q^{1-c^*} & 0 \\ 0 & 1 \end{pmatrix}\right) \tilde{f}_1 \cdot \omega_1(p^{1-c^*}) \\ &= M^*(\chi_1, v_1) f_1^* \otimes \omega_1^{-1} \text{ if } c_1 = 0. \end{aligned}$$

A direct computation shows that

$$\begin{aligned} \Psi(W_2, W_3, f_1^*) &= \frac{\zeta_q(2)}{\zeta_q(1)} \int_{\mathbf{Q}_q^\times} \int_{\mathbf{Q}_q} W_2\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}\right) W_3\left(\begin{pmatrix} -y & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}\right) \chi_1 |\cdot|^{\frac{1}{2}}(y) f_1^*\left(\begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}\right) dx \frac{d^\times y}{|y|} \\ &= \frac{\chi_1 |\cdot|^{\frac{1}{2}}(q^{c_1 - c^*}) \zeta_q(2) |q|^{c^*}}{L(1, \chi_1 v_1^{-1}) \zeta_q(1)} \int_{\mathbf{Q}_q} W_2\left(\begin{pmatrix} y & 0 \\ 0 & 1 \end{pmatrix}\right) W_3\left(\begin{pmatrix} -y & 0 \\ 0 & 1 \end{pmatrix}\right) \chi_1 |\cdot|^{-\frac{1}{2}}(y) d^\times y \\ &= \frac{\zeta_q(2) \chi_1 (q^{c_1 - c^*}) |q|^{\frac{c_1 + c^*}{2}}}{\zeta_q(1) L(1, \chi_1 v_1^{-1})}. \end{aligned}$$

The last equality follows from the fact that either $L(s, \pi_2) = 1$ or $L(s, \pi_3) = 1$ in case (IIb). By Corollary 5.2, the above equation and Lemma 6.4 (1), we obtain

$$\begin{aligned} I_q(\phi_q^* \otimes \tilde{\phi}_q^*) &= \frac{\mathcal{I}_q(W_2 \otimes W_3 \otimes f_1^*, \tilde{W}_2 \otimes \tilde{W}_3 \otimes \tilde{f}_1^*)}{\zeta_q(2)^2 L(1/2, \Pi_q) B_{\Pi_q}} \cdot \frac{\langle \rho(\tau_{c_1}) W_{\pi_1}, \tilde{W}_{\pi_1} \rangle}{\langle \rho(\tau_{c_1}) f_1, \tilde{f}_1 \rangle} \cdot \frac{\zeta_q(2)^3}{\zeta_q(1)^3} \\ &= \frac{\gamma(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1) \Psi(W_2, W_3, f_1^*)^2}{L(1/2, \Pi_q) B_{\Pi_q}} \cdot \chi_1^2 |\cdot| (q^{-c_1}) \varepsilon(1/2, \pi_1)^2 \omega_1(-1) L(1, \chi_1 v_1^{-1})^2 \\ &= \frac{\chi_1^{-2} (q^{c^*}) |q|^{c^*} \varepsilon(1/2, \pi_2 \otimes \pi_3 \otimes \chi_1)}{B_{\Pi_q}} \cdot \frac{\zeta_q(2)^2}{\zeta_q(1)^2} \cdot \varepsilon(1/2, \pi_1)^2 \omega_1(-1). \end{aligned}$$

The lemma follows. \square

6.6. **The p -adic interpolation of normalized local zeta integrals $\mathcal{J}_{\Pi_{\underline{Q},q}}^*$.** In this subsection, we compute the normalized local zeta integrals $\mathcal{J}_{\Pi_{\underline{Q},q}}^* = \mathcal{J}_{\Pi_q}^*$ in (3.29) and show these integrals can be p -adically interpolated by an Iwasawa function in $\underline{Q} \in \mathfrak{X}_{\mathcal{R}}^+$. We begin with recalling some facts. If $\mathcal{F} \in \mathbf{I}[[q]]$ is a primitive Hida family of tame conductor N and $\underline{Q} \in \mathfrak{X}_{\mathbf{I}}^+$ be a classical point, as in the introduction we denote by $V_{\mathcal{F}_Q}$ the associated p -adic Galois representation, and for each prime ℓ , let $\text{WD}_{\ell}(V_{\mathcal{F}_Q})$ the representation of the Weil-Deligne group $W'_{\mathbf{Q}_{\ell}}$ attached to $V_{\mathcal{F}_Q}$. Let $\ell \neq p$ be a prime. On the automorphic side, denote by $\text{Rec}_{\mathbf{Q}_{\ell}}$ the local Langlands reciprocity map from the set of isomorphism classes of irreducible representations of $\text{GL}_n(\mathbf{Q}_{\ell})$ to the set of isomorphism classes of n -dimensional representations of Weil-Deligne group $W'_{\mathbf{Q}_{\ell}}$ over $\overline{\mathbf{Q}_p}$ ([HT01a]). Then

$$(6.6) \quad \text{Rec}_{\mathbf{Q}_q}(\pi_{\mathcal{F}_Q,\ell} \otimes |\cdot|_{\ell}^{\frac{1-kQ}{2}}) = \text{WD}_{\ell}(V_{\mathcal{F}_Q}); \quad \varepsilon_{\ell}(1/2, \pi_{\mathcal{F}_Q}) = |N|_{\ell}^{\frac{kQ}{2}} \varepsilon(\text{WD}_{\ell}(V_{\mathcal{F}_Q})).$$

We recall the standard fact for the p -adic interpolation of local constants in Hida families.

Lemma 6.11. *There exists $\varepsilon_{\ell}(\mathcal{F}) \in \mathbf{I}^{\times}$ such that*

$$\varepsilon_{\ell}(\mathcal{F})(Q) = \varepsilon(\text{WD}_q(V_{\mathcal{F}_Q}))$$

for every classical point $Q \in \mathfrak{X}_{\mathbf{I}}^+$. Moreover, if $\mathcal{G} \in \mathbf{I}[[q]]$ is another primitive Hida family, then there exists $\varepsilon(\mathcal{F} \otimes \mathcal{G}) \in (\mathbf{I} \widehat{\otimes}_{\mathcal{O}} \mathbf{I})^{\times}$ such that

$$\varepsilon_{\ell}(\mathcal{F} \otimes \mathcal{G})(Q_1, Q_2) = \varepsilon(\text{WD}_{\ell}(V_{\mathcal{F}_{Q_1}} \otimes V_{\mathcal{G}_{Q_2}}))$$

for every classical points $(Q_1, Q_2) \in \mathfrak{X}_{\mathbf{I}}^+ \times \mathfrak{X}_{\mathbf{I}}^+$.

PROOF. This is a simple consequence of the description of $\rho_{\mathcal{F}}|_{G_{\mathbf{Q}_{\ell}}}$ together with the rigidity of automorphic types of Hida families in §3.2. We can actually make explicit the construction of $\varepsilon_{\ell}(\mathcal{F})$ as follows. Let $Q \in \mathfrak{X}_{\mathbf{I}}^+$ be any arithmetic point. If $\pi_{\mathcal{F}_Q,\ell}$ is a principal series, then $\rho_{\mathcal{F},\ell} \otimes \langle \varepsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{1/2}|_{G_{\mathbf{Q}_{\ell}}} \simeq \alpha_{\mathcal{F},\ell} \xi_1 \varepsilon_{\text{cyc}}^{1/2} \oplus \alpha_{\mathcal{F},\ell}^{-1} \xi_2 \varepsilon_{\text{cyc}}^{1/2}$ is reducible with $\xi_1, \xi_2 : G_{\mathbf{Q}_{\ell}} \rightarrow \overline{\mathbf{Q}}^{\times}$ finite order characters and $\alpha_{\mathcal{F},\ell} : G_{\mathbf{Q}_{\ell}} \rightarrow \mathbf{I}^{\times}$ unramified, and it is not difficult to see that

$$\varepsilon_{\ell}(\mathcal{F}) = \varepsilon(0, \xi) \varepsilon(0, \xi') \cdot \alpha_{\mathcal{F},\ell}(\text{Frob}_{\ell}^{n_1 - n_2}) \langle \varepsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{\frac{1}{2}}(\text{Frob}_{\ell}^{n_1 + n_2}) \cdot |\ell|_{\ell}^{n_1 + n_2},$$

where $n_1 = c(\xi_1)$ and $n_2 = c(\xi_2)$. If $\pi_{\mathcal{F},Q}$ is special, then $\rho_{\mathcal{F},\ell}|_{G_{\mathbf{Q}_{\ell}}} \otimes \langle \varepsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{1/2}$ is a non-split extension of ξ by $\xi \varepsilon_{\text{cyc}}$ for a finite order character $\xi : G_{\mathbf{Q}_{\ell}} \rightarrow \overline{\mathbf{Q}}^{\times}$, and letting $n' = c(\xi)$, we have

$$\varepsilon(\mathcal{F}) = \varepsilon(0, \xi)^2 \langle \varepsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{-1} \varepsilon_{\text{cyc}}(\text{Frob}_{\ell}^{n'}) \cdot \begin{cases} - \langle \varepsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{1/2} & \text{if } n' = 0 \\ 1 & \text{if } n' > 0 \end{cases}.$$

If $\pi_{\mathcal{F}_Q,\ell}$ is supercuspidal, then $\rho_{\mathcal{F},\ell}|_{G_{\mathbf{Q}_{\ell}}} = \rho_0 \otimes \langle \varepsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{-1/2}$ for some irreducible representation $\rho_0 : G_{\mathbf{Q}_{\ell}} \rightarrow \text{GL}_2(\overline{\mathbf{Q}})$ of finite image and of conductor $\ell^{n''}$, and we have

$$\varepsilon_{\ell}(\mathcal{F}) = \varepsilon(\text{WD}_{\ell}(\rho_0)) \cdot \langle \varepsilon_{\text{cyc}} \rangle_{\mathbf{I}}^{\frac{1}{2}}(\text{Frob}_{\ell}^{n''}).$$

The case $\rho_{\mathcal{F}} \otimes \rho_{\mathcal{G}}$ can be treated in the same manner by the formulae of ε -factors in [GJ78]. We omit the details. \square

We recall that the finite set Σ_{exc} in (1.5) is given by

$$\Sigma_{\text{exc}} = \left\{ q \in \Sigma_f^{(\text{IIa})} \sqcup \Sigma_g^{(\text{IIa})} \sqcup \Sigma_h^{(\text{IIa})} \mid \text{either of } \pi_{f,q}, \pi_{g,q}, \pi_{h,q} \text{ is supercuspidal of type } \mathbf{1} \right\}.$$

Proposition 6.12. *With Hypothesis 6.1, for each $q \mid N$ and $q \notin \Sigma^{-}$, there exists a unique element $I_{\mathbf{F},q} \in \mathcal{R}^{\times}$ such that*

$$\mathcal{J}_{\Pi_{\underline{Q},q}}^* = I_{\mathbf{F},q}(\underline{Q}) \cdot \begin{cases} (1 + q^{-1}) & \text{if } q \in \Sigma_{\text{exc}}, \\ 1 & \text{otherwise.} \end{cases}$$

for all $\underline{Q} \in \mathfrak{X}_{\mathcal{R}}^+$.

PROOF. We shall express $\mathcal{S}_{\Pi_q}^*$ in terms of epsilon factors of Galois representation under the setting in §6.1. As before, let $(f, g, h) = (\mathbf{f}_{Q_1}, \mathbf{g}_{Q_2}, \mathbf{h}_{Q_3})$ be p -stabilized newform of weight k_1, k_2, k_3 . Let $\chi_{\mathbf{F}} : G_{\mathbf{Q}} \rightarrow \mathcal{R}^\times$ be the unique character such that $\chi_{\mathbf{F}}^{-2} = (\det \rho_{\mathbf{f}} \otimes \det \rho_{\mathbf{g}} \otimes \det \rho_{\mathbf{h}}) \varepsilon_{\text{cyc}}^{-1}$. Then $\chi_{\mathbf{F}}$ is unramified at q . If χ_F is the specialization of $\chi_{\mathbf{F}}$ at \underline{Q} , then

$$\text{Rec}_{\mathbf{Q}_q}(\omega_F^{-1/2} |\cdot|_q^{\frac{w_{\underline{Q}}+1}{2}}) = \chi_{\mathbf{F}\underline{Q}}|_{W_{\mathbf{Q}_q}}.$$

As before, $c_2 = c_q(\pi_g)$, $c_3 = c_q(\pi_h)$ and $c^* = \max\{c_2, c_3\}$. Write $|\cdot|$ for $|\cdot|_q$. Recall that

$$\mathcal{S}_{\Pi_{\underline{Q},q}}^* = I_q(\phi_q^* \otimes \tilde{\phi}_q^*) \cdot B_{\Pi_q} \cdot \frac{\zeta_q(1)^2}{|N|^2 \zeta_q(2)^2} \cdot \omega_{F,q}^{-1}(\mathbf{d}_f) |\mathbf{d}_F^{\kappa}|.$$

Here $\mathbf{d}_F^{\kappa} = \mathbf{d}_f^{\kappa_1} \mathbf{d}_g^{\kappa_2} \mathbf{d}_h^{\kappa_3}$ is a product of the adjustment of levels defined in §3.4. Let Frob_q be the geometric Frobenius element in the Weil group $W_{\mathbf{Q}_q}$.

Case (Ia) and (Ib): Suppose we are in the situation of either §6.2 or §6.3. Then we have $v_q(\mathbf{d}_f) = 0$, $v_q(\mathbf{d}_g) = c^* - c_2$ and $v_q(\mathbf{d}_h) = c^* - c_3$. Thus

$$\omega_{F,q}^{-1}(\mathbf{d}_f) |\mathbf{d}_F^{\kappa}| = |q|^{\kappa_2(c^* - c_2) + \kappa_3(c^* - c_3)} \quad (\kappa_i = k_i - 2).$$

In Case (Ia) with $c_3 = 0$, by Proposition 6.3 we obtain

$$\mathcal{S}_{\Pi_q}^* = \omega_2 \omega_3 (q^{-c_2}) \varepsilon(1/2, \pi_2)^2 |q|^{(\kappa_3 - 2)c_2}.$$

Hence, we find that $I_{\mathbf{F},q} = \det \rho_{\mathbf{g}} \det \rho_{\mathbf{h}} (\text{Frob}_q^{c^*}) |q|_{-2c_2} \cdot \varepsilon(\mathbf{g})^2$. Consider Case (Ia) with $c_3 > 0$ ($c^* = c_2$). Let $\alpha_q^*(\mathbf{h}) : W_{\mathbf{Q}_q} \rightarrow \mathbf{I}^\times$ be the unramified character sending Frob_q to $\mathbf{a}(q, \mathbf{h})$ and let $\alpha_q(\mathbf{h}) = \mathbf{a}(q, \mathbf{h}) := \chi_3 |\cdot|^{-\frac{1-k_3}{2}}(q)$. By local Langlands correspondence for $\text{GL}(2)$,

$$\varepsilon(\text{WD}_q(V_f \otimes V_g)) = \varepsilon\left(\frac{2 - k_1 - k_2}{2}, \pi_f \otimes \pi_g\right).$$

This implies that

$$\varepsilon(1/2, \pi_1 \otimes \pi_2 \otimes \chi_3) = \varepsilon(\text{WD}_q(V_f \otimes V_g) \otimes \alpha_q^*(\mathbf{h}) \chi_F).$$

By Proposition 6.3 and (6.6), we thus obtain

$$\mathcal{S}_{\Pi_q}^* = \varepsilon(\text{WD}_q(V_f \otimes V_g) \otimes \alpha_q^*(\mathbf{h}) \chi_{\mathbf{F}\underline{Q}}) \cdot \alpha_q(\mathbf{h})^{-2c^*} |q|^{2c^*} \cdot \det V_h(\text{Art}_q(-1)) \cdot \varepsilon(V_h)^2.$$

Here $\text{Art} : \mathbf{Q}_q^\times \rightarrow W_{\mathbf{Q}_q}^{ab}$ is the Artin map. Therefore, by Lemma 6.11 we find that

$$I_{\mathbf{F},q} = \varepsilon(\mathbf{f} \otimes \mathbf{g}) \cdot \alpha_q^*(\mathbf{h}) \chi_{\mathbf{F}\underline{Q}}(\text{Frob}_q^{c'}) \cdot \alpha_q^*(\mathbf{h}) \varepsilon_{\text{cyc}}(\text{Frob}_q^{-2c^*}) \det \rho_{\mathbf{h}}(\text{Art}_q(-1)) \cdot \varepsilon(\mathbf{h})^2,$$

where c' is the exponent of the conductor of $\pi_{f,q} \times \pi_{g,q}$. In case (Ib) with $L(s, \pi_2 \otimes \pi_3) = 1$, we see from Proposition 6.6 that

$$\begin{aligned} \mathcal{S}_{\Pi_q}^* &= \varepsilon(\text{WD}_p(V_g \otimes V_h) \otimes \alpha_q^*(f) \chi_{\mathbf{F}\underline{Q}}) \cdot \alpha_q(f)^{2c^*} \chi_F(q^{c^*}) \\ &\quad \times \varepsilon(\text{WD}_q(V_g))^2 \varepsilon(\text{WD}_q(V_h))^2 \cdot |q|^{2(c_2 + c_3 - 2c^*)}. \end{aligned}$$

It follows that

$$I_{\mathbf{F},q} = \varepsilon(\mathbf{g} \otimes \mathbf{h}) \cdot \alpha_q^*(\mathbf{f})^2 \chi_{\mathbf{F}}(\text{Frob}_q^{c^*}) \cdot \alpha_q^*(\mathbf{f}) \chi_{\mathbf{F}}(\text{Frob}_q^{c''}) |q|^{2(c_2 + c_3 - 2c^*)},$$

where c'' is the exponent of the conductor of $\pi_{g,q} \times \pi_{h,q}$. If $L(s, \pi_{g,q} \otimes \pi_{h,q}) \neq 1$, then $\mathcal{S}_{\Pi_q}^* = 2 |q|^{-1}$.

We proceed to treat Case (IIa) and (IIb). So $\pi_{f,q}$ is principal series while $\pi_{g,q}$ and $\pi_{h,q}$ are discrete series.

Case (IIa): In the setting of §6.4, we have $v_q(\mathbf{d}_f) = r = \lceil \frac{c^*}{2} \rceil$ and $v_q(\mathbf{d}_g) = v_q(\mathbf{d}_h) = 0$; then

$$\omega_{F,q}^{-1}(\mathbf{d}_f) |\mathbf{d}_F^{\kappa}|_q = \omega_{F,q}(q^{-r}) |q|^{\kappa_1 r}.$$

By Proposition 6.9 and (6.6), we find that

$$\mathcal{S}_{\Pi_q}^* = \varepsilon(\text{WD}_q(V_g \otimes V_h) \otimes \alpha_{f,q}^* \chi_F) \cdot \alpha_{f,q}^*(\text{Frob}_q^{-2r}).$$

Case (IIb): In the setting of §6.5, we have $v_q(\mathbf{d}_f) = c^* - c_1$ and $v_q(\mathbf{d}_g) = v_q(\mathbf{d}_h) = 0$. Then

$$\omega_{F,q}^{-1}(\mathbf{d}_f) |\mathbf{d}_F^{\kappa}|_q = \omega_{F,q}(q^{c_1 - c^*}) |q|^{\kappa_1(c^* - c_1)}.$$

If $c_1 > 0$, we set $\alpha_q(\mathbf{f}) := \mathbf{a}(q, \mathbf{f})$. If $c_1 = 0$, then set $\alpha_q(\mathbf{f}) := \mathbf{a}(q, \mathbf{f}) - \beta_q(\mathbf{f})$, where $\beta(q, \mathbf{f})$ is a root of the Hecke polynomial of \mathbf{f} at q fixed in Definition 3.3. Define $\alpha_{\mathbf{f},q}^* : W_{\mathbf{Q}_q} \rightarrow \mathbf{I}_1^\times$ to be the unramified character with $\alpha_{\mathbf{f},q}^*(\text{Frob}_q) = \alpha_q(\mathbf{f})$. By definition, $\text{Rec}_{\mathbf{Q}_q}(\chi_1 \omega_F^{1/2} \cdot | \cdot |^{\frac{1-k_{\mathbf{Q}_1}}{2}}) = \alpha_{\mathbf{f},q}^*$ the specialization of $\alpha_{\mathbf{f},q}^*$ at Q_1 . From Proposition 6.10, we obtain the following expression of $\mathcal{S}_{\Pi_q}^*$:

$$\begin{aligned} \mathcal{S}_{\Pi_q}^* &= \varepsilon(\text{WD}_q(V_g \otimes V_h) \otimes \alpha_{\mathbf{f},q}^* \chi_F) \cdot \alpha_{\mathbf{f},q}^*(\text{Frob}_q^{-2c^*}) \cdot \varepsilon(\text{WD}_q(V_f))^2 |q|^{2c_1} \\ &\quad \times \det V_f(\text{Art}_q(-1)) \begin{cases} (1 + |q|) & \text{if } \pi_2 \text{ is of type } \mathbf{1} \\ 1 & \text{if } \pi_2 \text{ is of type } \mathbf{2} \end{cases}. \end{aligned}$$

In either case, it is easy to see by Lemma 6.11 that

$$I_{\mathbf{F},q} = \varepsilon_q(\mathbf{f} \otimes \mathbf{g}) \cdot \alpha_{\mathbf{f},q}^* \chi_F(\text{Frob}_q^{c'}) \cdot \alpha_{\mathbf{f},q}^*(\text{Frob}_q^{-2c^*}) \cdot \varepsilon_q(\mathbf{f})^2 \det \rho_{\mathbf{f}}(\text{Art}_q(-1)) |q|^{2c_1},$$

where c' is the exponent of the conductor of $\pi_{\mathbf{f},q} \times \pi_{\mathbf{g},q}$. This completes the proof in all cases. \square

7. THE INTERPOLATION FORMULAE

7.1. Proof of the main results. We complete the proofs of the main results in this section. We retain the notation in the introduction. For $\underline{Q} = (Q_1, Q_2, Q_3)$, recall that $\omega_{\underline{Q}}^{1/2} = \omega^{a - \frac{w_{\underline{Q}} - 3}{2}} \epsilon_{Q_1}^{1/2} \epsilon_{Q_2}^{1/2} \epsilon_{Q_3}^{1/2}$ and that

$$\Pi_{\underline{Q}} := \pi_{\mathbf{f}_{Q_1}} \times \pi_{\mathbf{g}_{Q_2}} \times \pi_{\mathbf{h}_{Q_3}} \times \omega_{\underline{Q}}^{-1/2}.$$

In terms of L -functions attached to Galois representations in the introduction, we have

$$L(s + \frac{1}{2}, \Pi_{\underline{Q}}) = \Gamma_{\mathbf{V}_{\underline{Q}}^\dagger}(s) \cdot L(\mathbf{V}_{\underline{Q}}^\dagger, s),$$

where $\Gamma_{\mathbf{V}_{\underline{Q}}^\dagger}(s) = L(s + \frac{1}{2}, \Pi_{\underline{Q}, \infty})$ is the Γ -factor of $\mathbf{V}_{\underline{Q}}^\dagger$ in (1.4). The set Σ^- in Definition 3.9 is given by

$$\Sigma^- = \left\{ \ell \mid N \mid \varepsilon(\text{WD}_\ell(\mathbf{V}_{\underline{Q}}^\dagger)) = -1 \text{ for some } \underline{Q} \in \mathfrak{X}_{\mathcal{R}}^+ \right\}.$$

Theorem 7.1. *Suppose that $p > 3$, (ev) and (sf) hold. After enlarging the coefficient ring \mathcal{O} to some finite unramified extension over \mathcal{O} , the following statements hold.*

- (1) *If $\Sigma^- = \emptyset$ and \mathbf{f} satisfies the Hypothesis (CR), then there exists an element $\mathcal{L}_{\mathbf{F}}^{\mathbf{f}} \in \mathcal{R}$ such that for every $\underline{Q} = (Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathcal{R}}^{\mathbf{f}}$ in the unbalanced range dominated by \mathbf{f} , we have*

$$(\mathcal{L}_{\mathbf{F}}^{\mathbf{f}}(\underline{Q}))^2 = \Gamma_{\mathbf{V}_{\underline{Q}}^\dagger}(0) \cdot \frac{L(\mathbf{V}_{\underline{Q}}^\dagger, 0)}{(-1)^{k_{Q_1}} \Omega_{\mathbf{f}_{Q_1}}^2} \cdot \mathcal{E}_p(\text{Fil}_{\mathbf{f}}^+ \mathbf{V}_{\underline{Q}}) \cdot \prod_{\ell \in \Sigma_{\text{exc}}} (1 + \ell^{-1}),$$

where the period $\Omega_{\mathbf{f}_{Q_1}}$ is the period attached to the p -stabilized form \mathbf{f}_{Q_1} as in Definition 4.12.

- (2) *If $\#\Sigma^-$ is odd, \mathbf{f}, \mathbf{g} and \mathbf{h} all satisfy Hypothesis (CR, Σ^-), and N^- and N/N^- are relatively prime, there exists a unique element $\mathcal{L}_{\mathbf{F}}^{\text{bal}} \in \mathcal{R}$ such that for any arithmetic point $\underline{Q} \in \mathfrak{X}_{\text{bal}}$ in the balanced range, we have*

$$(\mathcal{L}_{\mathbf{F}}^{\text{bal}}(\underline{Q}))^2 = \Gamma_{\mathbf{V}_{\underline{Q}}^\dagger}(0) \cdot \frac{L(\mathbf{V}_{\underline{Q}}^\dagger, 0)}{\Omega_{\mathbf{f}_{Q_1}^D} \Omega_{\mathbf{g}_{Q_2}^D} \Omega_{\mathbf{h}_{Q_3}^D}} \cdot \mathcal{E}_p(\text{Fil}_{\text{bal}}^+ \mathbf{V}_{\underline{Q}}) \cdot \prod_{\ell \in \Sigma_{\text{exc}}} (1 + \ell^{-1}).$$

PROOF. By the observation in Remark 6.2, there exists Dirichlet characters $\underline{\chi} = (\chi_1, \chi_2, \chi_3)$ modulo M with $M^2 \mid N$ such that $\chi_1 \chi_2 \chi_3 = 1$ and the triple

$$\mathbf{F}' := (\mathbf{f} \otimes \chi_1, \mathbf{g} \otimes \chi_2, \mathbf{h} \otimes \chi_3)$$

of primitive Hida families attached to the Dirichlet twists $(\mathbf{f} | [\chi_1], \mathbf{g} | [\chi_2], \mathbf{h} | [\chi_3])$ satisfies Hypothesis 6.1 at all classical points. Enlarging \mathcal{O} if necessary, we may choose a square root $\sqrt{I_{\mathbf{F}'}} \in \mathcal{R}^\times$ of $I_{\mathbf{F}'} := \prod_{q \mid N/N^-} I_{\mathbf{F}',q}$

defined in Proposition 6.12. On the other hand, by Proposition 7.5 and Proposition 7.7 in the next subsection, there exist $u_1 \in \mathbf{I}_1^\times$ and $u_2 \in \mathcal{R}^\times$ such that for all arithmetic points $\underline{Q} \in \mathfrak{X}_{\mathcal{R}}^+$, we have the equalities

$$\begin{aligned}\Omega_{(\mathbf{f} \otimes \chi_1)_{Q_1}} &= u_1(Q_1) \cdot \Omega_{\mathbf{f}_{Q_1}}; \\ \Omega_{\mathbf{f}_{Q_1}^D \otimes \chi_1} \Omega_{\mathbf{g}_{Q_2}^D \otimes \chi_2} \Omega_{\mathbf{h}_{Q_3}^D \otimes \chi_3} &= u_2^2(\underline{Q}) \cdot \Omega_{\mathbf{f}_{Q_1}^D} \Omega_{\mathbf{g}_{Q_2}^D} \Omega_{\mathbf{h}_{Q_3}^D}.\end{aligned}$$

Now we define

$$\mathcal{L}_{\mathbf{F}}^{\mathbf{f}} := \mathcal{L}_{\mathbf{F}'}^{\mathbf{f} \otimes \chi_1} \cdot 2\sqrt{I_{\mathbf{F}'}}^{-1} \cdot u_1; \quad \mathcal{L}_{\mathbf{F}}^{\text{bal}} := \Theta_{\mathbf{F}'D^*} \cdot 2^{-\frac{\#\Sigma^- + 1}{2}} \sqrt{I_{\mathbf{F}'}}^{-1} \cdot u_2.$$

Then we can verify directly that $\mathcal{L}_{\mathbf{F}}^{\mathbf{f}}$ (resp. $\mathcal{L}_{\mathbf{F}}^{\text{bal}}$) enjoys the desired interpolation formulae by Corollary 3.13 (resp. Corollary 4.13) combined with Proposition 6.12, the p -adic computation Proposition 5.4 (resp. Proposition 5.6) and Remark 5.7. \square

Remark 7.2. The reason for the appearance of the fudge factor $\prod_{\ell \in \Sigma_{\text{exc}}} (1 + \ell^{-1})$ is not clear to the author, but a similar factor H_0 appeared in p -adic L -functions for adjoint representations [Hid88a, Corollary 7.12].

7.2. The comparison between the canonical periods of Hida families with twists. Let $\mathbf{f} \in e\mathbf{S}(N, \psi, \mathbf{I})$ be a primitive Hida family of tame conductor N and of brach character ψ . We assume that \mathbf{f} satisfies (CR). Let $Q \in \mathfrak{X}_{\mathbf{I}}^+$ be an arithmetic point. Let $\mathbf{f}_Q \in e\mathcal{S}_{k_Q}(Np^*, \psi\omega^{2-k_Q}\epsilon_Q)$ be the specialization of \mathbf{f} at Q and $\pi = \pi_{\mathbf{f}_Q}$ be the unitary cuspidal automorphic representation of $\text{GL}_2(\mathbf{A})$ associated with \mathbf{f}_Q . Let $q \neq p$ be a prime number such that π is *minimal* at q . Let χ be a Dirichlet character modulo a power of q . Let

$$\mathbf{f}^\# = \mathbf{f} \otimes \chi$$

be the primitive Hida family corresponding to the twist $\mathbf{f}|\chi$ and let $N^\#$ be the tame conductor of $\mathbf{f}^\#$. We assume that $\pi \otimes \chi$ is not minimal at q . Namely,

$$N^\# = Nq^t \text{ for some } t > 0.$$

The goal of this subsection is to show the two periods $\Omega_{\mathbf{f}_Q}$ and $\Omega_{\mathbf{f}_Q^\#}$ defined in Definition 3.12 are equal up to a unit in \mathbf{I} by the method of level-raising. We will also prove the same result for the periods of the primitive Jacquet-Langlands lifts \mathbf{f}^D and $\mathbf{f}^{\#D}$.

Remark 7.3. We recall some generalities on congruence ideals following the discussion in [Hid88a, page 363-366]. Let R be a domain. Let T be a finite reduced R -algebra with a R -algebra homomorphism $\lambda : T \rightarrow R$. For any T -module M , we denote

$$M[\lambda] := \{x \in M \mid rx = 0 \text{ for all } r \in \text{Ker } \lambda\}.$$

Then

$$C(\lambda) := \lambda(T[\lambda]) = \lambda(\text{Ann}_T(\text{Ker } \lambda)).$$

Let H be a free T -module of rank d . Suppose that T is Gorenstein, i.e. $T \simeq \text{Hom}_R(T, R)$ as T -modules and that we have a perfect pairing $\langle \cdot, \cdot \rangle : H \times H \rightarrow R$ such that $\langle tx, y \rangle = \langle x, ty \rangle$ for $t \in T$. Then $T[\lambda]$ is free R -module of rank one and hence $H[\lambda]$ is free R -module of rank d with a basis $\{e_1, \dots, e_d\}$. We have

$$C(\lambda)^d = (\det \langle e_i, e_j \rangle).$$

Lemma 7.4. *We have $C(\mathbf{f}^\#) = C(\mathbf{f}) \cdot E_q(\mathbf{f})$, where*

$$E_q(\mathbf{f}) = \begin{cases} (q-1)(\mathbf{a}(q, \mathbf{f})^2 - \psi_{\mathbf{I}}(q)(1+q)^2) & \text{if } q \nmid N, \\ 1 - q^{-1} & \text{if } \pi_q \text{ is a ramified principal series,} \\ 1 - q^{-2} & \text{if } \pi_q \text{ is unramified special,} \\ 1 & \text{if } \pi_q \text{ is supercuspidal} \end{cases}$$

(recall that $\psi_{\mathbf{I}}$ is the \mathbf{I} -adic character $\psi \langle \epsilon_{\text{cyc}} \rangle^{-2} \langle \epsilon_{\text{cyc}} \rangle_{\mathbf{I}}$).

PROOF. We shall follow the notation in §3.3. Let $\mathbf{T}^\# := \mathbf{T}(N^\#, \mathbf{I})$ and let $\mathfrak{m}^\#$ be the maximal ideal of $\mathbf{T}^\#$ containing the operator U_q , $\{T_q - \mathbf{a}(q, \mathbf{f})\}_{q \nmid Np}$ and $\{U_q - \mathbf{a}(q, \mathbf{f})\}_{q \mid Np, q \neq q}$. The twisting morphism $[[\chi^{-1}]$ induces an isomorphism

$$[[\chi^{-1}] : e\mathbf{S}(N^\#, \mathbf{I})_{\mathfrak{m}_{\mathbf{f}^\#}} \simeq e\mathbf{S}(N^\#, \mathbf{I})_{\mathfrak{m}^\#}[U_q = 0]$$

as \mathbf{T}^\sharp -modules. Let $r_0 = 2$ if $q \nmid N$, $r_0 = 1$ if $q \mid N$ and π_q is not supercuspidal and $r_0 = 0$ if π_q is supercuspidal. For brevity, we put

$$\mathbf{S}(Nq^r) := e\mathbf{S}(Nq^r, \mathbf{I})_{\mathfrak{m}^\sharp} \otimes_{\mathbf{I}} \text{Frac } \mathbf{I} \text{ for } r \in \mathbf{Z}_{\geq 0}.$$

According to the possible list of tame conductors of newforms in $e\mathbf{S}(Nq^r, \mathbf{I})_{\mathfrak{m}^\sharp}$ [DT94, page 436], all newforms in $e\mathbf{S}(Nq^r, \mathbf{I})_{\mathfrak{m}^\sharp}$ have tame conductor dividing Nq^{r_0} . It follows that $U_q = 0$ on $\mathbf{S}(Nq^{r_0})$ and that

$$\mathbf{S}(Nq^{r+1}) = \mathbf{S}(Nq^r) \oplus V_q \mathbf{S}(Nq^r) \text{ if } r \geq r_0.$$

Here recall that $V_q(\sum \mathfrak{a}_n q^n) = q \sum \mathfrak{a}_n q^{qn}$. Combined with the relation $U_q V_q = q$, the above facts implies that

$$e\mathbf{S}(N^\sharp, \mathbf{I})_{\mathfrak{m}_{\mathbf{f}^\sharp}} \otimes_{\mathbf{I}} \text{Frac } \mathbf{I} \simeq \mathbf{S}(Nq^t)[U_q = 0] = \mathbf{S}(Nq^{r_0})[U_q = 0] = \mathbf{S}(Nq^{r_0}).$$

and hence

$$(7.1) \quad \mathbf{T}(N^\sharp, \mathbf{I})_{\mathfrak{m}_{\mathbf{f}^\sharp}} \simeq \mathbf{T}_{\mathfrak{m}^\sharp}^\sharp = \mathbf{T}(Nq^{r_0}, \mathbf{I})_{\mathfrak{m}^\sharp}.$$

We are going to apply the discussion in Remark 7.3 to compare the congruence ideals. For each positive integer M not divisible by p , put

$$\mathbf{H}_p(M) = \varprojlim_{n \rightarrow \infty} \mathbf{H}_{\text{ét}}^1(X_1(Mp^n)/\overline{\mathbf{Q}}, \mathbf{Z}_p) \otimes_{\mathbf{Z}_p} \mathcal{O}.$$

Let $\{, \}_M : \mathbf{H}_p(M) \times \mathbf{H}_p(M) \rightarrow \Lambda$ denote the Hecke-equivariant perfect pairing defined in [Oht95, Definition (4.1.17)]. Let $\mathbf{H}_p(M)_{\mathfrak{m}} := (\mathbf{H}_p(M) \otimes_{\Lambda} \mathbf{I})_{\mathfrak{m}}$. By [Wil95, Corollary 1 and 2, page 482], $\mathbf{H}_p(M)_{\mathfrak{m}}$ is a free $\mathbf{T}(M, \mathbf{I})_{\mathfrak{m}}$ -module of rank two and $\mathbf{T}(M, \mathbf{I})_{\mathfrak{m}}$ is Gorenstein under the Hypothesis (CR). Let $\mathbf{H} = \mathbf{H}_p(N)_{\mathfrak{m}}$ and $\mathbf{H}^\sharp = \mathbf{H}_p(Nq^{r_0})_{\mathfrak{m}^\sharp}$. Suppose that we have an injective \mathbf{I} -linear map $i_q : \mathbf{H} \rightarrow \mathbf{H}^\sharp$ such that

- (i) $i_q(\mathbf{H}[\lambda_{\mathbf{f}}]) \subset \mathbf{H}^\sharp[\lambda_{\mathbf{f}^\sharp}]$;
- (ii) the \mathbf{I} -submodule $i_q(\mathbf{H})$ is a direct summand of \mathbf{H}^\sharp .

Let i_q^* be the adjoint map of i_q . Recall that $i^* : \mathbf{H}^\sharp \rightarrow \mathbf{H}$ is the unique map such that $\{i_q(x), y\}_{Nq^{r_0}} = \{x, i_q^*(y)\}_N$. We have

$$(7.2) \quad C(\mathbf{f}^\sharp)^2 = C(\mathbf{f})^2 \det(i_q^* i_q|_{\mathbf{H}[\lambda_{\mathbf{f}}]}).$$

We proceed to construct the map i_q and compute the composition $i_q i_q^*$. Let $\lambda = \lambda_{\mathbf{f}}$. For an integer d relatively prime to Np , S_d denotes the Hecke operator $[\Gamma_N \begin{pmatrix} d & 0 \\ 0 & d \end{pmatrix} \Gamma_N]$. Then we have $S_d = \sigma_d \langle d \rangle_{\mathbf{I}} \langle d \rangle^{-2} \in \mathbf{T}$, where σ_d is the diamond operator.

Case $q \nmid N$ ($r_0 = 2$): Define $i_q : \mathbf{H} \rightarrow \mathbf{H}^\sharp$ by

$$i_q(x) = qx - V_q T_q x - S_q V_q^2 x.$$

Then one verifies directly that $U_q i_q = 0$, which implies (i). The property (ii) is a consequence of Ihara's lemma [Rib84, Theorem 4.1]. A direct computation shows that

$$i_q^* = q[\Gamma_N \Gamma_{Nq}] - S_q^{-1} T_q [\Gamma_N \begin{pmatrix} q & 0 \\ 0 & 1 \end{pmatrix} \Gamma_{Nq}] + S_q^{-1} [\Gamma_N \begin{pmatrix} q^2 & 0 \\ 0 & 1 \end{pmatrix} \Gamma_{Nq}],$$

and hence $i_q^* i_q|_{\mathbf{H}[\lambda]}$ is a scalar given by

$$i_q^* i_q|_{\mathbf{H}[\lambda]} = \lambda(S_q)^{-1} q(1-q)(\lambda(T_q)^2 - (1+q)^2 \lambda(S_q)).$$

Note that $\lambda(S_q) = \psi_{\mathbf{I}}(q)$.

Case $q \mid N$ ($r_0 = 1$): Define $i_q : \mathbf{H} \rightarrow \mathbf{H}^\sharp$ by

$$i_q(x) = x - q^{-1} V_q \mathbf{U}_q x.$$

A direct computation shows that the adjoint map i_q^* is given by

$$i_q^* = [\Gamma_N \begin{pmatrix} q & 0 \\ 0 & 1 \end{pmatrix} \Gamma_{Nq}] - q^{-1} \mathbf{U}_q [\Gamma_N \Gamma_{Nq}]$$

and that

$$i_q^* i_q = -q^{-1} \left([\Gamma_N \begin{pmatrix} q & 0 \\ 0 & 1 \end{pmatrix} \Gamma_{Nq}] V_q - q^{-1} \mathbf{U}_q [\Gamma_N \Gamma_{Nq}] V_q \right) \mathbf{U}_q.$$

Let $s = v_q(N)$ and $\tau_{q^s} := \begin{pmatrix} 0 & 1 \\ -q^s & 0 \end{pmatrix} \in \mathrm{GL}_2(\mathbf{Q}_q)$. It is easy to see that

$$\tau_{q^s} \cdot [\Gamma_N \Gamma_{Nq}] V_q \cdot \tau_{q^s}^{-1} = S_q^{-1} \cdot \mathbf{U}_q.$$

The restriction of $[\Gamma_N \Gamma_{Nq}] V_q$ to $\mathbf{H}[\lambda]$ is given by

$$\lambda(S_q)^{-1} \cdot (\tau_{q^s}^{-1} \mathbf{U}_q \tau_{q^s} |_{\mathbf{H}[\lambda_f]}) = \lambda(\mathbf{U}_q)^{-1} \cdot \begin{cases} q & \text{if } \pi_{f,q} \text{ is a ramified principal series,} \\ 1 & \text{if } \pi_{f,q} \text{ is unramified special.} \end{cases}$$

We thus find that

$$i_q^* i_q |_{\mathbf{H}[\lambda]} = -q^{-1} \lambda(\mathbf{U}_q) (q - q^{-1} \lambda(\mathbf{U}_q) \lambda(S_q^{-1}) (\tau_q^{-1} \mathbf{U}_q \tau_q |_{\mathbf{H}[\lambda]})) = -\lambda(\mathbf{U}_q) E_q(1, \mathrm{Ad} \rho_f).$$

The assertion follows from (7.2) and the above computation of $i_q^* i_q |_{\mathbf{H}[\lambda]}$. \square

Proposition 7.5. *There exists a unit $u \in \mathbf{I}^\times$ such that for any arithmetic point Q , we have*

$$\Omega_{f_Q} = u(Q) \cdot \Omega_{f_Q^\sharp}.$$

PROOF. Let f_Q° and $f_Q^{\sharp\circ}$ be the newforms corresponding to f_Q and f_Q^\sharp of conductors Np^n and $N^\sharp p^n$ respectively. From (2.18), we see that

$$\frac{\|f_Q^{\sharp\circ}\|_{\Gamma_0(N^\sharp p^n)}^2}{\|f_Q^\circ\|_{\Gamma_0(Np^n)}^2} = \frac{[\mathrm{SL}_2(\mathbf{Z}) : \Gamma_0(N^\sharp)]}{[\mathrm{SL}_2(\mathbf{Z}) : \Gamma_0(N)]} \cdot \frac{\varepsilon(1/2, \pi_q) B_{\pi_q \otimes \chi_q}}{\varepsilon(1/2, \pi_q \otimes \chi_q) B_{\pi_q}}.$$

A direct computation shows that if $q \nmid N$, then the right hand side equals

$$\frac{N^\sharp}{N} \cdot L(1, \pi_q, \mathrm{Ad})^{-1} = q^{-3} \cdot (\psi_{\mathbf{I}}^{-1}(q) E_q(\mathbf{f}))(Q),$$

and if $q \mid N$, then it is equal to

$$\frac{N^\sharp}{N} \begin{cases} 1 - q^{-1} & \text{if } q \mid N \text{ and } \pi_q \text{ is a ramified principal series,} \\ 1 - q^{-2} & \text{if } \pi_q \text{ is special,} \\ 1 & \text{if } \pi_q \text{ is supercuspidal.} \end{cases}$$

In any case, it is clear that there exists a unit $u' \in \mathbf{I}^\times$ such that

$$\frac{\|f_Q^{\sharp\circ}\|_{\Gamma_0(N^\sharp p^n)}^2}{\|f_Q^\circ\|_{\Gamma_0(Np^n)}^2} = u'(Q) \cdot E_q(\mathbf{f})(Q)$$

for all arithmetic points Q . Therefore, the assertion follows from the definitions of periods in Definition 3.12 and Lemma 7.4 by noting that $\mathcal{E}_p(f_Q, \mathrm{Ad}) = \mathcal{E}_p(f_Q^\sharp, \mathrm{Ad})$. \square

The definite case. Now we consider periods of definite quaternionic Hida families. Assume that \mathbf{f} satisfies Hypothesis (CR, Σ^-). Let $\mathbf{f}^D \in e\mathbf{S}^D(N, \psi, \mathbf{I})$ be the primitive Jacquet-Langlands lift of \mathbf{f} . Let q^c be the conductor of χ . Let \mathcal{P}_χ be the element in the group ring $\mathcal{O}[\mathrm{GL}_2(\mathbf{Q}_q)]$ defined by

$$\mathcal{P}_\chi := \mathfrak{g}(\chi^{-1})^{-1} \sum_{b \in (\mathbf{Z}_q/q^c \mathbf{Z}_q)^\times} \chi(b) \cdot \begin{pmatrix} 1 & bq^{-c} \\ 0 & 1 \end{pmatrix},$$

where $\mathfrak{g}(\chi^{-1})$ is the Gauss sum. We define

$$\mathbf{f}^D |[\chi](x) := \mathcal{P}_\chi(\mathbf{f}^D)(x) \chi(\nu(x)) \in e\mathbf{S}^D(Nq^{2c}, \psi\chi^2, \mathbf{I}).$$

Lemma 7.6. *The quaternionic form $\mathbf{f}^D |[\chi]$ is a primitive Jacquet-Langlands lift of \mathbf{f}^\sharp . In other words, $\mathbf{f}^D |[\chi] \in e\mathbf{S}^D(N^\sharp, \psi\chi^2, \mathbf{I})[\lambda_{f^\sharp}]$ is a generator over \mathbf{I} .*

PROOF. We first note that $\mathbf{U}_q(\mathbf{f}^D|[\chi]) = 0$ as χ is ramified. Since $\pi_q \otimes \chi_q$ is not minimal, the operator \mathbf{U}_q is nilpotent on $\mathcal{V}_{\pi_q \otimes \chi_q}^N(\mathbf{Z}_q)$ and $\dim_{\mathbf{C}} \mathcal{V}_{\pi_q \otimes \chi_q}^N(\mathbf{Z}_q)[\mathbf{U}_q = 0] = 1$. This implies that $\mathbf{f}^D|[\chi] \in e\mathbf{S}^D(N^\sharp, \psi\chi^2, \mathbf{I})[\lambda_{\mathbf{f}_\chi^D}]$.

To see that $\mathbf{f}^D|[\chi]$ is primitive, it suffices to show that $\mathbf{f}^D|[\chi]$ is non-vanishing modulo the maximal ideal $\mathfrak{m}_{\mathbf{I}}$ of \mathbf{I} . Let $f' := \mathbf{f}^D \otimes \chi \circ \nu \pmod{\mathfrak{m}_{\mathbf{I}}} \in \mathcal{S}_2^D(N^\sharp p^t, \psi\chi^2, \overline{\mathbb{F}}_p)$ for some positive integer t . Define two operators on $\mathcal{S}_2^D(N^\sharp p^t, \psi\chi^2, \overline{\mathbb{F}}_p)$ by

$$L_1 = \sum_{a \in \mathbf{Z}_q/q^c \mathbf{Z}_q} \psi_{\mathbf{Q}_q}(aq^{-c}) \rho\left(\begin{pmatrix} 1 & aq^{-c} \\ 0 & 1 \end{pmatrix}\right); \quad L_2 = \sum_{b \in (\mathbf{Z}_q/q^c \mathbf{Z}_q)^\times} \chi^{-1}(b) \rho\left(\begin{pmatrix} b & 0 \\ 0 & 1 \end{pmatrix}\right).$$

Then

$$\begin{aligned} L_2 L_1(\mathbf{f}^D|[\chi] \pmod{\mathfrak{m}_{\mathbf{I}}}) &= \sum_b \sum_a \psi_{\mathbf{Q}_q}(abq^{-c}) \rho\left(\begin{pmatrix} 1 & aq^{-c} \\ 0 & 1 \end{pmatrix}\right) f' \\ &= q^c f' - q^{c-1} \sum_{b \in \mathbf{Z}_q/q \mathbf{Z}_q} \rho\left(\begin{pmatrix} 1 & bq^{-1} \\ 0 & 1 \end{pmatrix}\right) f' \\ &= q^{c-1} \left(q - \begin{pmatrix} q^{-1} & 0 \\ 0 & 1 \end{pmatrix} \mathbf{U}_q \right) f'. \end{aligned}$$

Suppose that $\mathbf{f}^D|[\chi] \pmod{\mathfrak{m}_{\mathbf{I}}} = 0$. Then we deduce from the above equation that either

$$\left(q - \begin{pmatrix} q^{-1} & 0 \\ 0 & 1 \end{pmatrix} \mathbf{a}(f, q) \chi(q) \right) f' = 0 \text{ if } q \mid N,$$

or

$$\left(q - \mathbf{a}(f, q) \begin{pmatrix} q^{-1} & 0 \\ 0 & 1 \end{pmatrix} - \psi(q) \begin{pmatrix} q^{-2} & 0 \\ 0 & 1 \end{pmatrix} \right) f' = 0 \text{ if } q \nmid N.$$

In any case, this implies that $f' = 0$ by *Ihara's lemma* for definite quaternion algebras [CH16, Lemma 5.5] and hence $\mathbf{f}^D \pmod{\mathfrak{m}_{\mathbf{I}}} = 0$, which is a contradiction. \square

Proposition 7.7. *Let $\mathbf{f}^{\sharp D}$ be a primitive Jacquet-Langlands lift of \mathbf{f}^\sharp . There exists $u \in \mathbf{I}^\times$ such that for every arithmetic point $Q \in \mathfrak{X}_{\mathbf{I}}^+$, we have*

$$\Omega_{\mathbf{f}_Q^D} = u^2(Q) \cdot \Omega_{\mathbf{f}^{\sharp D}}.$$

PROOF. Let $\mathbf{f}' := \mathbf{f}^D|[\chi]$. Then $\mathbf{f}^{\sharp D} = u \cdot \mathbf{f}'$ for some $u \in \mathbf{I}^\times$. Let $f = \mathbf{U}_p^{-n} \mathbf{f}'_Q$ and let $\varphi = \Phi(f)$ be the vector-valued automorphic form associated with f . Let $\varphi_\chi := \mathcal{P}_\chi \otimes \chi \circ \nu$. By Lemma 4.4 and (4.5), we have

$$\frac{\eta_{\mathbf{f}'}(Q)}{\eta_{\mathbf{f}^D}(Q)} = \frac{\langle \mathbf{U}_p^{-n} \mathbf{f}'_Q, \mathbf{f}'_Q \rangle_{N^\sharp p^n}}{\langle \mathbf{U}_p^{-n} \mathbf{f}'_Q, \mathbf{f}'_Q \rangle_{N p^n}} = S_1 \cdot S_2$$

where

$$S_1 := \left(\frac{N^\sharp}{N} \right)^{\frac{k_Q - 2}{2}} \cdot \frac{\text{vol}(\widehat{R}_N^\times)}{\text{vol}(\widehat{R}_{N^\sharp}^\times)}, \quad S_2 := \frac{\chi_p(p^{-n}) \langle \rho(\boldsymbol{\tau}_{N^\sharp p^n}^D) \varphi_\chi, \varphi'_\chi \rangle}{\langle \rho(\boldsymbol{\tau}_{N p^n}^D) \varphi, \varphi' \rangle}.$$

It is easy to see that

$$S_1 = \frac{[\text{SL}_2(\mathbf{Z}) : \Gamma_0(N^\sharp)] (N^\sharp)^{\frac{k_Q - 2}{2}}}{[\text{SL}_2(\mathbf{Z}) : \Gamma_0(N)] N^{\frac{k_Q - 2}{2}}}.$$

On the other hand,

$$\begin{aligned} S_2 &= \frac{\chi(\widehat{N}^\sharp) \langle \rho(\boldsymbol{\tau}_{N^\sharp p^n}^D) \mathcal{P}_\chi(\varphi), \mathcal{P}_\chi(\varphi') \rangle}{\langle \rho(\boldsymbol{\tau}_{N p^n}^D) \varphi, \varphi' \rangle} \\ &= \chi(\widehat{N}^\sharp) \cdot \frac{\langle \rho(\boldsymbol{\tau}_{N^\sharp p^n, q}^D) \mathcal{P}_\chi(W_{\pi_q}), \mathcal{P}_\chi(W_{\pi_q}) \otimes \omega_q^{-1} \rangle}{\langle \rho(\boldsymbol{\tau}_{N p^n, q}^D) W_{\pi_q}, W_{\pi_q} \otimes \omega_q^{-1} \rangle}. \end{aligned}$$

Note that $\mathcal{P}_\chi W_{\pi_q} \left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \right) = \mathbb{I}_{\mathbf{Z}_q^\times}(a) \chi_q^{-1}(a)$, so we get $\mathcal{P}_\chi W_{\pi_q} \otimes \chi_q = W_{\pi_q \otimes \chi_q}$, so we find that

$$S_2 = \chi(\widehat{N}) \cdot \frac{\langle \rho(\tau_{N^\sharp}^D) W_{\pi_q \otimes \chi}, W_{\pi_q \otimes \chi} \otimes \chi_q^{-2} \omega_q^{-1} \rangle}{\langle \rho(\tau_{N,q}) W_{\pi_q}, W_{\pi_q} \otimes \omega_q^{-1} \rangle} = \chi(\widehat{N}) \cdot \frac{B_{\pi_q \otimes \chi_q}}{B_{\pi_q}}.$$

From the above computations of S_1 and S_2 , we see that

$$\begin{aligned} \frac{\eta_{\mathbf{f}'}(Q)}{\eta_{\mathbf{f}^D}(Q)} &= \frac{\chi(\widehat{N}) \varepsilon(1/2, \pi_q \otimes \chi_q) (N^\sharp)^{\frac{k_Q-2}{2}}}{\varepsilon(1/2, \pi_q) N^{\frac{k_Q-2}{2}}} \cdot \frac{[\mathrm{SL}_2(\mathbf{Z}) : \Gamma_0(N^\sharp)]}{[\mathrm{SL}_2(\mathbf{Z}) : \Gamma_0(N)]} \frac{\varepsilon(1/2, \pi_q) B_{\pi_q \otimes \chi_q}}{\varepsilon(1/2, \pi_q \otimes \chi_q) B_{\pi_q}} \\ &= \frac{\varepsilon^{\Sigma^-}(\mathbf{f}_Q^\sharp)}{\varepsilon^{\Sigma^-}(\mathbf{f}_Q)} \cdot \frac{\|\mathbf{f}_Q^{\sharp\circ}\|_{\Gamma_0(N^\sharp)}^2}{\|\mathbf{f}_Q^\circ\|_{\Gamma_0(N)}^2} \quad (\text{by (2.18)}), \end{aligned}$$

and the lemma follows. \square

Remark 7.8. If \mathbf{f} satisfies (CR, Σ^-) , then $\eta_{\mathbf{f}^D}$ indeed generates the congruence ideal associated with the homomorphism $\lambda_{\mathbf{f}} : \mathbf{T}^D(N, \psi, \mathbf{I}) \rightarrow \mathbf{I}$. This strengthens [CH16, Prop. 6.1] by replacing (CR^+) there with a weaker hypothesis (CR, Σ^-) here. Note that $\mathbf{T}^D(N, \psi, \mathbf{I})$ is isomorphic to the N^- -new quotient of $\mathbf{T}(N, \psi, \mathbf{I})$. In particular, this implies that the congruence ideal $(\eta_{\mathbf{f}^D})$ contains $(\eta_{\mathbf{f}})$ and $(\eta_{\mathbf{f}^D}) = (\eta_{\mathbf{f}})$ if the residual Galois representation $\rho_{\mathbf{f}} \pmod{\mathfrak{m}_{\mathbf{I}}}$ is ramified at all $\ell \in \Sigma^-$. This implies Hida's canonical period of \mathbf{f} is an integral multiple of the Gross period of \mathbf{f} .

8. APPLICATIONS TO ANTICYCLOTOMIC p -ADIC L -FUNCTIONS

8.1. Primitive Hida families of CM forms. In this section, we show that when \mathbf{g} and \mathbf{h} are primitive Hida families of CM forms, then the unbalanced p -adic triple product L -function specializes to a product of theta elements á la Bertolini and Darmon in [BD96]. As a consequence, the anticyclotomic exceptional zero conjecture can be deduced from the theorem of Greenberg and Stevens. Let K be an imaginary quadratic field over \mathbf{Q} of the absolute discriminant D_K . Suppose that $p\mathcal{O}_K = \mathfrak{p}\bar{\mathfrak{p}}$, where \mathfrak{p} is the prime induced by the fixed embedding $\bar{\mathbf{Q}} \hookrightarrow \mathbf{C} \simeq \bar{\mathbf{Q}}_p$. Let K_∞ be the \mathbf{Z}_p^\times -extension of K and let $\Gamma_\infty = \mathrm{Gal}(K_\infty/K)$ be the Galois group. Let $K_{\mathfrak{p}^\infty}$ be the \mathfrak{p} -ramified \mathbf{Z}_p -extension in K_∞ and $\Gamma_{\mathfrak{p}^\infty} = \mathrm{Gal}(K_{\mathfrak{p}^\infty}/K)$ be the Galois group. For each ideal \mathfrak{a} prime to $\mathfrak{p}\mathfrak{c}$, define $\sigma_{\mathfrak{a}} \in \mathrm{Gal}(K(\mathfrak{c}\mathfrak{p}^\infty)/K)$ be the image of \mathfrak{a} under the geometrically normalized Artin map sending \mathfrak{q} to the geometric Frobenius $\mathrm{Frob}_{\mathfrak{q}}$. For each place w of K , we let $\mathrm{Art}_w : K_w^\times \rightarrow G_K^{ab}$ denote the restriction of the Artin map to K_w^\times . Then $\mathrm{Art}_{\mathfrak{p}}$ induces an embedding $\Lambda \rightarrow \mathcal{O}[[\Gamma_{\mathfrak{p}^\infty}]]$ given by $[z] \mapsto \mathrm{Art}_{\mathfrak{p}}(z)|_{K_{\mathfrak{p}^\infty}}$. Let $I_{\mathfrak{p}}^w := \mathrm{Art}_{\mathfrak{p}}(1 + p\mathbf{Z}_p)|_{K_{\mathfrak{p}^\infty}} \subset \Gamma_{\mathfrak{p}^\infty}$. Let $\mathfrak{p}^b := [\Gamma_{\mathfrak{p}^\infty} : I_{\mathfrak{p}}^w]$. Note that $b = 0$ if the class number h_K of K is prime to p . Fixing a topological generator $\gamma_{\mathfrak{p}}$ of $\Gamma_{\mathfrak{p}^\infty}$ such that $\gamma_{\mathfrak{p}}^{\mathfrak{p}^b} = \mathrm{Art}_{\mathfrak{p}}(1 + p)|_{K_{\mathfrak{p}^\infty}}$, let $l : \mathrm{Gal}(K_\infty/K) \rightarrow \mathbf{Z}_p$ be the *logarithm* defined by the equation

$$\sigma|_{K_{\mathfrak{p}^\infty}} = \gamma_{\mathfrak{p}}^{l(\sigma)}.$$

For each variable S , let $\Psi_S : \Gamma_\infty \rightarrow \mathcal{O}[[S]]^\times$ be the universal character defined by

$$\Psi_S(\sigma) = (1 + S)^{l(\sigma)}, \quad \sigma|_{K_{\mathfrak{p}^\infty}} = \gamma_{\mathfrak{p}}^{l(\sigma)}.$$

For any p -adic character $\psi : G_K \rightarrow \mathcal{O}^\times$ of conductor $\mathfrak{c}\mathfrak{p}^\infty$, we define

$$\theta_\psi(S)(q) = \sum_{(\mathfrak{a}, \mathfrak{p}\mathfrak{c})=1} \psi(\sigma_{\mathfrak{a}}) \cdot \Psi_S^{-1}(\sigma_{\mathfrak{a}}) q^{|\mathfrak{a}|} \in \mathcal{O}[[S]][[q]].$$

Let $\mathcal{V} : G_{\mathbf{Q}} \rightarrow G_K^{ab}$ be the transfer map. If ψ is geometric of weight $(1, 0)$, namely

$$\psi(\mathrm{Art}_{\mathfrak{p}}(z_1)\mathrm{Art}_{\bar{\mathfrak{p}}}(z_2)) = z_1 \text{ for } z_1, z_2 \in \mathbf{Z}_p^\times \text{ sufficiently close to } 1,$$

then $\varepsilon_{\mathrm{cyc}}^{-1} \psi \circ \mathcal{V}$ descends to a Dirichlet character, and $\theta_\psi(S)$ is a primitive Hida families in $e\mathbf{S}(C, \psi_K^\dagger, \mathcal{O}[[S]])$, where

$$C = \#(\mathcal{O}_K/\mathfrak{c})D_K; \quad \psi_K^\dagger := (\varepsilon_{\mathrm{cyc}}^{-1} \psi \circ \mathcal{V}) \cdot \tau_{K/\mathbf{Q}}.$$

Here $\tau_{K/\mathbf{Q}}$ is the quadratic character associated with K/\mathbf{Q} .

8.2. Anticyclotomic p -adic L -functions for modular forms. Let N be a positive integer relatively prime to p . Let $f \in \mathcal{S}_{2r}(Np, \mathbf{1})$ be a p -stabilized newform of weight $2r \geq 2$, tame conductor N and trivial nebentypus and let χ be a ring class character of K with the conductor $c\mathcal{O}_K$. We recall the anticyclotomic p -adic L -functions associated with (f, χ) in the definite setting. Decompose $N = N^+N^-$, where N^+ (resp. N^-) is a product of primes split (resp. non-split) in K . Suppose that

- $(Np, cD_K) = 1$,
- N^- is a square-free product of an *odd* number of primes,
- the residual Galois representation $\bar{\rho}_{f,p}$ satisfies (CR, $\text{supp } N^-$).

Let f° be the normalized newform of conductor $N^\circ = Np^{n_p}$ corresponding to f . Enlarging \mathcal{O} so that it contains all Fourier coefficients of f , let $\mathbb{T} := \mathbb{T}_{2r}(N^\circ, \mathbf{1})$ be the Hecke algebra of level $\Gamma_0(N^\circ)$ and let $\lambda_{f^\circ} : \mathbb{T} \rightarrow \mathcal{O}$ be the homomorphism induced by f° . Denote by \mathbb{T}_{N^-} be the N^- -new quotient of the \mathbb{T} . Then λ_{f° factors through \mathbb{T}_{N^-} , and we denote by λ_{f°, N^-} the resulting morphism. Let $\eta_{f^\circ} \in \mathcal{O}$ (resp. η_{f°, N^-}) be the congruence number corresponding to λ_{f° (resp. λ_{f°, N^-}). It is clear that η_{f°, N^-} is a divisor of the congruence number η_{f° of f° .

Let K_∞^- be the anticyclotomic \mathbf{Z}_p -extension of K . Let \mathbf{c} be the complex conjugation. We define the logarithm $\tilde{l} : \Gamma_\infty^- \rightarrow \mathbf{Z}_p$ by $\tilde{l}(\sigma) := l(\sigma^{1-\mathbf{c}}|_{K_{p^\infty}})$. Then the map \tilde{l} factors through the Galois group $\Gamma_\infty^- := \text{Gal}(K_\infty^-/K)$ and induces an isomorphism $\tilde{l} : \Gamma_\infty^- \simeq \mathbf{Z}_p$ as K_{p^∞} and the cyclotomic \mathbf{Z}_p -extension K_∞^+ are linearly disjoint. Let γ_- be the generator of Γ^- such that $\tilde{l}(\gamma_-) = 1$. If $\zeta \in \mu_{p^\infty}$ is a p -power root of unity, denote by $\epsilon_\zeta : \Gamma_\infty^- \rightarrow \mu_{p^\infty}$ the character defined by $\epsilon_\zeta(\gamma_-) = \zeta$. Fixing a factorization $N^+\mathcal{O}_K = \mathfrak{N}\bar{\mathfrak{N}}$, by [BD96], [CH16, Thm. A] and [Hum17, Thm. A], there exists a unique Iwasawa function $\mathcal{L}_{f/K, \chi}(W) \in \mathcal{O}[[W]]$ such that for each primitive p^n -th root of unity ζ ,

$$(8.1) \quad \begin{aligned} \mathcal{L}_{f/K, \chi}(\zeta - 1)^2 &= (2\pi)^{-2r} \Gamma(r)^2 \cdot \frac{L(f^\circ/K \otimes \chi \epsilon_\zeta, r)}{\Omega_{f^\circ, N^-}} \cdot \alpha_p(f)^{-2n} p^{(2r-1)n} \cdot \mathcal{E}_p(f, \zeta)^{2-n_p} \\ &\quad \times u_K^2 \sqrt{D_K} cD_K^{k-2} \chi \epsilon_\zeta(\sigma_{\mathfrak{N}}) \cdot \varepsilon_p(f^\circ), \end{aligned}$$

where

- $\alpha_p(f) \in \mathcal{O}^\times$ is the p -th Fourier coefficient of f ,
- $L(f^\circ/K \otimes \chi \epsilon_\zeta, s)$ is the Rankin-Selberg L -function of f° and the CM form $\theta_{\psi \epsilon_\zeta}$ attached to $\chi \epsilon_\zeta$,

$$\mathcal{E}_p(f, \zeta) := \begin{cases} (1 - \alpha_p(f)^{-1} p^{r-1} \chi(\mathfrak{p}))(1 - \alpha_p(f) p^{r-1} \chi(\bar{\mathfrak{p}})) & \text{if } \zeta = 1, \\ 1 & \text{if } \zeta \neq 1. \end{cases}$$

- Ω_{f°, N^-} is the Gross period of f° defined by

$$\Omega_{f^\circ, N^-} := 2^{2r} \cdot \|f^\circ\|_{\Gamma_0(N_{f^\circ})}^2 \cdot \eta_{f^\circ, N^-}^{-1}.$$

- $u_K = \#(\mathcal{O}_K^\times)/2$ and $\varepsilon_p(f^\circ) \in \{\pm 1\}$ is the local root number of f° at p .

When $\chi = \mathbf{1}$ is the trivial character, we write \mathcal{L}_f for $\mathcal{L}_{f, \mathbf{1}}$.

8.3. Factorization of p -adic triple product L -functions. Let $\mathbf{f} \in e\mathbf{S}(N, \omega^{k-2}, \mathbf{1})$ be the primitive Hida family passing through f at some arithmetic point Q_1 of weight $k_{Q_1} = 2r$ and trivial finite part $\epsilon_{Q_1} = 1$. Fixing $\mathbf{v} \in \bar{\mathbf{Z}}^\times$ such that $\mathbf{v}^{p^b} = 1 + p$, let $\psi = \Psi_{\mathbf{v}-1}$ be a geometric p -adic character of G_K of weight $(1, 0)$. Let $\ell \nmid Np$ be a rational prime split in K and let χ be a ring class character of conductor $\ell^m \mathcal{O}_K$ for some $m > 0$. Let $C = D_K \ell^{2m}$ and put

$$\mathbf{g} = \theta_{\psi \chi}(S_2) \in e\mathbf{S}(C, \omega^{-1} \tau_{K/Q}, \mathcal{O}[[S_2]]); \quad \mathbf{h} = \theta_{\psi \chi^{-1}}(S_3) \in e\mathbf{S}(C, \omega^{-1} \tau_{K/Q}, \mathcal{O}[[S_3]]).$$

Let $\mathbf{F} = (\mathbf{f}, \mathbf{g}, \mathbf{h})$ be the triple of primitive Hida families and let $\mathcal{L}_{\mathbf{F}}^f \in \mathcal{R} = \mathbf{I}[[S_1, S_2]]$ be the associated unbalanced p -adic L -function in Theorem 7.1 with $a = -r$ in (ev).

Proposition 8.1. *Set*

$$W_2 = \mathbf{v}^{-1}(1 + S_2)^{1/2}(1 + S_3)^{1/2} - 1; \quad W_3 = (1 + S_2)^{1/2}(1 + S_3)^{-1/2} - 1.$$

Then we have

$$\mathcal{L}_{\mathbf{F}}^f(Q_1, S_1, S_2) = \pm \mathbf{w}^{-1} \cdot \mathcal{L}_{f/K}(W_2) \cdot \mathcal{L}_{f/K, \chi^2}(W_3) \cdot \frac{\eta_{f^\circ}}{\eta_{f^\circ, N^-}} \in \mathcal{O}[[S_1, S_2]],$$

where $\mathbf{w} = \mathbf{w}(W_2, W_3)$ is a unit in $\mathcal{O}[[S_1, S_2]]$ given by

$$\mathbf{w} := u_K^2 D_K^{2r-3/2} \ell^{m/2} \psi(\sigma_{\mathfrak{N}}) \cdot \Psi_{W_1} \Psi_{W_2}(\sigma_{\mathfrak{N}}^{1-c}).$$

PROOF. For $i = 2, 3$, taking ζ_i primitive p^{n_i} -th roots of unity with $n_i > 0$, we let $Q_2 = \zeta_2 \zeta_3 \mathbf{v} - 1$ and $Q_3 = \zeta_2 \zeta_3^{-1} \mathbf{v} - 1$, so \mathbf{g}_{Q_2} and \mathbf{h}_{Q_3} are CM forms of weight one. Let $\underline{Q} = (Q_1, Q_2, Q_3)$. Let

$$\mathcal{X}_1 := \psi \Psi_{S_2}^{-1/2} \Psi_{S_3}^{-1/2} \circ \mathcal{V} : G_{\mathbf{Q}} \rightarrow \mathcal{O}[[S_1, S_2]]^\times$$

be a square root of $\det V_{\mathbf{g}} \det V_{\mathbf{h}}$. We have the decomposition of Galois representations

$$\mathrm{Ind}_K^{\mathbf{Q}} \psi \chi \Psi_{S_2}^{-1} \otimes \mathrm{Ind}_K^{\mathbf{Q}} \psi \chi^{-1} \Psi_{S_3}^{-1} \otimes \mathcal{X}_1^{-1} = \mathrm{Ind}_K^{\mathbf{Q}} \Psi_{W_2}^{c-1} \oplus \mathrm{Ind}_K^{\mathbf{Q}} \chi^2 \Psi_{W_3}^{c-1}.$$

Following the notation in the introduction, this implies that

$$\begin{aligned} \mathbf{V}_{\underline{Q}}^\dagger &= V_f(r) \otimes \mathrm{Ind}_K^{\mathbf{Q}} \epsilon_2 \oplus V_f(r) \otimes \mathrm{Ind}_K^{\mathbf{Q}} \chi^2 \epsilon_3; \\ \mathrm{Fil}_f^+ \mathbf{V}_{\underline{Q}}^\dagger &= \alpha_{f,p} \epsilon_{\mathrm{cyc}}^r \otimes (\epsilon_{2,p} \oplus \epsilon_{2,p}^{-1} \oplus \chi_p^2 \epsilon_{3,p} \oplus \chi_p^{-2} \epsilon_{3,p}^{-1}), \end{aligned}$$

where $\epsilon_i = \epsilon_{\zeta_i} : \Gamma_\infty^- \rightarrow \mu_{p^\infty}$ is the finite order character with $\epsilon_i(\gamma^-) = \zeta_i$, $i = 2, 3$. Now we explicate the items that appear in the formula of $\mathcal{L}_{\mathbf{F}}^f(\underline{Q})$ in Theorem 7.1:

- The L -values

$$\Gamma_{\mathbf{V}_{\underline{Q}}^\dagger}(0) \cdot L(\mathbf{V}_{\underline{Q}}^\dagger, s) = 4(2\pi)^{-4r} \Gamma(r)^4 \cdot L(f^\circ/K \otimes \epsilon_2, r) \cdot L(f^\circ/K \otimes \chi^2 \epsilon_3, r),$$

- By definition, ϵ_2 and ϵ_3 are of conductors $p^{n_2} \mathcal{O}_K$ and $p^{n_3} \mathcal{O}_K$, so the modified Euler factor at p is given by

$$\begin{aligned} \mathcal{E}_p(\mathrm{Fil}_f^+ \mathbf{V}_{\underline{Q}}^\dagger) &= \frac{1}{\varepsilon(r, \alpha_{f,p} \chi_p^2 \epsilon_{3,p}) \varepsilon(r, \alpha_{f,p} \chi_p^{-2} \epsilon_{3,p}^{-1}) \varepsilon(r, \alpha_{f,p} \epsilon_{2,p}) \varepsilon(r, \alpha_{f,p} \epsilon_{2,p}^{-1})} \\ &= \alpha_p(f)^{-2(n_2+n_3)} \cdot |p|^{(1-2r)(n_2+n_3)} \cdot \epsilon_{2,p}(-1) \epsilon_{3,p}(-1) \\ &= \alpha_p(f)^{-2(n_2+n_3)} \cdot |p|^{(1-2r)(n_2+n_3)}. \end{aligned}$$

- $\Omega_f = 2^{2r} \|f^\circ\|_{\Gamma_0(N^\circ)}^2 \cdot \eta_{f^\circ}^{-1}$ and $\Sigma_{\mathrm{exc}} = \emptyset$.

Comparing with the interpolation formula of Θ_f in (8.1), we find that

$$\mathcal{L}_{\mathbf{F}}^f(Q_1, \mathbf{v} \zeta_2 \zeta_3 - 1, \mathbf{v} \zeta_2 \zeta_3^{-1} - 1)^2 = \mathbf{w}(\zeta_2 - 1, \zeta_3 - 1)^{-2} \cdot \mathcal{L}_{f/K}(\zeta_2 - 1)^2 \mathcal{L}_{f/K, \chi^2}(\zeta_3 - 1)^2$$

for all non-trivial p -power roots of unity ζ_2, ζ_3 , and hence the proposition follows. \square

Remark 8.2 (An Euler system construction for $\mathcal{L}_{f/K}$). This anticyclotomic p -adic L -function $\mathcal{L}_{f/K}$ in the definite setting is constructed by using Gross points in definite quaternion algebras, and a priori there is no obvious Euler system construction. Below we explain how $\mathcal{L}_{f/K}$ can be actually recovered by the Euler system of generalized Kato classes à la Darmon and Rotger. Suppose that the weight $k_{Q_1} = 2$. In [DR17], Darmon and Rotger introduce a one-variable generalized Kato classes $\kappa(f, \mathbf{gh}) \in H^1(\mathbf{Q}, V_f \otimes V_{\mathbf{g}} \otimes_{\mathcal{O}[[S]]} V_{\mathbf{h}})$ and prove that the image of $\kappa(f, \mathbf{gh})$ under certain Perrin-Riou's big logarithm map, which we denote by \mathfrak{L} , is given by the one-variable unbalanced p -adic L -function $\mathcal{L}_{\mathbf{F}}^f(Q_1, \mathbf{v}S - 1, \mathbf{v}S - 1)$ ([DR17, Theorem 5.3]). On the other hand, in virtue of Proposition 8.1 combined with a result of Vatsal on the non-vanishing of central L -values with anticyclotomic twist, we conclude that when χ is sufficiently ramified,

$$\mathfrak{L}(\kappa(f, \mathbf{gh})) = \mathcal{L}_{\mathbf{F}}^f(Q_1, \mathbf{v}S - 1, \mathbf{v}S - 1) = \mathcal{L}_{f/K}(S) \cdot (\text{non-zero constant}).$$

in a forthcoming work joint with F. Castella, we will apply the above observation to prove a Rubin's formula for generalized Kato classes.

8.4. **An improved p -adic L -function.** Let

$$Z = (1 + T_1)^{-1}(1 + T_2)(1 + T_3) \in \mathcal{R}_0.$$

In this subsection, we introduce an ‘‘two-variable’’ improved p -adic L -function $\mathcal{L}_{\mathbf{F}}^* \in \mathcal{R}/(Z)$ for $\mathbf{F} = (\mathbf{f}, \mathbf{g}, \mathbf{h})$ a triple of primitive Hida families as in §3.5. We further assume that $\psi_1^{-1}\omega^{1+a}$ is unramified at p . Let $\alpha_p(?) := \mathbf{a}(p, ?)$ for $? \in \{\mathbf{f}, \mathbf{g}, \mathbf{h}\}$ be the \mathbf{U}_p -eigenvalues. Then we have

Proposition 8.3. *There exists an improved p -adic L -function $\mathcal{L}_{\mathbf{F}}^* \in \mathcal{R}/(Z)$ such that*

$$\mathcal{L}_{\mathbf{F}}^{\mathbf{f}}(\text{mod } Z) = \left(1 - \frac{\psi_1 \omega^{-a-1}(p) \alpha_p(\mathbf{g}) \alpha_p(\mathbf{h})}{\alpha_p(\mathbf{f})}\right) \cdot \mathcal{L}_{\mathbf{F}}^*.$$

Moreover, let $\underline{Q} = (Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathcal{R}}^{\mathbf{f}}$ with $Z(\underline{Q}) = 0$, then

$$(\mathcal{L}_{\mathbf{F}}^*(\underline{Q}))^2 = \frac{L(1/2, \Pi_{\underline{Q}})}{(-1)^{k_{Q_1}} \Omega_{\mathbf{f}_{Q_1}}^2} \cdot \mathcal{E}^*(\Pi_{\underline{Q}, p}),$$

where

$$\mathcal{E}^*(\Pi_{\underline{Q}, p}) := \frac{L_p(\text{Fil}_{\mathbf{f}}^+ \mathbf{V}_{\underline{Q}}^{\dagger}, 0)}{\varepsilon(\text{WD}_p(\text{Fil}_{\mathbf{f}}^+ \mathbf{V}_{\underline{Q}}^{\dagger}))} \cdot \frac{L_p(U'_{\underline{Q}}, s)^2}{L_p(\mathbf{V}_{\underline{Q}}^{\dagger} / \text{Fil}_{\mathbf{f}}^+ \mathbf{V}_{\underline{Q}}^{\dagger}, s)} \Big|_{s=0},$$

where $U'_{\underline{Q}} = (\text{Fil}^0 V_{\mathbf{f}_{Q_1}})^{\vee} \otimes \text{Fil}^0 V_{\mathbf{g}_{Q_2}} \otimes \text{Fil}^0 V_{\mathbf{h}_{Q_3}} \otimes \psi_1^{-1} \omega^{a+1}$.

PROOF. Let $\mathbf{G} := \mathbf{g}^* \cdot \mathbf{h}^* (\text{mod } Z)$. Then the argument in Lemma 3.4 shows that

$$\mathbf{G} \in \mathbf{S}(N, \psi_{1,(p)} \overline{\psi_1}^{(p)}, \mathbf{I}_1) \widehat{\otimes}_{\mathbf{I}_1} \mathcal{R}/(Z),$$

so we can define \mathbf{G}^{aux} as \mathbf{H}^{aux} in (3.8), replacing \mathbf{H} by \mathbf{G} and define $\mathcal{L}_{\mathbf{F}}^*$ by

$$\mathcal{L}_{\mathbf{F}}^* := \mathbf{a}(1, 1_{\mathbf{f}}^*(\text{Tr}_{N/N_1}(\mathbf{G}^{\text{aux}}))) \in \mathcal{R}/(Z).$$

In what follows, we shall keep the notation in §3.8. For each $\underline{Q} = (Q_1, Q_2, Q_3) \in \mathfrak{X}_{\mathcal{R}}^{\mathbf{f}}$ with $\mathcal{R}(\underline{Q}) = 0$, i.e. $k_{Q_1} = k_{Q_2} + k_{Q_3}$ and $\epsilon_{Q_1} = \epsilon_{Q_2} \epsilon_{Q_3}$, let $F = (f, g, h) = (\mathbf{f}_{Q_1}, \mathbf{g}_{Q_2}, \mathbf{h}_{Q_3})$. Applying the proof of Proposition 3.7 to the improved p -adic L -function $\mathcal{L}_{\mathbf{F}}^*$, we obtain

$$(8.2) \quad \frac{\mathcal{L}_{\mathbf{F}}^{\mathbf{f}}(\underline{Q})}{I(\rho(\mathbf{t}_n) \phi_{\mathbf{F}}^*)} = \frac{\mathcal{L}_{\mathbf{F}}^*(\underline{Q})}{I(\rho(\mathbf{t}_n) \phi_{\mathbf{F}}^{*,*})},$$

where $\phi_{\mathbf{F}}^{*,*} := \rho(\mathcal{J}_{\infty}) \varphi_{\mathbf{f}}^* \boxtimes \varphi_{\mathbf{g}}^* \boxtimes \varphi_{\mathbf{h}}^*$, and $I(\rho(\mathbf{t}_n) \phi_{\mathbf{F}}^{*,*})$ is the global trilinear period integral

$$I(\rho(\mathbf{t}_n) \phi_{\mathbf{F}}^{*,*}) = \int_{\mathbf{A} \times \text{GL}_2(\mathbf{Q}) \backslash \text{GL}_2(\mathbf{A})} \phi_{\mathbf{F}}^{*,*}(xt_n, x, x) d^{\tau} x.$$

Letting $\alpha_1 = \omega_{F,p}^{-1/2}(p) \mathbf{a}(p, f) p^{1 - \frac{k_{Q_1}}{2}}$, $\alpha_2 = \mathbf{a}(p, g) p^{1 - \frac{k_{Q_2}}{2}}$ and $\alpha_3 = \mathbf{a}(p, h) p^{1 - \frac{k_{Q_3}}{2}}$, one verifies that

$$\phi_{\mathbf{F}}^* = 1 \otimes 1 \otimes (1 - |p| \alpha_3 \cdot \pi_h \left(\begin{pmatrix} p^{-1} & 0 \\ 0 & 1 \end{pmatrix} \right)) \cdot \phi_{\mathbf{F}}^{*,*}$$

and that

$$I(\rho(\mathbf{t}_n) \phi_{\mathbf{F}}^*) = I(\rho(\mathbf{t}_n) \phi_{\mathbf{F}}^{*,*}) - |p|^{\frac{3}{2}} \alpha_1 \alpha_2 \alpha_3 \cdot I(\rho(\mathbf{t}_{n-1}) \phi_{\mathbf{F}}^{*,*})$$

for n sufficiently large. From the above equation, (8.2) and Proposition 3.7, we can deduce that

$$\mathcal{L}_{\mathbf{F}}^{\mathbf{f}}(\underline{Q}) = (1 - |p|^{\frac{1}{2}} \omega_{g,p} \omega_{h,p}(p) \alpha_1^{-1} \alpha_2 \alpha_3) \cdot \mathcal{L}_{\mathbf{F}}^*(\underline{Q}).$$

Now as in Theorem 7.1, we apply the above construction to a suitable Dirichlet twist \mathbf{F}' of \mathbf{F} and define $\mathcal{L}_{\mathbf{F}}^* := \mathcal{L}_{\mathbf{F}'}^* \cdot 2\sqrt{I_{\mathbf{F}'}}^{-1}$. Then $\mathcal{L}_{\mathbf{F}}^*$ clearly does the job.

To see the interpolation formula, applying the proof of Corollary 3.13 and Theorem 7.1 to $\mathcal{L}_{\mathbf{f}}^*$, we can show that

$$(\mathcal{L}_{\mathbf{F}}^*(\underline{Q}))^2 = \frac{1}{4} \cdot \frac{L(1/2, \Pi_{\underline{Q}})}{(-1)^{k_{Q_1}} \Omega_{\mathbf{f}_{Q_1}}^2} \cdot \mathcal{I}_{\Pi_{\underline{Q}, p}}^* \cdot \prod_{q|N} I_{\mathbf{F}, q}(\underline{Q}) \cdot \prod_{\ell \in \Sigma_{\text{exc}}} (1 + \ell^{-1}),$$

where $\mathcal{J}_{\Pi_{\mathbf{Q},p}}^*$ is the improved p -adic zeta integral defined in Remark 5.5. Then the interpolation formula follows from the expression of $\mathcal{J}_{\Pi_{\mathbf{Q},p}}^*$ given in Remark 5.5. \square

Remark 8.4. The method used here can be also applied to the construction of improved p -adic L -functions in the balanced case.

8.5. An alternative proof of anticyclotomic exceptional zero conjecture. We return to the setting in §8.2 and §8.3. Suppose that $f = f^\circ$ is the newform attached to an elliptic curve E/\mathbf{Q} of conductor Np with split multiplicative reduction at p . For a ring class character χ , put

$$\mathcal{L}_p(f/K \otimes \chi, s) := \mathcal{L}_{f/K, \chi}(\mathbf{v}^s - 1) \text{ for } s \in \mathbf{Z}_p.$$

Then we know $\mathcal{L}_p(f/K, 0) = 0$. Write $\mathfrak{p}^{h_K} = \varpi \mathcal{O}_K$ with $\varpi \in K^\times$ and let $\log_{\varpi/\overline{\varpi}}: \mathbf{C}_p^\times \rightarrow \mathbf{C}_p$ be the p -adic logarithm such that $\log_{\varpi/\overline{\varpi}}(\varpi/\overline{\varpi}) = 0$. We provide a Greenberg-Stevens style proof of the anticyclotomic exceptional zero conjecture for elliptic curves that was proved in [BD99].

Theorem 8.5 (Bertolini and Darmon). *Let q_E be the Tate period of E . Then we have*

$$\frac{d\mathcal{L}_p(f/K, s)}{ds} \Big|_{s=0} = \pm \frac{\log_{\varpi/\overline{\varpi}}(q_E)}{\text{ord}_p(q_E)} \cdot \sqrt{\frac{L(E/K, 1)u_K^2 D_K^{1/2}}{4\pi^2 \Omega_{f,N}}}.$$

PROOF. By [CH16, Theorem D], we can choose a ring class character χ of ℓ -power conductor with $\ell \nmid Np$ split in K such that $\mathcal{L}_p(f/K \otimes \chi^2, 0) \neq 0$. Let $\mathbf{f} = \mathbf{f}(T) \in \mathbf{Z}_p[[T]][[q]]$ be the primitive Hida family passing through f at the weight two specialization $T = \mathbf{u}^2 - 1$ with $\mathbf{u} := 1 + p$. Let $\mathbf{F} = (\mathbf{f}(T), \boldsymbol{\theta}_{\psi_\chi}(S_2), \boldsymbol{\theta}_{\psi_{\chi^{-1}}}(S_3))$ be the triple of Hida families and let $\mathcal{L}_{\mathbf{F}}^{\mathbf{f}} = \mathcal{L}_{\mathbf{F}}^{\mathbf{f}}(T, S_2, S_3)$ be the unbalanced p -adic L -function attached to \mathbf{F} in Theorem 7.1. Fixing a lift $\tilde{\mathcal{L}}_{\mathbf{F}}^* \in \mathcal{R}$ of $\mathcal{L}_{\mathbf{F}}^*$ (mod Z), we define analytic functions on \mathbf{Z}_p^3 :

$$\begin{aligned} \mathcal{L}_p(k_1, k_2, k_3) &:= \mathcal{L}_{\mathbf{F}}^{\mathbf{f}}(\mathbf{u}^{k_1} - 1, \mathbf{v}^{k_2} - 1, \mathbf{v}^{k_3} - 1); \\ \mathcal{L}_p^*(k_1, k_2, k_3) &:= \tilde{\mathcal{L}}_{\mathbf{F}}^*(\mathbf{u}^{k_1} - 1, \mathbf{v}^{k_2} - 1, \mathbf{v}^{k_3} - 1) \end{aligned}$$

for $(k_1, k_2, k_3) \in \mathbf{Z}_p^3$. Let $a_{\mathbf{f}}(k_1) = \alpha_p(\mathbf{f})(\mathbf{u}^{k_1} - 1)$,

$$a_{\mathbf{g}}(k_2) = \alpha_p(\mathbf{g})(\mathbf{v}^{k_2} - 1) = \chi(\text{Frob}_{\overline{\mathbf{p}}})\mathbf{v}^{l(\text{Frob}_{\overline{\mathbf{p}}})(1-k_2)}; \quad a_{\mathbf{h}}(k_3) = \chi^{-1}(\text{Frob}_{\overline{\mathbf{p}}})\mathbf{v}^{l(\text{Frob}_{\overline{\mathbf{p}}})(1-k_3)}.$$

It is clear that

$$a_{\mathbf{f}}(2) = 1; \quad a_{\mathbf{g}}(1)a_{\mathbf{g}}(1) = 1.$$

By Proposition 8.3, there exists $\mathcal{H}(T_1, S_1, S_2) \in \mathcal{R}$ and $H(k_1, k_2, k_3) = \mathcal{H}(\mathbf{u}^{k_1} - 1, \mathbf{v}^{k_2} - 1, \mathbf{v}^{k_3} - 1)$ such that

$$(8.3) \quad \mathcal{L}_p(k_1, k_2, k_3) = \left(1 - \frac{a_{\mathbf{g}}(k_2)a_{\mathbf{h}}(k_3)}{a_{\mathbf{f}}(k_1)}\right) \cdot \mathcal{L}_p^*(k_1, k_2, k_3) + H(k_1, k_2, k_3) \cdot (\mathbf{u}^{-k_1+k_2+k_3} - 1)$$

(the nebentypus $\psi_1 = 1, \psi_2 = \psi_3 = \omega^{-1}$ and $a = -1$). We may assume $L(f/K, 1) \neq 0$, so the root numbers of f and its quadratic twist $f \otimes \tau_{K/\mathbf{Q}}$ are $+1$. This in turns implies that the root numbers of \mathbf{f} and $\mathbf{f} \otimes \tau_{K/\mathbf{Q}}$ are -1 , and hence the one-variable Iwasawa function $\mathcal{L}_p(k_1, 1, 1)$ vanishes identically. Taking the derivative with respect to k_1 on the both sides of (8.3), we find that

$$0 = \frac{\partial \mathcal{L}_p}{\partial k_1}(2, 1, 1) = a'_{\mathbf{f}}(2) \cdot \mathcal{L}_p^*(2, 1, 1) - H(2, 1, 1) \cdot \log_p \mathbf{u}.$$

This implies that

$$H(2, 1, 1) \cdot \log_p \mathbf{u} = a'_{\mathbf{f}}(2) \cdot \mathcal{L}_p^*(2, 1, 1);$$

By an elementary calculation and a theorem of Greenberg-Stevens [GS93, Theorem 3.18],

$$a'_{\mathbf{g}}(1) = \frac{\log_p \overline{\varpi}}{h_K}; \quad a'_{\mathbf{f}}(2) = -\frac{1}{2} \cdot \frac{\log_p(q_E)}{\text{ord}_p(q_E)}.$$

It follows that

$$\begin{aligned} \frac{\partial \mathcal{L}_p}{\partial k_2}(2, 1, 1) &= \frac{\partial \mathcal{L}_p}{\partial k_3}(2, 1, 1) = \left(-\frac{\log_p \overline{\varpi}}{h_K}\right) \cdot \mathcal{L}_p^*(2, 1, 1) + H(2, 1, 1) \cdot \log_p \mathbf{u} \\ &= \left(-\frac{\log_p \overline{\varpi}}{h_K} - \frac{1}{2} \frac{\log_p(q_E)}{\text{ord}_p(q_E)}\right) \cdot \mathcal{L}_p^*(2, 1, 1). \end{aligned}$$

By Proposition 8.1, we have

$$\mathcal{L}_p(2, k_2, k_3) = v(k_2, k_3) \cdot \mathcal{L}_p(f/K, \frac{k_2 + k_3 - 2}{2}) \mathcal{L}_p(f \otimes \chi^2, \frac{k_2 - k_3}{2})$$

for some nowhere vanishing analytic function $v(k_2, k_3)$. Letting $v = v(1, 1) \neq 0$, we find that

$$\begin{aligned} v \cdot \mathcal{L}'_p(f/K, 0) \mathcal{L}_p(f/K \otimes \chi^2, 0) &= \frac{\partial \mathcal{L}_p}{\partial k_2}(2, 1, 1) + \frac{\partial \mathcal{L}_p}{\partial k_3}(2, 1, 1) \\ &= (-1) \left(\frac{\log_p(q_E)}{\text{ord}_p(q_E)} + \frac{2 \log_p \overline{\omega}}{h_K} \right) \cdot \mathcal{L}_p^*(2, 1, 1) \\ &= (-1) \frac{\log_{\overline{\omega}/\overline{\omega}}(q_E)}{\text{ord}_p(q_E)} \cdot \mathcal{L}_p^*(2, 1, 1). \end{aligned}$$

On the other hand, the interpolation formula in Proposition 8.3 shows that

$$\mathcal{L}_p^*(2, 1, 1)^2 = v^2 \cdot (2\pi)^{-2} \frac{L(f/K, 1)}{\Omega_{f, N^-}} \cdot u_K^2 \sqrt{D_K} \cdot \mathcal{L}_p(f/K \otimes \chi^2, 0)^2.$$

Combining the above two equations, we obtain

$$(\mathcal{L}'_p(f/K, 0))^2 = \left(\frac{\log_{\overline{\omega}/\overline{\omega}}(q_E)}{\text{ord}_p(q_E)} \right)^2 \cdot \frac{L(f/K, 1)}{4\pi^2 \Omega_{f, N^-}} \cdot u_K^2 \sqrt{D_K},$$

and the theorem follows. \square

REFERENCES

- [BD96] M. Bertolini and H. Darmon, *Heegner points on Mumford-Tate curves*, Invent. Math. **126** (1996), no. 3, 413–456.
- [BD99] Massimo Bertolini and Henri Darmon, *p -adic periods, p -adic L -functions, and the p -adic uniformization of Shimura curves*, Duke Math. J. **98** (1999), no. 2, 305–334.
- [BD07] ———, *Hida families and rational points on elliptic curves*, Invent. Math. **168** (2007), no. 2, 371–431. MR 2289868
- [BDP13] M. Bertolini, H. Darmon, and K. Prasanna, *Generalized Heegner cycles and p -adic Rankin L -series*, Duke Math. J. **162** (2013), no. 6, 1033–1148.
- [Bum97] D. Bump, *Automorphic forms and representations*, Cambridge Studies in Advanced Mathematics, vol. 55, Cambridge University Press, Cambridge, 1997.
- [Cas73] W. Casselman, *On some results of Atkin and Lehner*, Math. Ann. **201** (1973), 301–314.
- [CC17] S.-Y. Chen and Yao Cheng, *On Deligne’s conjecture for central values of certain automorphic representation on $\text{GL}(3) \times \text{GL}(2)$* , preprint.
- [CH16] Masataka Chida and Ming-Lun Hsieh, *Special values of anticyclotomic L -functions for modular forms*, J. reine angew. Math. (2016), to appear. Published Online: DOI 10.1515/crelle-2015-0072.
- [Coa89a] John Coates, *On p -adic L -functions*, Astérisque (1989), no. 177-178, Exp. No. 701, 33–59, Séminaire Bourbaki, Vol. 1988/89. MR 1040567
- [Coa89b] ———, *On p -adic L -functions attached to motives over \mathbf{Q} II*, Boletim da Sociedade Brasileira de Matemática - Bulletin/Brazilian Mathematical Society **20** (1989), no. 1, 101–112.
- [Col16] Dan Collins, *Anticyclotomic p -adic L -functions and Ichino’s formula*, preprint. arXiv:1612.06948.
- [CPR89] John Coates and Bernadette Perrin-Riou, *On p -adic L -functions attached to motives over \mathbf{Q}* , Algebraic number theory, Adv. Stud. Pure Math., vol. 17, Academic Press, Boston, MA, 1989, pp. 23–54.
- [DDT94] Henri Darmon, Fred Diamond, and Richard Taylor, *Fermat’s last theorem*, Current developments in mathematics, 1995 (Cambridge, MA), Int. Press, Cambridge, MA, 1994, pp. 1–154.
- [Del79] P. Deligne, *Valeurs de fonctions L et périodes d’intégrales*, Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979, With an appendix by N. Koblitz and A. Ogus, pp. 313–346.
- [Dim14] Mladen Dimitrov, *On the local structure of ordinary Hecke algebras at classical weight one points*, Automorphic forms and Galois representations. Vol. 2, London Math. Soc. Lecture Note Ser., vol. 415, Cambridge Univ. Press, Cambridge, 2014, pp. 1–16.
- [DLR15] Henri Darmon, Alan Lauder, and Victor Rotger, *Stark points and p -adic iterated integrals attached to modular forms of weight one*, Forum Math. Pi **3** (2015), e8, 95. MR 3456180
- [DR14] Henri Darmon and Victor Rotger, *Diagonal cycles and Euler systems I: A p -adic Gross-Zagier formula*, Ann. Sci. Éc. Norm. Supér. (4) **47** (2014), no. 4, 779–832.
- [DR17] ———, *Diagonal cycles and Euler systems II: The Birch and Swinnerton-Dyer conjecture for Hasse-Weil-Artin L -functions*, J. Amer. Math. Soc. **30** (2017), no. 3, 601–672.
- [DT94] F. Diamond and R. Taylor, *Nonoptimal levels of mod l modular representations*, Invent. Math. **115** (1994), no. 3, 435–462.

- [FO12] Olivier Fouquet and Tadashi Ochiai, *Control theorems for Selmer groups of nearly ordinary deformations*, J. Reine Angew. Math. **666** (2012), 163–187. MR 2920885
- [Fon94] Jean-Marc Fontaine, *Représentations l -adiques potentiellement semi-stables*, Astérisque (1994), no. 223, 321–347, Périodes p -adiques (Bures-sur-Yvette, 1988). MR 1293977
- [GJ78] Stephen Gelbart and Hervé Jacquet, *A relation between automorphic representations of $GL(2)$ and $GL(3)$* , Ann. Sci. École Norm. Sup. (4) **11** (1978), no. 4, 471–542. MR 533066
- [Gre94] Ralph Greenberg, *Iwasawa theory and p -adic deformations of motives*, Motives (Seattle, WA, 1991), Proc. Sympos. Pure Math., vol. 55, Amer. Math. Soc., Providence, RI, 1994, pp. 193–223. MR 1265554
- [GS93] Ralph Greenberg and Glenn Stevens, *p -adic L -functions and p -adic periods of modular forms*, Invent. Math. **111** (1993), no. 2, 407–447.
- [GS16] Matthew Greenberg and Marco Adamo Seveso, *Triple product p -adic L -functions for balanced weights*, arXiv:1506.05681.
- [Hel07] David Helm, *On maps between modular Jacobians and Jacobians of Shimura curves*, Israel J. Math. **160** (2007), 61–117.
- [Hid85] Haruzo Hida, *A p -adic measure attached to the zeta functions associated with two elliptic modular forms. I*, Invent. Math. **79** (1985), no. 1, 159–195. MR 774534
- [Hid88a] H. Hida, *Modules of congruence of Hecke algebras and L -functions associated with cusp forms*, Amer. J. Math. **110** (1988), no. 2, 323–382. MR 935010 (89i:11058)
- [Hid88b] ———, *On p -adic hecke algebras for $GL(2)$ over totally real fields*, Annals of Mathematics **128** (1988), 295–384.
- [Hid88c] Haruzo Hida, *A p -adic measure attached to the zeta functions associated with two elliptic modular forms. II*, Ann. Inst. Fourier (Grenoble) **38** (1988), no. 3, 1–83.
- [Hid90] ———, *p -adic L -functions for base change lifts of GL_2 to GL_3* , Automorphic forms, Shimura varieties, and L -functions, Vol. II (Ann Arbor, MI, 1988), Perspect. Math., vol. 11, Academic Press, Boston, MA, 1990, pp. 93–142.
- [Hid93] ———, *Elementary theory of L -functions and Eisenstein series*, London Mathematical Society Student Texts, vol. 26, Cambridge University Press, Cambridge, 1993.
- [Hid94] ———, *On the critical values of L -functions of $GL(2)$ and $GL(2) \times GL(2)$* , Duke Math. J. **74** (1994), no. 2, 431–529.
- [Hid06] H. Hida, *Hilbert modular forms and Iwasawa theory*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, Oxford, 2006.
- [Hid16] Haruzo Hida, *Arithmetic of adjoint L -values, p -adic aspects of modular forms*, World Sci. Publ., Hackensack, NJ, 2016, pp. 185–236.
- [HK91] Michael Harris and Stephen S. Kudla, *The central critical value of a triple product L -function*, Ann. of Math. (2) **133** (1991), no. 3, 605–672. MR 1109355
- [HT01a] M. Harris and R. Taylor, *The geometry and cohomology of some simple Shimura varieties*, Annals of Mathematics Studies, vol. 151, Princeton University Press, Princeton, NJ, 2001, With an appendix by Vladimir G. Berkovich.
- [HT01b] Michael Harris and Jacques Tilouine, *p -adic measures and square roots of special values of triple product L -functions*, Math. Ann. **320** (2001), no. 1, 127–147. MR 1835065
- [Hu17] YueKe Hu, *The subconvexity bound for triple product L -functions in level aspect*, Amer. J. Math. **139** (2017), no. 1, 215–259.
- [Hun17] Pin-Chi Hung, *On the non-vanishing mod l of central L -values with anticyclotomic twists for Hilbert modular forms*, J. Number Theory **173** (2017), 170–209. MR 3581914
- [Ich08] Atsushi Ichino, *Trilinear forms and the central values of triple product L -functions*, Duke Math. J. **145** (2008), no. 2, 281–307.
- [II10] Atsushi Ichino and Tamutsu Ikeda, *On the periods of automorphic forms on special orthogonal groups and the Gross-Prasad conjecture*, Geom. Funct. Anal. **19** (2010), no. 5, 1378–1425.
- [Ish17] Isao Ishikawa, *On the construction of twisted triple product p -adic L -functions*, 2017, Thesis (Ph.D.)—Kyoto University.
- [Jac72] H. Jacquet, *Automorphic forms on $GL(2)$. Part II*, Lecture Notes in Mathematics, Vol. 278, Springer-Verlag, Berlin, 1972.
- [JL70] H. Jacquet and R. P. Langlands, *Automorphic forms on $GL(2)$* , Lecture Notes in Mathematics, Vol. 114, Springer-Verlag, Berlin, 1970.
- [LV11] Matteo Longo and Stefano Vigni, *Quaternion algebras, Heegner points and the arithmetic of Hida families*, Manuscripta Math. **135** (2011), no. 3-4, 273–328.
- [Miy06] Toshitsune Miyake, *Modular forms*, english ed., Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2006, Translated from the 1976 Japanese original by Yoshitaka Maeda. MR 2194815
- [MV10] Philippe Michel and Akshay Venkatesh, *The subconvexity problem for GL_2* , Publ. Math. Inst. Hautes Études Sci. (2010), no. 111, 171–271.
- [MW86] B. Mazur and A. Wiles, *On p -adic analytic families of Galois representations*, Compositio Math. **59** (1986), no. 2, 231–264.
- [NPS14] Paul D. Nelson, Ameya Pitale, and Abhishek Saha, *Bounds for Rankin-Selberg integrals and quantum unique ergodicity for powerful levels*, J. Amer. Math. Soc. **27** (2014), no. 1, 147–191.
- [Oht95] Masami Ohta, *On the p -adic Eichler-Shimura isomorphism for Λ -adic cusp forms*, J. Reine Angew. Math. **463** (1995), 49–98. MR 1332907
- [Oht00] ———, *Ordinary p -adic étale cohomology groups attached to towers of elliptic modular curves. II*, Math. Ann. **318** (2000), no. 3, 557–583. MR 1800769

- [Orl87] Tobias Orloff, *Special values and mixed weight triple products (with an appendix by Don Blasius)*, Invent. Math. **90** (1987), no. 1, 169–188.
- [Pra90] Dipendra Prasad, *Tilinear forms for representations of $GL(2)$ and local ϵ -factors*, Compositio Math. **75** (1990), no. 1, 1–46.
- [PSR87] I. Piatetski-Shapiro and Stephen Rallis, *Rankin triple L functions*, Compositio Math. **64** (1987), no. 1, 31–115. MR 911357
- [PW11] R. Pollack and T. Weston, *On anticyclotomic μ -invariants of modular forms*, Compos. Math. **147** (2011), no. 5, 1353–1381.
- [Ram00] Dinakar Ramakrishnan, *Modularity of the Rankin-Selberg L -series, and multiplicity one for $SL(2)$* , Ann. of Math. (2) **152** (2000), no. 1, 45–111. MR 1792292
- [Rib84] Kenneth A. Ribet, *Congruence relations between modular forms*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Warsaw, 1983), PWN, Warsaw, 1984, pp. 503–514. MR 804706
- [Sch02] Ralf Schmidt, *Some remarks on local newforms for $GL(2)$* , J. Ramanujan Math. Soc. **17** (2002), no. 2, 115–147.
- [SU06] Christopher Skinner and Eric Urban, *Sur les déformations p -adiques de certaines représentations automorphes*, J. Inst. Math. Jussieu **5** (2006), no. 4, 629–698.
- [Tat79] J. Tate, *Number theoretic background*, Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979, pp. 3–26. MR 546607
- [Tay04] Richard Taylor, *Galois representations*, Ann. Fac. Sci. Toulouse Math. (6) **13** (2004), no. 1, 73–119. MR 2060030
- [Wal85] J.-L. Waldspurger, *Sur les valeurs de certaines fonctions L automorphes en leur centre de symétrie*, Compositio Math. **54** (1985), no. 2, 173–242.
- [Wil88] A. Wiles, *On ordinary λ -adic representations associated to modular forms*, Invent. Math. **94** (1988), no. 3, 529–573.
- [Wil95] ———, *Modular elliptic curves and Fermat’s last theorem*, Ann. of Math. (2) **141** (1995), no. 3, 443–551.

INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, TAIPEI 10617, TAIWAN
E-mail address: mlhsieh@math.sinica.edu.tw